

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
26 July 2001 (26.07.2001)

PCT

(10) International Publication Number  
WO 01/053456 A3(51) International Patent Classification<sup>7</sup>: C07H 21/02,  
21/04, C07K 14/00, C12N 1/21, 9/00, 15/00, 15/63, 15/85,  
15/8609/491,404  
09/552,317  
09/596,196  
09/643,31325 January 2000 (25.01.2000) US  
25 April 2000 (25.04.2000) US  
17 June 2000 (17.06.2000) US  
31 August 2000 (31.08.2000) US

(21) International Application Number: PCT/US00/35061

(22) International Filing Date:

22 December 2000 (22.12.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

09/471,275 23 December 1999 (23.12.1999) US  
09/488,725 21 January 2000 (21.01.2000) US(63) Related by continuation (CON) or continuation-in-part  
(CIP) to earlier applications:

US	09/488,725 (CIP)
Filed on	21 January 2000 (21.01.2000)
US	09/491,404 (CIP)
Filed on	25 January 2000 (25.01.2000)
US	09/552,317 (CIP)
Filed on	25 April 2000 (25.04.2000)
US	09/596,196 (CIP)
Filed on	17 June 2000 (17.06.2000)
US	09/653,274 (CIP)
Filed on	31 August 2000 (31.08.2000)

[Continued on next page]

(54) Title: METHODS AND MATERIALS RELATING TO PROTHROMBINASE-LIKE POLYPEPTIDES AND POLYNUCLEOTIDES

FROM FIG. 1-1

PROTHROMBINASE-LIKE: 156 TIGSVTKTPSGLYIIHPEGSSYPFEVMDYRGGGWTVIQKRIDGIIIDFQRLWCDYLG 215  
IG K S Y + P+ + FEV CDW+ GGGWTV+Q R+DG +F R W DY G  
PROTHROMBINASE FG12: 218 AIG---KRSSETYRVTPDPKNSSFEVYCDWETMGGGWTVLQARLDGSTNFTRTWQDYKAG 274

PROTHROMBINASE-LIKE: 216 FGDLLGEFWLGLKKIFYIVNQKNTSFMLYVALESEDDTLAYASYDNFLEDETRFFKMH 275  
FG+L EFWLG KI + K +L+LE + YA YD F++ +E +++H+  
PROTHROMBINASE FG12: 275 FGNLRREFWLGNDKIHLTKSKE---WILRIDLEDFNGVELYALDYQFYVANEFLKYRLHV 332

PROTHROMBINASE-LIKE: 276 GRYSGNAGDAFRGLKKEDNQAMPFSTSDVDNDGCRPAVLNGOSVKSCHLHNKTGWFF 335  
G Y+G AGDA R K N+ F+T D DND P+ G +C L+ +GWFF  
PROTHROMBINASE FG12: 333 GNYNGTAGDALP-FNKHYNHOLKFFFTPDKNDRY-PS---G---NCG-LYSSGWFF 381

PROTHROMBINASE-LIKE: 336 NECGLANLNG-IHHFSGKLLATG!QWGTW---TKNNSPVKIKSVSMKIRRMYPN-YFK 383  
+ C ANLNG +H + + G! WGTW P KS + + M P +FK  
PROTHROMBINASE FG12: 382 DACLSANLNGKYYHOKYRGVRNG!FWGTWPGVSEAHPPGGYKSSFKAKMMIRPKHFK 438

BLASTX ALIGNMENT OF SEQ ID NO: 4 (PROTHROMBINASE-LIKE WITH HUMAN PROTHROMBINASE FG 12 PROTEIN  
(PROTHROMBINASE Fg1))The Genbank database indicates that the fibrinogen-like protein  
is homologous (37% identity)

```
>gi |5730075 ref |NP_006673.1||fibrinogen-like 2 precursor; fibroleukin
>pir || S47273 pT49 protein-human >pir ||I37391 fibrinogen-like
protein expressed in T lymphocytes (pT49) - human
>emb |CAA85298.1 | (Z36531) fibrinogen-like protein [Homo sapiens]
>gb |AAD10825.1 | (AF104015) fibrinogen-like protein 2 [Homo sapeins]
Length = 439
```

Score = 402 (141.5 bits), Expect = 5.7e-36, P = 5.7e-36  
Identities = 109/291 (37%), Positives = 155/291 (53%)

PROTHROMBINASE-LIKE: 98 L RNMDEQQASLDYLSNQVNELMNRVLLTTEVFRKQLOPPHPRVQSHGL---DCTDIKD 155  
+ N +D + A+L ++ VN L + ++ +Q+ RPVQ H + DC+D  
PROTHROMBINASE FG12: 169 IENYDYSKVANLTFV---VNSLOGKCKSCPSQ---EQIGS---RPVQ-HLIVKDCSDYY- 217

TO FIG. 1-2

WO 01/053456 A3



(71) **Applicant** (*for all designated States except US*): **HYSEQ, INC.** [US/US]; 670 Almanor Avenue, Sunnyvale, CA 94086 (US).

LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(72) **Inventors; and**

(75) **Inventors/Applicants** (*for US only*): **HALEY, Dana, A.** [US/US]; 1430 Glenwood Avenue, San Jose, CA 95125 (US). **BOYLE, Bryan, J.** [US/US]; 1947 10th Avenue, San Francisco, CA 94116 (US). **SUK-YUE HO, Alice** [US/US]; 30573 Meridian Circle, Union City, CA 94587 (US). **ZHOU, Ping** [CN/US]; 1461 Japaul Lane, San Jose, CA 95132 (US). **MIZE, Nancy, K.** [US/US]; 662 Mountain View Avenue, Mountain View, CA 94041 (US). **KUO, Chiauyun** [US/US]; 2121 Limewood Drive, San Jose, CA 95132 (US). **ARTERBURN, Matthew, C.** [US/US]; 5910 Via Del Cielo, Pleasanton, CA 94566 (US). **TANG, Y., Tom** [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). **LIU, Chenghua** [CN/US]; 1125 Ranchero Way #14, San Jose, CA 95117 (US). **DRMANAC, Radoje** [YU/US]; 850 East Greenwich Place, Palo Alto, CA 94303 (US).

(84) **Designated States** (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

— *with international search report*

(88) **Date of publication of the international search report:**  
30 May 2002

(48) **Date of publication of this corrected version:**  
5 December 2002

(74) **Agent:** **ELRIFI, Ivor, R.**; Mintz, Levin, Cohn, Ferris, Glovsky and Popeo, P.C., One Financial Center, Boston, MA 02111 (US).

(15) **Information about Corrections:**

see PCT Gazette No. 49/2002 of 5 December 2002, Section II

**Previous Correction:**

see PCT Gazette No. 27/2002 of 4 July 2002, Section II

(81) **Designated States** (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

## METHODS AND MATERIALS RELATING TO PROTHROMBINASE-LIKE POLYPEPTIDES AND POLYNUCLEOTIDES

### 1. TECHNICAL FIELD

5           The present invention provides novel polynucleotides and proteins encoded by such polynucleotides, along with uses for these polynucleotides and proteins, for example in therapeutic, diagnostic and research methods. In particular, the invention relates to a novel prothrombinase-like polypeptide.

### 10   2. BACKGROUND ART

          Identified polynucleotide and polypeptide sequences have numerous applications in, for example, diagnostics, forensics, gene mapping; identification of mutations responsible for genetic disorders or other traits, to assess biodiversity, and to produce many other types of data and products dependent on DNA and amino acid sequences. Proteins are known to have  
15   biological activity, for example, by virtue of their secreted nature in the case of leader sequence cloning, by virtue of their cell or tissue source in the case of PCR-based techniques, or by virtue of structural similarity to other genes of known biological activity. It is to these polypeptides and the polynucleotides encoding them that the present invention is directed. In particular, this invention is directed to novel soluble prothrombinase-like polypeptides and  
20   polynucleotides.

          The ability of the body to control the flow of blood following vascular injury is paramount to continued survival. The process of blood clotting and then the subsequent dissolution of the clot, following repair of the injured tissue, is termed hemostasis. Hemostasis consists of 1) vascular constriction, 2) platelet activation and clumping, 3) formation of  
25   thrombus, and 4) clot dissolution following tissue repair. In order to initiate the clotting process, inactive prothrombin is converted to thrombin by prothrombinase complex, which is composed of platelet phospholipids, Ca ions, prothrombin and factors Va and Xa. Prothrombinase has been identified and characterized as an immune procoagulant (International Patent Application No. WO98/51335). The generation of thrombin by cleavage of  
30   prothrombin is a central regulatory process. Thrombin activates platelets to start the initial cascade and also cleaves fibrinogen, a major plasma protein. Fibrin forms a network of fibers, which finally leads to formation of a clot. Thus, prothrombinase is a procoagulant and regulation of its activity is crucial for survival. Fibrinogen cleavage products are also shown

to display vasoconstrictor and chemotactic activities and are mitogenic for several cell types (Herrick et al, (1999) *Int. J. Biochem. Cell Biol.* 31, 741-746).

Dysregulation of coagulation pathways can result in severe pathological conditions including hemophilia, myocardial infarction, glomerular diseases, diabetes, tumor growth, 5 fulminant viral hepatitis, and atherosclerosis among others. Many of these pathologies involve fibrin deposition and necrosis of the affected tissue. Thrombosis or blood clotting has also been associated with cancer (Dovorak, H.F. *Hum. Pathol.* 18(3), pp. 275-284 (1987)). Prothrombinase is also associated with viral hepatitis. (Levy et al., *Am. J. Pathol.* 156(4), pp. 1217-1225 (2000)). Some researchers have shown that hyperexpression of prothrombinase is 10 associated with early stages of Type 1 diabetes (van der Planken et al., *Clin. Appl. Thromb. Hemost.* 6(2), pp. 65-68 (2000)). Prothrombinase or prothrombinase-like protein may be beneficial in wound healing by increasing coagulation (Thiagarajan et al., *Biochemistry* 39(21) pp. 6459-6465 (2000)). Inhibition of prothrombinase or prothrombinase-like protein may reduce abnormal clotting, increase pathogenesis or viral hepatitis, decrease liver necrosis, or 15 treat diabetes Type 1.

Recently, it is becoming apparent that there are inducible prothrombinase genes that are induced either upon viral infection or cytokine treatment (Levy et al (2000) *Am. J. Pathol.* 150, 1217-1225). Antibodies to the prothrombinase fgl2 protein were able to abrogate the fulminant hepatitis in murine model of the disease and reduce the liver necrosis. Similarly, it 20 was shown that fgl2 expression is up regulated upon cytokine treatment of pregnant mice leading to spontaneous abortion (Clark et al (1999) *Am. J. Reproductive Immunol.* 42, 37-43). It would be beneficial to identify other inducible prothrombinase-like proteins that are expressed only under certain pathological conditions.

25

### 3. SUMMARY OF THE INVENTION

This invention is based on the discovery of novel prothrombinase-like polypeptides, novel isolated polynucleotides encoding such polypeptides, including recombinant DNA molecules, cloned genes or degenerate variants thereof, especially naturally occurring variants 30 such as allelic variants, antisense polynucleotide molecules, and antibodies that specifically recognize one or more epitopes present on such polypeptides, as well as hybridomas producing such antibodies. Specifically, the polynucleotides of the present invention are based on a prothrombinase-like polynucleotide isolated from a cDNA library prepared from human ovary



(Hyseq clone identification numbers 7735656 (SEQ ID NO: 1) and 7727065 (SEQ ID NO: 11)).

The compositions of the present invention additionally include vectors such as expression vectors containing the polynucleotides of the invention, cells genetically engineered to contain such polynucleotides and cells genetically engineered to express such polynucleotides.

The compositions of the invention provide isolated polynucleotides that include, but are not limited to, a polynucleotide comprising the nucleotide sequence set forth in the SEQ ID NO: 1-3, 10-13, and 15; or a fragment of SEQ ID NO: 1-3, 10-13 and 15; a polynucleotide comprising the full length protein coding sequence of the SEQ ID NO: 1-3, 10-13 and 15 (for example, SEQ ID NO: 4, 9, 14, 18); and a polynucleotide comprising the nucleotide sequence of the mature protein coding sequence of any of SEQ ID NO:1-3, 10-13 and 15. The polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes under stringent hybridization conditions to (a) the complement of any of the nucleotide sequences set forth in SEQ ID NO:1-3, 10-13 and 15; (b) a nucleotide sequence encoding any of SEQ ID NO: 4-9, 14, and 16-18; a polynucleotide which is an allelic variant of any polynucleotides recited above having at least 70% polynucleotide sequence identity to the polynucleotides; a polynucleotide which encodes a species homolog (e.g. orthologs) of any of the peptides recited above; or a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptide comprising SEQ ID NO: 4-9, 14 and 16-18.

A collection as used in this application can be a collection of only one polynucleotide. The collection of sequence information or unique identifying information of each sequence can be provided on a nucleic acid array. In one embodiment, segments of sequence information are provided on a nucleic acid array to detect the polynucleotide that contains the segment. The array can be designed to detect full-match or mismatch to the polynucleotide that contains the segment. The collection can also be provided in a computer-readable format.

This invention further provides cloning or expression vectors comprising at least a fragment of the polynucleotides set forth above and host cells or organisms transformed with these expression vectors. Useful vectors include plasmids, cosmids, lambda phage derivatives, phagemids, and the like, that are well known in the art. Accordingly, the invention also provides a vector including a polynucleotide of the invention and a host cell containing the polynucleotide. In general, the vector contains an origin of replication functional in at least one organism, convenient restriction endonuclease sites, and a selectable marker for the host cell. Vectors according to the invention include expression vectors, replication vectors, probe generation

vectors, and sequencing vectors. A host cell according to the invention can be a prokaryotic or eukaryotic cell and can be a unicellular organism or part of a multicellular organism.

The compositions of the present invention include polypeptides comprising, but not limited to, an isolated polypeptide selected from the group comprising the amino acid sequence of SEQ ID NO: 4-9, 14, and 16-18; or the corresponding full length or mature protein. Polypeptides of the invention also include polypeptides with biological activity that are encoded by (a) any of the polynucleotides having a nucleotide sequence set forth in the SEQ ID NO: 1-3, 10-13 and 15; or (b) polynucleotides that hybridize to the complement of the polynucleotides of (a) under stringent hybridization conditions. Biologically or immunologically active variants of any of the protein sequences listed as SEQ ID NO: 4-9, 14 and 16-18 and substantial equivalents thereof that retain biological or immunological activity are also contemplated. The polypeptides of the invention may be wholly or partially chemically synthesized but are preferably produced by recombinant means using the genetically engineered cells (e.g. host cells) of the invention.

The invention also provides compositions comprising a polypeptide of the invention. Pharmaceutical compositions of the invention may comprise a polypeptide of the invention and an acceptable carrier, such as a hydrophilic, e.g., pharmaceutically acceptable, carrier.

The invention also relates to methods for producing a polypeptide of the invention comprising culturing host cells comprising an expression vector containing at least a fragment of a polynucleotide encoding the polypeptide of the invention in a suitable culture medium under conditions permitting expression of the desired polypeptide, and purifying the protein or peptide from the culture or from the host cells. Preferred embodiments include those in which the protein produced by such a process is a mature form of the protein.

Polynucleotides according to the invention have numerous applications in a variety of techniques known to those skilled in the art of molecular biology. These techniques include use as hybridization probes, use as oligomers, or primers, for PCR, use in an array, use in computer-readable media, use for chromosome and gene mapping, use in the recombinant production of protein, and use in generation of antisense DNA or RNA, their chemical analogs and the like. For example, when the expression of an mRNA is largely restricted to a particular cell or tissue type, polynucleotides of the invention can be used as hybridization probes to detect the presence of the particular cell or tissue mRNA in a sample using, *e.g.*, *in situ* hybridization.

In other exemplary embodiments, the polynucleotides are used in diagnostics as expressed sequence tags for identifying expressed genes or, as well known in the art and

exemplified by Vollrath et al., Science 258:52-59 (1992), as expressed sequence tags for physical mapping of the human genome.

The polypeptides according to the invention can be used in a variety of conventional procedures and methods that are currently applied to other proteins. For example, a  
5 polypeptide of the invention can be used to generate an antibody that specifically binds the polypeptide. Such antibodies, particularly monoclonal antibodies, are useful for detecting or quantitating the polypeptide in tissue. The polypeptides of the invention can also be used as molecular weight markers, and as a food supplement.

Methods are also provided for preventing, treating, or ameliorating a medical condition  
10 which comprises the step of administering to a mammalian subject a therapeutically effective amount of a composition comprising a peptide of the present invention and a pharmaceutically acceptable carrier.

In particular, the polypeptides and polynucleotides of the invention can be utilized, for example, as part of methods for the prevention and/or treatment of diseases associated with the  
15 dysregulation of coagulation pathways, such as hemophilia, myocardial infarction, glomerular disease, diabetes, fulminant viral hepatitis and atherosclerosis among others. The polynucleotides and polypeptides may also be beneficial in the treatment of viral infections and in the treatment of some forms of cancer.

The methods of the invention also provides methods for the treatment of disorders as  
20 recited herein which comprise the administration of a therapeutically effective amount of a composition comprising a polynucleotide or polypeptide of the invention and a pharmaceutically acceptable carrier to a mammalian subject exhibiting symptoms or tendencies related to disorders as recited herein. In addition, the invention encompasses methods for treating diseases or disorders as recited herein comprising the step of administering a  
25 composition comprising compounds and other substances that modulate the overall activity of the target gene products and a pharmaceutically acceptable carrier. Compounds and other substances can effect such modulation either on the level of target gene/protein expression or target protein activity. Specifically, methods are provided for preventing, treating or ameliorating a medical condition, including viral diseases, which comprises administering to a  
30 mammalian subject, including but not limited to humans, a therapeutically effective amount of a composition comprising a polypeptide of the invention, a polynucleotide of the invention or a therapeutically effective amount of a composition comprising a binding partner of (e.g., antibody specifically reactive for) prothrombinase-like polypeptides of the invention. The mechanics of the particular condition or pathology will dictate whether the polypeptides, or

nucleotides of the invention or binding partners (or inhibitors) of the polypeptides would be beneficial to the individual in need of treatment.

According to this method, polypeptides of the invention can be administered to produce an *in vitro* or *in vivo* inhibition of cellular function. A polypeptide of the invention can be  
5 administered *in vivo* alone or as an adjunct to other therapies. Conversely, protein or other active ingredients of the present invention may be included in formulations of a particular agent to minimize side effects of such an agent.

The invention further provides methods for manufacturing medicaments useful in the above-described methods.

10 The present invention further relates to methods for detecting the presence of the polynucleotides or polypeptides of the invention in a sample (e.g., tissue or sample). Such methods can, for example, be utilized as part of prognostic and diagnostic evaluation of disorders as recited herein and for the identification of subjects exhibiting a predisposition to such conditions.

15 The invention provides a method for detecting a polypeptide of the invention in a sample comprising contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex and detecting formation of the complex, so that if a complex is formed, the polypeptide is detected.

The invention also provides kits comprising polynucleotide probes and/or monoclonal  
20 antibodies, and optionally quantitative standards, for carrying out methods of the invention. Furthermore, the invention provides methods for evaluating the efficacy of drugs, and monitoring the progress of patients, involved in clinical trials for the treatment of disorders as recited above.

The invention also provides methods for the identification of compounds that modulate  
25 (i.e., increase or decrease) the expression or activity of the polynucleotides and/or polypeptides of the invention. Such methods can be utilized, for example, for the identification of compounds that can ameliorate symptoms of disorders as recited herein. Such methods can include, but are not limited to, assays for identifying compounds and other substances that interact with (e.g., bind to) the polypeptides of the invention.

30 The invention provides a method for identifying a compound that binds to the polypeptide of the present invention comprising contacting the compound with the polypeptide under conditions and for a time sufficient to form a polypeptide/compound complex and detecting the complex, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide is identified.

Also provided is a method for identifying a compound that binds to the polypeptide comprising contacting the compound with the polypeptide in a cell for a time sufficient to form a polypeptide/compound complex wherein the complex drives expression of a reporter gene sequence in the cell and detecting the complex by detecting reporter gene sequence expression  
 5 so that if the polypeptide/compound complex is detected a compound that binds to the polypeptide is identified.

#### 4. BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the BLASTX amino acid sequence alignment between the protein  
 10 encoded by SEQ ID NO:3 (i.e. SEQ ID NO:4) (also identified as "prothrombinase-like") and human prothrombinase Fg12 protein ("prothrombinase Fg12") (SEQ ID NO:19), indicating that the two sequence share 53 % similarity over 291 amino acid residues of SEQ ID NO:4 and 37% identity over 291 amino acid residues of SEQ ID NO:4, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine,  
 15 H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine. Gaps are presented as dashes.

Figure 2 shows the BLASTP sequence alignment between the protein encoded by SEQ ID :13 (i.e. SEQ ID NO:14) identified as "prothrombinase-like" and a portion of human  
 20 prothrombinase Fg12 protein (amino acids 1-269 of SEQ ID NO:19), indicating that the two sequences share 54 % similarity over 132 amino acid residues of SEQ ID NO:14 and 37% . identity over the same 132 amino acid residues of SEQ ID NO:14, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine,  
 25 P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine. Gaps are presented as dashes.

Figure 3 shows the BLASTP alignment between the protein encoded by SEQ ID : 13 (i.e. SEQ ID NO: 14) identified as "prothrombinase-like" and human prothrombinase Fg12 protein (amino acids 163-270 of SEQ ID NO: 19), indicating that the two sequences share 34%  
 30 similarity over that 115 amino acid residue portion of SEQ ID NO: 14 and 40% identity over the same 115 amino acid residue portion of SEQ ID NO: 14, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine,

P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine,  
W=Tryptophan, Y=Tyrosine. Gaps are presented as dashes.

Figure 4 shows the MSI GeneAtlas alignment of SEQ ID NO:14 with the blood coagulation, pdb number =1fzc\_B indicating that the two sequences share 32% identity over  
5 224 amino acids and 69% over the same 224 amino acids, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine. The score 86.22, the estimated p-value is 7.2e-04 and the upper limit p value is  
10 1.1e-11. The MSI GeneAtlas Report allows a functional prediction based protein structure homology (both secondary and tertiary protein structure), which utilizes threading (SEQFold), high throughput modeling and sequence profile-based searching (PB90) methodologies.

## 5. DETAILED DESCRIPTION OF THE INVENTION

15 The prothrombinase-like polypeptide of SEQ ID NO: 4 is an approximately 388-amino acid protein with a predicted molecular mass of approximately 43 kDa unglycosylated. Protein database searches with the BLASTX algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990), herein incorporated by reference) indicate that SEQ ID NO: 4 is homologous to human prothrombinase Fg12 protein.  
20 Figure 1 shows the BLASTX amino acid sequence alignment between the protein encoded by SEQ ID NO:3 (i.e. SEQ ID NO: 4) (also identified as "prothrombinase-like") and the human prothrombinase Fg12 protein (SEQ ID NO: 19) indicating that the two sequences share 53% similarity over 291 amino acid residues of SEQ ID NO: 4 and 37% identity over the same 291 amino acid residues of SEQ ID NO: 4.

25 A predicted approximately twenty residue signal peptide is encoded from approximately residue 1 to 20 of SEQ ID NO:4. The mature portion (also represented as SEQ ID No: 9) is useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide was predicted using the Kyte/Doolittle algorithm (J. Molecular Biology, vol. 157, pp. 105-131 (1982)). One of skill in the art will recognize that  
30 the actual cleavage site may be different than that predicted by the computer program.

Using eMATRIX software package (Stanford University, Stanford CA), prothrombinase-like is expected to have fibrinogen beta and gamma chains in the C-terminal domain. Specifically, at residues 181 through 217 of SEQ ID NO:4, at residues 222 through 234 of SEQ ID NO:4, at residues 271 through 285 of SEQ ID NO:4, at residues 324 through

353 of SEQ ID NO:4. The domains corresponding to SEQ ID NO:4 are as follows, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine,

5 V=Valine, W=Tryptophan, Y=Tyrosine:

Residues 181-217 VMCDMDYRGGGWTVIQKRIDGIDFQRLWCDYLDGFG (designated as SEQ ID NO:5),

Residues 324-353 CSHLHNKTGWWFNECGLANLNGIHHFSGKL (designated as SEQ ID NO:6),

10 Residues 271-285 FKMHLGRYSGNAGDA (designated as SEQ ID No:7),

Residues 222-234 EFWLGLKKIFYIV (designated as SEQ ID NO:8).

A predicted soluble, secreted splice variant of SEQ ID NO: 4 is SEQ ID NO: 14. It is an approximately 326 amino acid protein with a predicted molecular mass of approximately 37 kDa unglycosylated. Protein database searches with the BLASTX algorithm (Altschul S.F. et al., J. Mol. Evol. 36:290-300 (1993) and Altschul S.F. et al., J. Mol. Biol. 21:403-10 (1990),  
15 herein incorporated by reference) indicate that SEQ ID NO: 14 is homologous to human prothrombinase Fg12 protein. Figures 2 and 3 shows the BLASTP amino acid sequence alignment between the protein encoded by SEQ ID NO:13 (i.e. SEQ ID NO: 14) (also identified as "prothrombinase-like") and the human prothrombinase Fg12 protein (SEQ ID  
20 NO: 19).

Using eMATRIX software package (Stanford University, Stanford CA), prothrombinase-like is expected to have fibrinogen beta and gamma chains in the C-terminal domain. Specifically, at residues 181 through 217 of SEQ ID NO:14, at residues 262 through 291 of SEQ ID NO:14. The domains corresponding to SEQ ID NO:14 are as follows, wherein  
25 A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine:  
Residues 181-217 VMCDMDYRGGGWTVIQKRIDGIDFQRLWCDYLDGFG (designated as  
30 SEQ ID NO:5),  
Residues 262-292 CSHLHNKTGWWFNECGLANLNGIHHFSGKL (designated as SEQ ID NO:6).

A predicted approximately twenty residue signal peptide is encoded from approximately residue 1 to residue 20 of both SEQ ID NO: 14 (SEQ ID NO: 16). The extracellular portion is

useful on its own. This can be confirmed by expression in mammalian cells and sequencing of the cleaved product. The signal peptide region was predicted using the Kyte-Doolittle hydrophobicity prediction algorithm (J. Mol Biol, 157, pp. 105-31 (1982), incorporated herein by reference). One of skill in the art will recognize that the actual cleavage site may be

5 different than that predicted by the computer program.

Using Molecular Simulations Inc. GeneAtlas software (Molecular Simulations Inc., San Diego, CA) and the sequence profile-based searching protocol PB90 using optimized PSI-BLAST and the Seqfold based searching protocol on the PSI-BLAST defined domains, the membrane-bound prothrombinase-like polypeptide SEQ ID NO:14 was determined to have a  
10 region at residues 82 to 321 (SEQ ID NO:17) with characteristic motifs to blood coagulation (SEQ ID NO: 20) (See Figure 4). The domains corresponding to SEQ ID NO: 14 are as follows wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine,  
15 T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine:

Blood coagulation domain (designated as amino acids 82 – 321 of SEQ ID NO: 14) with PSI-BLAST e-value of 1.3e-66, protein database identification number entry = lfzc\_B (Research Collaboratory for Structural Bioinformatics. <http://www.rcsb.org/pdb>).

The polypeptides and polynucleotides of the invention can be utilized, for example, as  
20 part of methods for the prevention and/or treatment of disorders mediated by loss or overexpression of prothrombinase-like polypeptide. Such disorders include diseases associated with the dysregulation of coagulation pathways, such as hemophilia, myocardial infarction, glomerular disease, diabetes, fulminant viral hepatitis and atherosclerosis among others. The polynucleotides and polypeptides may also be beneficial in the treatment of viral infections and  
25 in the treatment of some forms of cancer.

## 5.1 DEFINITIONS

It must be noted that as used herein and in the appended claims, the singular forms “a”, “an” and “the” include plural references unless the context clearly dictates otherwise.

30 The term "active" refers to those forms of the polypeptide which retain the biologic and/or immunologic activities of any naturally occurring polypeptide. According to the invention, the terms “biologically active” or “biological activity” refer to a protein or peptide having structural, regulatory or biochemical functions of a naturally occurring molecule. The term “prothrombinase-like” refers to biological activity that is similar to the biological activity



of a human prothrombinase. Likewise "biologically active" or "biological activity" refers to the capability of the natural, recombinant or synthetic prothrombinase-like peptide, or any peptide thereof, to induce a specific biological response in appropriate animals or cells and to bind with specific antibodies.

5       The term "activated cells" as used in this application are those cells which are engaged in extracellular or intracellular membrane trafficking, including the export of secretory or enzymatic molecules as part of a normal or disease process.

          The terms "complementary" or "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence 5'-AGT-3' binds to the  
10       complementary sequence 3'-TCA-5'. Complementarity between two single-stranded molecules may be "partial" such that only some of the nucleic acids bind or it may be "complete" such that total complementarity exists between the single stranded molecules. The degree of complementarity between the nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands.

15       The term "embryonic stem cells (ES)" refers to a cell that can give rise to many differentiated cell types in an embryo or an adult, including the germ cells. The term "germ line stem cells (GSCs)" refers to stem cells derived from primordial stem cells that provide a steady and continuous source of germ cells for the production of gametes. The term  
20       "primordial germ cells (PGCs)" refers to a small population of cells set aside from other cell lineages particularly from the yolk sac, mesenteries, or gonadal ridges during embryogenesis that have the potential to differentiate into germ cells and other cells. PGCs are the source from which GSCs and ES cells are derived. The PGCs, the GSCs and the ES cells are capable of self-renewal. Thus these cells not only populate the germ line and give rise to a plurality of terminally differentiated cells that comprise the adult specialized organs, but are able to  
25       regenerate themselves.

          The term "expression modulating fragment," EMF, means a series of nucleotides which modulates the expression of an operably linked ORF or another EMF.

          As used herein, a sequence is said to "modulate the expression of an operably linked sequence" when the expression of the sequence is altered by the presence of the EMF. EMFs  
30       include, but are not limited to, promoters, and promoter modulating sequences (inducible elements). One class of EMFs is nucleic acid fragments which induce the expression of an operably linked ORF in response to a specific regulatory factor or physiological event.

          The terms "nucleotide sequence" or "nucleic acid" or "polynucleotide" or "oligonucleotide" are used interchangeably and refer to a heteropolymer of nucleotides or the

sequence of these nucleotides. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA) or to any DNA-like or RNA-like material. Generally, nucleic acid segments provided by this invention may be assembled from  
5 fragments of the genome and short oligonucleotide linkers, or from a series of oligonucleotides, or from individual nucleotides, to provide a synthetic nucleic acid which is capable of being expressed in a recombinant transcriptional unit comprising regulatory elements derived from a microbial or viral operon, or a eukaryotic gene.

The terms "oligonucleotide fragment" or a "polynucleotide fragment", "portion," or  
10 "segment" or "probe" or "primer" are used interchangeable and refer to a sequence of nucleotide residues which are at least about 5 nucleotides, more preferably at least about 7 nucleotides, more preferably at least about 9 nucleotides, more preferably at least about 11 nucleotides and most preferably at least about 17 nucleotides. The fragment is preferably less than about 500 nucleotides, preferably less than about 200 nucleotides, more preferably less  
15 than about 100 nucleotides, more preferably less than about 50 nucleotides and most preferably less than 30 nucleotides. Preferably the probe is from about 6 nucleotides to about 200 nucleotides, preferably from about 15 to about 50 nucleotides, more preferably from about 17 to 30 nucleotides and most preferably from about 20 to 25 nucleotides. Preferably the fragments can be used in polymerase chain reaction (PCR), various hybridization procedures  
20 or microarray procedures to identify or amplify identical or related parts of mRNA or DNA molecules. A fragment or segment may uniquely identify each polynucleotide sequence of the present invention. Preferably the fragment comprises a sequence substantially similar to a portion of SEQ ID NO: 1-3, 10-13 and 15.

Probes may, for example, be used to determine whether specific mRNA molecules are  
25 present in a cell or tissue or to isolate similar nucleic acid sequences from chromosomal DNA as described by Walsh et al. (Walsh, P.S. et al., 1992, PCR Methods Appl 1:241-250). They may be labeled by nick translation, Klenow fill-in reaction, PCR, or other methods well known in the art. Probes of the present invention, their preparation and/or labeling are elaborated in Sambrook, J. et al., 1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor  
30 Laboratory, NY; or Ausubel, F.M. et al., 1989, Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, both of which are incorporated herein by reference in their entirety.

The nucleic acid sequences of the present invention also include the sequence information from any of the nucleic acid sequences of SEQ ID NO: 1-3, 10-13 and 15. The

sequence information can be a segment of SEQ ID NO: 1-3, 10-13 and 15 that uniquely identifies or represents the sequence information of SEQ ID NO: 1-3, 10-13 and 15. One such segment can be a twenty-mer nucleic acid sequence because the probability that a twenty-mer is fully matched in the human genome is 1 in 300. In the human genome, there are three  
5 billion base pairs in one set of chromosomes. Because  $4^{20}$  possible twenty-mers exist, there are 300 times more twenty-mers than there are base pairs in a set of human chromosome. Using the same analysis, the probability for a seventeen-mer to be fully matched in the human genome is approximately 1 in 5. When these segments are used in arrays for expression studies, fifteen-mer segments can be used. The probability that the fifteen-mer is fully  
10 matched in the expressed sequences is also approximately one in five because expressed sequences comprise less than approximately 5% of the entire genome sequence.

Similarly, when using sequence information for detecting a single mismatch, a segment can be a twenty-five mer. The probability that the twenty-five mer would appear in a human genome with a single mismatch is calculated by multiplying the probability for a full match  
15  $(1/4^{25})$  times the increased probability for mismatch at each nucleotide position  $(3 \times 25)$ . The probability that an eighteen mer with a single mismatch can be detected in an array for expression studies is approximately one in five. The probability that a twenty-mer with a single mismatch can be detected in a human genome is approximately one in five.

The term "open reading frame," ORF, means a series of nucleotide triplets coding for  
20 amino acids without any termination codons and is a sequence translatable into protein.

The terms "operably linked" or "operably associated" refer to functionally related nucleic acid sequences. For example, a promoter is operably associated or operably linked with a coding sequence if the promoter controls the transcription of the coding sequence. While operably linked nucleic acid sequences can be contiguous and in the same reading  
25 frame, certain genetic elements e.g. repressor genes are not contiguously linked to the coding sequence but still control transcription/translation of the coding sequence.

The term "pluripotent" refers to the capability of a cell to differentiate into a number of differentiated cell types that are present in an adult organism. A pluripotent cell is restricted in its differentiation capability in comparison to a totipotent cell.

30 The terms "polypeptide" or "peptide" or "amino acid sequence" refer to an oligopeptide, peptide, polypeptide or protein sequence or fragment thereof and to naturally occurring or synthetic molecules. A polypeptide "fragment," "portion," or "segment" is a stretch of amino acid residues of at least about 5 amino acids, preferably at least about 7 amino

acids, more preferably at least about 9 amino acids and most preferably at least about 17 or more amino acids. The peptide preferably is not greater than about 200 amino acids, more preferably less than 150 amino acids and most preferably less than 100 amino acids.

Preferably the peptide is from about 5 to about 200 amino acids. To be active, any  
5 polypeptide must have sufficient length to display biological and/or immunological activity.

The term "naturally occurring polypeptide" refers to polypeptides produced by cells that have not been genetically engineered and specifically contemplates various polypeptides arising from post-translational modifications of the polypeptide including, but not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation and acylation.

10 The term "translated protein coding portion" means a sequence which encodes for the full length protein which may include any leader sequence or a processing sequence.

The term "mature protein coding sequence" refers to a sequence which encodes a peptide or protein without any leader/signal sequence. The peptide may have the leader sequences removed during processing in the cell or the protein may have been produced  
15 synthetically or using a polynucleotide only encoding for the mature protein coding sequence.

The term "derivative" refers to polypeptides chemically modified by such techniques as ubiquitination, labeling (e.g., with radionuclides or various enzymes), covalent polymer attachment such as pegylation (derivatization with polyethylene glycol) and insertion or substitution by chemical synthesis of amino acids such as ornithine, which do not normally  
20 occur in human proteins.

The term "variant"(or "analog") refers to any polypeptide differing from naturally occurring polypeptides by amino acid insertions, deletions, and substitutions, created using, *e g.*, recombinant DNA techniques. Guidance in determining which amino acid residues may be replaced, added or deleted without abolishing activities of interest, may be found by comparing  
25 the sequence of the particular polypeptide with that of homologous peptides and minimizing the number of amino acid sequence changes made in regions of high homology (conserved regions) or by replacing amino acids with consensus sequence.

Alternatively, recombinant variants encoding these same or similar polypeptides may be synthesized or selected by making use of the "redundancy" in the genetic code. Various codon  
30 substitutions, such as the silent changes which produce various restriction sites, may be introduced to optimize cloning into a plasmid or viral vector or expression in a particular prokaryotic or eukaryotic system. Mutations in the polynucleotide sequence may be reflected in the polypeptide or domains of other peptides added to the polypeptide to modify the

properties of any part of the polypeptide, to change characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover rate.

Preferably, amino acid "substitutions" are the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, *i.e.*, conservative amino acid replacements. "Conservative" amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues involved. For example, nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine; polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine; positively charged (basic) amino acids include arginine, lysine, and histidine; and negatively charged (acidic) amino acids include aspartic acid and glutamic acid. "Insertions" or "deletions" are preferably in the range of about 1 to 20 amino acids, more preferably 1 to 10 amino acids. The variation allowed may be experimentally determined by systematically making insertions, deletions, or substitutions of amino acids in a polypeptide molecule using recombinant DNA techniques and assaying the resulting recombinant variants for activity.

Alternatively, where alteration of function is desired, insertions, deletions or non-conservative alterations can be engineered to produce altered polypeptides. Such alterations can, for example, alter one or more of the biological functions or biochemical characteristics of the polypeptides of the invention. For example, such alterations may change polypeptide characteristics such as ligand-binding affinities, interchain affinities, or degradation/turnover rate. Further, such alterations can be selected so as to generate polypeptides that are better suited for expression, scale up and the like in the host cells chosen for expression. For example, cysteine residues can be deleted or substituted with another amino acid residue in order to eliminate disulfide bridges.

The terms "purified" or "substantially purified" as used herein denotes that the indicated nucleic acid or polypeptide is present in the substantial absence of other biological macromolecules, *e.g.*, polynucleotides, proteins, and the like. In one embodiment, the polynucleotide or polypeptide is purified such that it constitutes at least 95% by weight, more preferably at least 99% by weight, of the indicated biological macromolecules present (but water, buffers, and other small molecules, especially molecules having a molecular weight of less than 1000 daltons, can be present).

The term "isolated" as used herein refers to a nucleic acid or polypeptide separated from at least one other component (*e.g.*, nucleic acid or polypeptide) present with the nucleic

acid or polypeptide in its natural source. In one embodiment, the nucleic acid or polypeptide is found in the presence of (if anything) only a solvent, buffer, ion, or other components normally present in a solution of the same. The terms "isolated" and "purified" do not encompass nucleic acids or polypeptides present in their natural source.

5           The term "recombinant," when used herein to refer to a polypeptide or protein, means that a polypeptide or protein is derived from recombinant (*e.g.*, microbial, insect, or mammalian) expression systems. "Microbial" refers to recombinant polypeptides or proteins made in bacterial or fungal (*e.g.*, yeast) expression systems. As a product, "recombinant microbial" defines a polypeptide or protein essentially free of native endogenous substances  
10   and unaccompanied by associated native glycosylation. Polypeptides or proteins expressed in most bacterial cultures, *e.g.*, *E. coli*, will be free of glycosylation modifications; polypeptides or proteins expressed in yeast will have a glycosylation pattern in general different from those expressed in mammalian cells.

          The term "recombinant expression vehicle or vector" refers to a plasmid or phage or  
15   virus or vector, for expressing a polypeptide from a DNA (RNA) sequence. An expression vehicle can comprise a transcriptional unit comprising an assembly of (1) a genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers, (2) a structural or coding sequence which is transcribed into mRNA and translated into protein, and (3) appropriate transcription initiation and termination sequences. Structural units intended  
20   for use in yeast or eukaryotic expression systems preferably include a leader sequence enabling extracellular secretion of translated protein by a host cell. Alternatively, where recombinant protein is expressed without a leader or transport sequence, it may include an amino terminal methionine residue. This residue may or may not be subsequently cleaved from the expressed recombinant protein to provide a final product.

25           The term "recombinant expression system" means host cells which have stably integrated a recombinant transcriptional unit into chromosomal DNA or carry the recombinant transcriptional unit extrachromosomally. Recombinant expression systems as defined herein will express heterologous polypeptides or proteins upon induction of the regulatory elements linked to the DNA segment or synthetic gene to be expressed. This term also means host cells  
30   which have stably integrated a recombinant genetic element or elements having a regulatory role in gene expression, for example, promoters or enhancers. Recombinant expression systems as defined herein will express polypeptides or proteins endogenous to the cell upon induction of the regulatory elements linked to the endogenous DNA segment or gene to be expressed. The cells can be prokaryotic or eukaryotic.

The term "secreted" includes a protein that is transported across or through a membrane, including transport as a result of signal sequences in its amino acid sequence when it is expressed in a suitable host cell. "Secreted" proteins include without limitation proteins secreted wholly (*e.g.*, soluble proteins) or partially (*e.g.*, receptors) from the cell in which they are expressed. "Secreted" proteins also include without limitation proteins that are transported across the membrane of the endoplasmic reticulum. "Secreted" proteins are also intended to include proteins containing non-typical signal sequences (*e.g.* Interleukin-1 Beta, see Krasney, P.A. and Young, P.R. (1992) Cytokine 4(2):134 -143) and factors released from damaged cells (*e.g.* Interleukin-1 Receptor Antagonist, see Arend, W.P. et. al. (1998) Annu. Rev. Immunol. 16:27-55)

Where desired, an expression vector may be designed to contain a "signal or leader sequence" which will direct the polypeptide through the membrane of a cell. Such a sequence may be naturally present on the polypeptides of the present invention or provided from heterologous protein sources by recombinant DNA techniques.

The term "stringent" is used to refer to conditions that are commonly understood in the art as stringent. Stringent conditions can include highly stringent conditions (*i.e.*, hybridization to filter-bound DNA in 0.5 M NaHPO<sub>4</sub>, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1X SSC/0.1% SDS at 68°C), and moderately stringent conditions (*i.e.*, washing in 0.2X SSC/0.1% SDS at 42°C). Other exemplary hybridization conditions are described herein in the examples.

In instances of hybridization of deoxyoligonucleotides, additional exemplary stringent hybridization conditions include washing in 6X SSC/0.05% sodium pyrophosphate at 37°C (for 14-base oligonucleotides), 48°C (for 17-base oligonucleotides), 55°C (for 20-base oligonucleotides), and 60°C (for 23-base oligonucleotides).

As used herein, "substantially equivalent" can refer both to nucleotide and amino acid sequences, for example a mutant sequence, that varies from a reference sequence by one or more substitutions, deletions, or additions, the net effect of which does not result in an adverse functional dissimilarity between the reference and subject sequences. Typically, such a substantially equivalent sequence varies from one of those listed herein by no more than about 35% (*i.e.*, the number of individual residue substitutions, additions, and/or deletions in a substantially equivalent sequence, as compared to the corresponding reference sequence, divided by the total number of residues in the substantially equivalent sequence is about 0.35 or less). Such a sequence is said to have 65% sequence identity to the listed sequence. In one

embodiment, a substantially equivalent, *e.g.*, mutant, sequence of the invention varies from a listed sequence by no more than 30% (70% sequence identity); in a variation of this embodiment, by no more than 25% (75% sequence identity); and in a further variation of this embodiment, by no more than 20% (80% sequence identity) and in a further variation of this embodiment, by no more than 10% (90% sequence identity) and in a further variation of this embodiment, by no more than 5% (95% sequence identity). Substantially equivalent, *e.g.*, mutant, amino acid sequences according to the invention preferably have at least 80% sequence identity with a listed amino acid sequence, more preferably at least 90% sequence identity. Substantially equivalent nucleotide sequence of the invention can have lower percent sequence identities, taking into account, for example, the redundancy or degeneracy of the genetic code. Preferably, nucleotide sequence has at least about 65% identity, more preferably at least about 75% identity, and most preferably at least about 95% identity. For the purposes of the present invention, sequences having substantially equivalent biological activity and substantially equivalent expression characteristics are considered substantially equivalent. For the purposes of determining equivalence, truncation of the mature sequence (*e.g.*, via a mutation which creates a spurious stop codon) should be disregarded. Sequence identity may be determined, *e.g.*, using the Jotun Hein method (Hein, J. (1990) *Methods Enzymol.* 183:626-645). Identity between sequences can also be determined by other methods known in the art, *e.g.* by varying hybridization conditions.

20       The term "totipotent" refers to the capability of a cell to differentiate into all of the cell types of an adult organism.

      The term "transformation" means introducing DNA into a suitable host cell so that the DNA is replicable, either as an extrachromosomal element, or by chromosomal integration. The term "transfection" refers to the taking up of an expression vector by a suitable host cell, whether or not any coding sequences are in fact expressed. The term "infection" refers to the introduction of nucleic acids into a suitable host cell by use of a virus or viral vector.

25       As used herein, an "uptake modulating fragment," UMF, means a series of nucleotides which mediate the uptake of a linked DNA fragment into a cell. UMFs can be readily identified using known UMFs as a target sequence or target motif with the computer-based systems described below. The presence and activity of a UMF can be confirmed by attaching the suspected UMF to a marker sequence. The resulting nucleic acid molecule is then incubated with an appropriate host under appropriate conditions and the uptake of the marker sequence is determined. As described above, a UMF will increase the frequency of uptake of a linked marker sequence.



Each of the above terms is meant to encompass all that is described for each, unless the context dictates otherwise.

## 5.2 NUCLEIC ACIDS OF THE INVENTION

5       The invention is based on the discovery of a novel secreted prothrombinase-like polypeptide, the polynucleotides encoding the prothrombinase-like polypeptide and the use of these compositions for the diagnosis, treatment or prevention of neurological conditions and disorders.

10       The isolated polynucleotides of the invention include, but are not limited to a polynucleotide comprising any of the nucleotide sequences of SEQ ID NO: 1-3, 10-13 and 15; a fragment of SEQ ID NO: 1-3, 10-13 and 15; a polynucleotide comprising the full length protein coding sequence of SEQ ID NO: 1-3, 10-13 and 15 (for example SEQ ID NO: 4, 9, 14, 18); and a polynucleotide comprising the nucleotide sequence encoding the mature protein coding sequence of the polynucleotides of any one of SEQ ID NO: 1-3, 10-13 and 15. The  
15       polynucleotides of the present invention also include, but are not limited to, a polynucleotide that hybridizes under stringent conditions to (a) the complement of any of the nucleotide sequences of the SEQ ID NO: 1-3, 10-13 or 15; (b) a polynucleotide encoding any one of the polypeptides of SEQ ID NO: 4-9, 14 or 16-18; (c) a polynucleotide which is an allelic variant of any polynucleotides recited above; (d) a polynucleotide which encodes a species homolog of  
20       any of the proteins recited above; or (e) a polynucleotide that encodes a polypeptide comprising a specific domain or truncation of the polypeptides of SEQ ID NO: 4-9, 14 or 16-18. Domains of interest may depend on the nature of the encoded polypeptide; e.g., domains in receptor-like polypeptides include ligand-binding, extracellular, transmembrane, or cytoplasmic domains, or combinations thereof; domains in immunoglobulin-like proteins  
25       include the variable immunoglobulin-like domains; domains in enzyme-like polypeptides include catalytic and substrate binding domains; and domains in ligand polypeptides include receptor-binding domains.

30       The polynucleotides of the invention include naturally occurring or wholly or partially synthetic DNA, e.g., cDNA and genomic DNA, and RNA, e.g., mRNA. The polynucleotides may include all of the coding region of the cDNA or may represent a portion of the coding region of the cDNA.

The present invention also provides genes corresponding to the cDNA sequences disclosed herein. The corresponding genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the preparation of probes or

primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials. Further 5' and 3' sequence can be obtained using methods known in the art. For example, full length cDNA or genomic DNA that corresponds to any of the polynucleotides of the SEQ ID NO:1-3, 10-13 and 5 15 can be obtained by screening appropriate cDNA or genomic DNA libraries under suitable hybridization conditions using any of the polynucleotides of the SEQ ID NO:1-3, 10-13 and 15 or a portion thereof as a probe. Alternatively, the polynucleotides of the SEQ ID NO: 1-3, 10-13 and 15 may be used as the basis for suitable primer(s) that allow identification and/or amplification of genes in appropriate genomic DNA or cDNA libraries.

10 The nucleic acid sequences of the invention can be assembled from ESTs and sequences (including cDNA and genomic sequences) obtained from one or more public databases, such as dbEST, gbpr, and UniGene. The EST sequences can provide identifying sequence information, representative fragment or segment information, or novel segment information for the full-length gene.

15 The polynucleotides of the invention also provide polynucleotides including nucleotide sequences that are substantially equivalent to the polynucleotides recited above. Polynucleotides according to the invention can have, *e.g.*, at least about 65%, at least about 70%, at least about 75%, at least about 80%, more typically at least about 90%, and even more typically at least about 95%, sequence identity to a polynucleotide recited above.

20 Included within the scope of the nucleic acid sequences of the invention are nucleic acid sequence fragments that hybridize under stringent conditions to any of the nucleotide sequences of the SEQ ID NO: 1-3, 10-13 and 15, or complements thereof, which fragment is greater than about 5 nucleotides, preferably 7 nucleotides, more preferably greater than 9 nucleotides and most preferably greater than 17 nucleotides. Fragments of, *e.g.* 15, 17, or 20 25 nucleotides or more that are selective for (*i.e.* specifically hybridize to any one of the polynucleotides of the invention) are contemplated. Probes capable of specifically hybridizing to a polynucleotide can differentiate polynucleotide sequences of the invention from other polynucleotide sequences in the same family of genes or can differentiate human genes from genes of other species, and are preferably based on unique nucleotide sequences.

30 The sequences falling within the scope of the present invention are not limited to these specific sequences, but also include allelic and species variations thereof. Allelic and species variations can be routinely determined by comparing the sequence provided in SEQ ID NO: 1-3, 10-13 and 15, a representative fragment thereof, or a nucleotide sequence at least 90% identical, preferably 95% identical, to SEQ ID NO: 1-3, 10-13 and 15 with a sequence from another isolate

of the same species. Furthermore, to accommodate codon variability, the invention includes nucleic acid molecules coding for the same amino acid sequences as do the specific ORFs disclosed herein. In other words, in the coding region of an ORF, substitution of one codon for another codon that encodes the same amino acid is expressly contemplated.

5           The nearest neighbor result for the nucleic acids of the present invention, including SEQ ID NO: 1-3, 10-13 and 15, can be obtained by searching a database using an algorithm or a program. Preferably, a BLAST which stands for Basic Local alignment Search Tool is used to search for local sequence alignments (Altshul, S.F. J Mol. Evol. 36 290-300 (1993) and Altschul S.F. et al. J. Mol. Biol. 21:403-410 (1990))

10           Species homologs (or orthologs) of the disclosed polynucleotides and proteins are also provided by the present invention. Species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source from the desired species.

          The invention also encompasses allelic variants of the disclosed polynucleotides or  
15 proteins; that is, naturally-occurring alternative forms of the isolated polynucleotide which also encode proteins which are identical, homologous or related to that encoded by the polynucleotides.

          The nucleic acid sequences of the invention are further directed to sequences which encode variants of the described nucleic acids. These amino acid sequence variants may be  
20 prepared by methods known in the art by introducing appropriate nucleotide changes into a native or variant polynucleotide. There are two variables in the construction of amino acid sequence variants: the location of the mutation and the nature of the mutation. Nucleic acids encoding the amino acid sequence variants are preferably constructed by mutating the polynucleotide to encode an amino acid sequence that does not occur in nature. These nucleic  
25 acid alterations can be made at sites that differ in the nucleic acids from different species (variable positions) or in highly conserved regions (constant regions). Sites at such locations will typically be modified in series, *e.g.*, by substituting first with conservative choices (*e.g.*, hydrophobic amino acid to a different hydrophobic amino acid) and then with more distant choices (*e.g.*, hydrophobic amino acid to a charged amino acid), and then deletions or  
30 insertions may be made at the target site. Amino acid sequence deletions generally range from about 1 to 30 residues, preferably about 1 to 10 residues, and are typically contiguous. Amino acid insertions include amino- and/or carboxyl-terminal fusions ranging in length from one to one hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Intrasequence insertions may range generally from about 1 to 10 amino

residues, preferably from 1 to 5 residues. Examples of terminal insertions include the heterologous signal sequences necessary for secretion or for intracellular targeting in different host cells and sequences such as FLAG or poly-histidine sequences useful for purifying the expressed protein.

5 In a preferred method, polynucleotides encoding the novel amino acid sequences are changed via site-directed mutagenesis. This method uses oligonucleotide sequences to alter a polynucleotide to encode the desired amino acid variant, as well as sufficient adjacent nucleotides on both sides of the changed amino acid to form a stable duplex on either side of the site being changed. In general, the techniques of site-directed mutagenesis are well known  
10 to those of skill in the art and this technique is exemplified by publications such as, Edelman et al., *DNA* 2:183 (1983). A versatile and efficient method for producing site-specific changes in a polynucleotide sequence was published by Zoller and Smith, *Nucleic Acids Res.* 10:6487-6500 (1982). PCR may also be used to create amino acid sequence variants of the novel nucleic acids. When small amounts of template DNA are used as starting material, primer(s)  
15 that differs slightly in sequence from the corresponding region in the template DNA can generate the desired amino acid variant. PCR amplification results in a population of product DNA fragments that differ from the polynucleotide template encoding the polypeptide at the position specified by the primer. The product DNA fragments replace the corresponding region in the plasmid and this gives a polynucleotide encoding the desired amino acid variant.

20 A further technique for generating amino acid variants is the cassette mutagenesis technique described in Wells et al., *Gene* 34:315 (1985); and other mutagenesis techniques well known in the art, such as, for example, the techniques in Sambrook et al., *supra*, and *Current Protocols in Molecular Biology*, Ausubel et al. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally  
25 equivalent amino acid sequence may be used in the practice of the invention for the cloning and expression of these novel nucleic acids. Such DNA sequences include those which are capable of hybridizing to the appropriate novel nucleic acid sequence under stringent conditions.

Polynucleotides encoding preferred polypeptide truncations of the invention can be used to generate polynucleotides encoding chimeric or fusion proteins comprising one or more  
30 domains of the invention and heterologous protein sequences.

The polynucleotides of the invention additionally include the complement of any of the polynucleotides recited above. The polynucleotide can be DNA (genomic, cDNA, amplified, or synthetic) or RNA. Methods and algorithms for obtaining such polynucleotides are well known to those of skill in the art and can include, for example, methods for determining

hybridization conditions that can routinely isolate polynucleotides of the desired sequence identities.

In accordance with the invention, polynucleotide sequences comprising the mature protein coding sequences corresponding to any one of SEQ ID NO: 1-3, 10-13 and 15, or  
5 functional equivalents thereof, may be used to generate recombinant DNA molecules that direct the expression of that nucleic acid, or a functional equivalent thereof, in appropriate host cells. Also included are the cDNA inserts of any of the clones identified herein.

A polynucleotide according to the invention can be joined to any of a variety of other nucleotide sequences by well-established recombinant DNA techniques (see Sambrook J et al.  
10 (1989) Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, NY). Useful nucleotide sequences for joining to polynucleotides include an assortment of vectors, e.g., plasmids, cosmids, lambda phage derivatives, phagemids, and the like, that are well known in the art. Accordingly, the invention also provides a vector including a polynucleotide of the invention and a host cell containing the polynucleotide. In general, the vector contains  
15 an origin of replication functional in at least one organism, convenient restriction endonuclease sites, and a selectable marker for the host cell. Vectors according to the invention include expression vectors, replication vectors, probe generation vectors, and sequencing vectors. A host cell according to the invention can be a prokaryotic or eukaryotic cell and can be a unicellular organism or part of a multicellular organism.

20 The present invention further provides recombinant constructs comprising a nucleic acid having any of the nucleotide sequences of the SEQ ID NO: 1-3, 10-13 and 15 or a fragment thereof or any other polynucleotides of the invention. In one embodiment, the recombinant constructs of the present invention comprise a vector, such as a plasmid or viral vector, into which a nucleic acid having any of the nucleotide sequences of the SEQ ID NO: 1-  
25 3, 10-13 and 15 or a fragment thereof is inserted, in a forward or reverse orientation. In the case of a vector comprising one of the ORFs of the present invention, the vector may further comprise regulatory sequences, including for example, a promoter, operably linked to the ORF. Large numbers of suitable vectors and promoters are known to those of skill in the art and are commercially available for generating the recombinant constructs of the present  
30 invention. The following vectors are provided by way of example. Bacterial: pBs, phagescript, PsiX174, pBluescript SK, pBs KS, pNH8a, pNH16a, pNH18a, pNH46a (Stratagene); pTrc99A, pKK223-3, pKK233-3, pDR540, pRIT5 (Pharmacia). Eukaryotic: pWLneo, pSV2cat, pOG44, PXTI, pSG (Stratagene) pSVK3, pBPV, pMSG, pSVL (Pharmacia).

The isolated polynucleotide of the invention may be operably linked to an expression control sequence such as the pMT2 or pED expression vectors disclosed in Kaufman et al., *Nucleic Acids Res.* 19, 4485-4490 (1991), in order to produce the protein recombinantly. Many suitable expression control sequences are known in the art. General methods of

5 expressing recombinant proteins are also known and are exemplified in R. Kaufman, *Methods in Enzymology* 185, 537-566 (1990). As defined herein "operably linked" means that the isolated polynucleotide of the invention and an expression control sequence are situated within a vector or cell in such a way that the protein is expressed by a host cell which has been transformed (transfected) with the ligated polynucleotide/expression control sequence.

10 Promoter regions can be selected from any desired gene using CAT (chloramphenicol transferase) vectors or other vectors with selectable markers. Two appropriate vectors are pKK232-8 and pCM7. Particular named bacterial promoters include lacI, lacZ, T3, T7, gpt, lambda PR, and trc. Eukaryotic promoters include CMV immediate early, HSV thymidine kinase, early and late SV40, LTRs from retrovirus, and mouse metallothionein-I. Selection of

15 the appropriate vector and promoter is well within the level of ordinary skill in the art. Generally, recombinant expression vectors will include origins of replication and selectable markers permitting transformation of the host cell, *e.g.*, the ampicillin resistance gene of *E. coli* and *S. cerevisiae* TRP1 gene, and a promoter derived from a highly expressed gene to direct transcription of a downstream structural sequence. Such promoters can be derived from

20 operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), a-factor, acid phosphatase, or heat shock proteins, among others. The heterologous structural sequence is assembled in appropriate phase with translation initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode

25 a fusion protein including an amino terminal identification peptide imparting desired characteristics, *e.g.*, stabilization or simplified purification of expressed recombinant product. Useful expression vectors for bacterial use are constructed by inserting a structural DNA sequence encoding a desired protein together with suitable translation initiation and termination signals in operable reading phase with a functional promoter. The vector will comprise one or

30 more phenotypic selectable markers and an origin of replication to ensure maintenance of the vector and to, if desirable, provide amplification within the host. Suitable prokaryotic hosts for transformation include *E. coli*, *Bacillus subtilis*, *Salmonella typhimurium* and various species within the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*, although others may also be employed as a matter of choice.

As a representative but non-limiting example, useful expression vectors for bacterial use can comprise a selectable marker and bacterial origin of replication derived from commercially available plasmids comprising genetic elements of the well known cloning vector pBR322 (ATCC 37017). Such commercial vectors include, for example, pKK223-3  
5 (Pharmacia Fine Chemicals, Uppsala, Sweden) and GEM 1 (Promega Biotech, Madison, WI, USA). These pBR322 "backbone" sections are combined with an appropriate promoter and the structural sequence to be expressed. Following transformation of a suitable host strain and growth of the host strain to an appropriate cell density, the selected promoter is induced or derepressed by appropriate means (*e.g.*, temperature shift or chemical induction) and cells are  
10 cultured for an additional period. Cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification.

Polynucleotides of the invention can also be used to induce immune responses. For example, as described in Fan et al., *Nat. Biotech.* 17:870-872 (1999), incorporated herein by reference, nucleic acid sequences encoding a polypeptide may be used to generate antibodies  
15 against the encoded polypeptide following topical administration of naked plasmid DNA or following injection, and preferably intramuscular injection of the DNA. The nucleic acid sequences are preferably inserted in a recombinant expression vector and may be in the form of naked DNA.

### 5.3 ANTISENSE

20 Another aspect of the invention pertains to isolated antisense nucleic acid molecules that are hybridizable to or complementary to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1-3, 10-13 and 15, or fragments, analogs or derivatives thereof. An "antisense" nucleic acid comprises a nucleotide sequence that is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded  
25 cDNA molecule or complementary to an mRNA sequence. In specific aspects, antisense nucleic acid molecules are provided that comprise a sequence complementary to at least about 10, 25, 50, 100, 250 or 500 nucleotides or an entire coding strand, or to only a portion thereof. Nucleic acid molecules encoding fragments, homologs, derivatives and analogs of a protein of any of SEQ ID NO:4-9, 14 and 16-18 or antisense nucleic acids complementary to a  
30 nucleic acid sequence of SEQ ID NO:1-3, 10-13 and 15 are additionally provided.

In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence of the invention. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues. In another embodiment, the antisense nucleic acid molecule is antisense to

a "noncoding region" of the coding strand of a nucleotide sequence of the invention. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

Given the coding strand sequences encoding a nucleic acid disclosed herein (*e.g.*, SEQ  
5 ID NO:1-3, 10-13 and 15, antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick or Hoogsteen base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of a mRNA, but more preferably is an oligonucleotide that is antisense to only a portion of the coding or noncoding region of a mRNA. For example, the antisense oligonucleotide can be complementary to the region  
10 surrounding the translation start site of a mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of the invention can be constructed using chemical synthesis or enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides  
15 or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used.

Examples of modified nucleotides that can be used to generate the antisense nucleic acid  
20 include: 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine,  
25 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil,  
30 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).



The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a protein according to the invention to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by  
5 conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule that binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention includes direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered  
10 systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies that bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of  
15 antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an  $\alpha$ -anomeric nucleic acid molecule. An  $\alpha$ -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual  $\beta$ -units, the  
20 strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res* 15: 6625-6641). The antisense nucleic acid molecule can also comprise a 2'-*o*-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res* 15: 6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett* 215: 327-330).

#### 25 5.4 RIBOZYMES AND PNA MOIETIES

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity that are capable of cleaving a single-stranded nucleic acid, such as a mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes (described in Haselhoff and Gerlach (1988)  
30 *Nature* 334:585-591)) can be used to catalytically cleave a mRNA transcripts to thereby inhibit translation of a mRNA. A ribozyme having specificity for a nucleic acid of the invention can be designed based upon the nucleotide sequence of a DNA disclosed herein (*i.e.*, SEQ ID NO:1-3, 10-13 and 15). For example, a derivative of a Tetrahymena L-19 IVS RNA can be

constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a encoding a mRNA according to the invention. See, *e.g.*, Cech *et al.* U.S. Pat. No. 4,987,071; and Cech *et al.* U.S. Pat. No. 5,116,742.

Alternatively, mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel *et al.*, (1993) *Science* 261:1411-1418.

Alternatively, gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region (*e.g.*, promoter and/or enhancers) to form triple helical structures that prevent transcription of the gene in target cells. See generally, Helene. (1991) *Anticancer Drug Des.* 6: 569-84; Helene. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher (1992) *Bioassays* 14: 807-15.

In various embodiments, the nucleic acids of the invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acids can be modified to generate peptide nucleic acids (see Hyrup *et al.* (1996) *Bioorg Med Chem* 4: 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup *et al.* (1996) above; Perry-O'Keefe *et al.* (1996) *PNAS* 93: 14670-675.

PNAs of the invention can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, *e.g.*, inducing transcription or translation arrest or inhibiting replication. PNAs of the invention can also be used, *e.g.*, in the analysis of single base pair mutations in a gene by, *e.g.*, PNA directed PCR clamping; as artificial restriction enzymes when used in combination with other enzymes, *e.g.*, S1 nucleases (Hyrup B. (1996) above); or as probes or primers for DNA sequence and hybridization (Hyrup *et al.* (1996), above; Perry-O'Keefe (1996), above).

In another embodiment, PNAs of the invention can be modified, *e.g.*, to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras can be generated that may

combine the advantageous properties of PNA and DNA. Such chimeras allow DNA recognition enzymes, *e.g.*, RNase H and DNA polymerases, to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup (1996) above). The synthesis of PNA-DNA chimeras can be performed as described in Hyrup (1996) above and Finn *et al.* (1996) *Nucl Acids Res* 24: 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry, and modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used between the PNA and the 5' end of DNA (Mag *et al.* (1989) *Nucl Acid Res* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn *et al.* (1996) above). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment. See, Petersen *et al.* (1975) *Bioorg Med Chem Lett* 5: 1119-11124.

15 In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.*, 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaitre *et al.*, 1987, *Proc. Natl. Acad. Sci.* 84:648-652; PCT Publication No. W088/09810) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization triggered cleavage agents (See, *e.g.*, Krol *et al.*, 1988, *BioTechniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon, 1988, *Pharm. Res.* 5: 539-549). To this end, the oligonucleotide may be conjugated to another molecule, *e.g.*, a peptide, a hybridization triggered cross-linking agent, a transport agent, a hybridization-triggered cleavage agent, etc.

## 25 5.5 HOSTS

The present invention further provides host cells genetically engineered to contain the polynucleotides of the invention. For example, such host cells may contain nucleic acids of the invention introduced into the host cell using known transformation, transfection or infection methods. The present invention still further provides host cells genetically engineered to express the polynucleotides of the invention, wherein such polynucleotides are in operative association with a regulatory sequence heterologous to the host cell which drives expression of the polynucleotides in the cell.

Knowledge of prothrombinase-like DNA sequences allows for modification of cells to permit, or increase, expression of prothrombinase-like polypeptide. Cells can be modified (e.g., by homologous recombination) to provide increased prothrombinase-like polypeptide expression by replacing, in whole or in part, the naturally occurring prothrombinase-like promoter with all or part of a heterologous promoter so that the cells prothrombinase-like polypeptide is expressed at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to prothrombinase-like encoding sequences. See, for example, PCT International Publication No. WO94/12650, PCT International Publication No. WO92/20808, and PCT International Publication No. WO91/09955. It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (e.g., *ada*, *dhfr*, and the multifunctional *CAD* gene which encodes carbamyl phosphate synthase, aspartate transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the prothrombinase-like coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the prothrombinase-like coding sequences in the cells.

The host cell can be a higher eukaryotic host cell, such as a mammalian cell, a lower eukaryotic host cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. Introduction of the recombinant construct into the host cell can be effected by calcium phosphate transfection, DEAE, dextran-mediated transfection, or electroporation (Davis, L. et al., *Basic Methods in Molecular Biology* (1986)). The host cells containing one of the polynucleotides of the invention, can be used in conventional manners to produce the gene product encoded by the isolated fragment (in the case of an ORF) or can be used to produce a heterologous protein under the control of the EMF.

Any host/vector system can be used to express one or more of the ORFs of the present invention. These include, but are not limited to, eukaryotic hosts such as HeLa cells, Cv-1 cell, COS cells, 293 cells, and Sf9 cells, as well as prokaryotic host such as *E. coli* and *B. subtilis*. The most preferred cells are those which do not normally express the particular polypeptide or protein or which expresses the polypeptide or protein at low natural level. Mature proteins can be expressed in mammalian cells, yeast, bacteria, or other cells under the control of appropriate promoters. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention. Appropriate cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al., in *Molecular Cloning: A Laboratory Manual*, Second Edition,

Cold Spring Harbor, New York (1989), the disclosure of which is hereby incorporated by reference.

Various mammalian cell culture systems can also be employed to express recombinant protein. Examples of mammalian expression systems include the COS-7 lines of monkey  
5 kidney fibroblasts, described by Gluzman, Cell 23:175 (1981). Other cell lines capable of expressing a compatible vector are, for example, the C127, monkey COS cells, Chinese Hamster Ovary (CHO) cells, human kidney 293 cells, human epidermal A431 cells, human Colo205 cells, 3T3 cells, CV-1 cells, other transformed primate cell lines, normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, primary explants, HeLa cells,  
10 mouse L cells, BHK, HL-60, U937, HaK or Jurkat cells. Mammalian expression vectors will comprise an origin of replication, a suitable promoter and also any necessary ribosome binding sites, polyadenylation site, splice donor and acceptor sites, transcriptional termination sequences, and 5' flanking nontranscribed sequences. DNA sequences derived from the SV40 viral genome, for example, SV40 origin, early promoter, enhancer, splice, and  
15 polyadenylation sites may be used to provide the required nontranscribed genetic elements. Recombinant polypeptides and proteins produced in bacterial culture are usually isolated by initial extraction from cell pellets, followed by one or more salting-out, aqueous ion exchange or size exclusion chromatography steps. Protein refolding steps can be used, as necessary, in completing configuration of the mature protein. Finally, high performance liquid  
20 chromatography (HPLC) can be employed for final purification steps. Microbial cells employed in expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents.

Alternatively, it may be possible to produce the protein in lower eukaryotes such as yeast or insects or in prokaryotes such as bacteria. Potentially suitable yeast strains include  
25 *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Kluyveromyces* strains, *Candida*, or any yeast strain capable of expressing heterologous proteins. Potentially suitable bacterial strains include *Escherichia coli*, *Bacillus subtilis*, *Salmonella typhimurium*, or any bacterial strain capable of expressing heterologous proteins. If the protein is made in yeast or bacteria, it may be necessary to modify the protein produced therein, for example by phosphorylation or  
30 glycosylation of the appropriate sites, in order to obtain the functional protein. Such covalent attachments may be accomplished using known chemical or enzymatic methods.

In another embodiment of the present invention, cells and tissues may be engineered to express an endogenous gene comprising the polynucleotides of the invention under the control of inducible regulatory elements, in which case the regulatory sequences of the endogenous

gene may be replaced by homologous recombination. As described herein, gene targeting can be used to replace a gene's existing regulatory region with a regulatory sequence isolated from a different gene or a novel regulatory sequence synthesized by genetic engineering methods. Such regulatory sequences may be comprised of promoters, enhancers, scaffold-attachment  
5 regions, negative regulatory elements, transcriptional initiation sites, regulatory protein binding sites or combinations of said sequences. Alternatively, sequences which affect the structure or stability of the RNA or protein produced may be replaced, removed, added, or otherwise modified by targeting. These sequence include polyadenylation signals, mRNA stability elements, splice sites, leader sequences for enhancing or modifying transport or secretion  
10 properties of the protein, or other sequences which alter or improve the function or stability of protein or RNA molecules.

The targeting event may be a simple insertion of the regulatory sequence, placing the gene under the control of the new regulatory sequence, *e.g.*, inserting a new promoter or enhancer or both upstream of a gene. Alternatively, the targeting event may be a simple  
15 deletion of a regulatory element, such as the deletion of a tissue-specific negative regulatory element. Alternatively, the targeting event may replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally occurring elements. Here, the naturally occurring sequences are deleted and new sequences are added. In all cases, the identification of the targeting event may  
20 be facilitated by the use of one or more selectable marker genes that are contiguous with the targeting DNA, allowing for the selection of cells in which the exogenous DNA has integrated into the host cell genome. The identification of the targeting event may also be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively selectable marker is linked to the exogenous DNA, but configured such that the  
25 negatively selectable marker flanks the targeting sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result in the stable integration of the negatively selectable marker. Markers useful for this purpose include the Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial xanthine-guanine phosphoribosyl-transferase (gpt) gene.

30 The gene targeting or gene activation techniques which can be used in accordance with this aspect of the invention are more particularly described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.; International Application No. PCT/US92/09627 (WO93/09222) by Selden et al.; and International Application No.

PCT/US90/06436 (WO91/06667) by Skoultchi et al., each of which is incorporated by reference herein in its entirety.

## 5.6 POLYPEPTIDES OF THE INVENTION

5       The isolated polypeptides of the invention include, but are not limited to, a polypeptide comprising: the amino acid sequence set forth as any one of SEQ ID NO: 4-9, 14 or 16-18 or an amino acid sequence encoded by any one of the nucleotide sequences SEQ ID NO: 1-3, 10-13, or 15 or the corresponding full length or mature protein. Polypeptides of the invention also include polypeptides preferably with biological or immunological activity that are encoded by:

10   (a) a polynucleotide having any one of the nucleotide sequences set forth in the SEQ ID NO: 1-3, 10-13, or 15 or (b) polynucleotides encoding any one of the amino acid sequences set forth as SEQ ID NO: 4-9, 14 or 16-18 or (c) polynucleotides that hybridize to the complement of the polynucleotides of either (a) or (b) under stringent hybridization conditions. The invention also provides biologically active or immunologically active variants of any of the amino acid

15   sequences set forth as SEQ ID NO: 4-9, 14 or 16-18 or the corresponding full length or mature protein; and "substantial equivalents" thereof (e.g., with at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, typically at least about 95%, more typically at least about 98%, or most typically at least about 99% amino acid identity) that retain biological activity. Polypeptides encoded by allelic variants

20   may have a similar, increased, or decreased activity compared to polypeptides comprising SEQ ID NO: 4-9, 14 or 16-18.

Fragments of the proteins of the present invention which are capable of exhibiting biological activity are also encompassed by the present invention. Fragments of the protein may be in linear form or they may be cyclized using known methods, for example, as

25   described in H. U. Saragovi, et al., Bio/Technology 10, 773-778 (1992) and in R. S. McDowell, et al., J. Amer. Chem. Soc. 114, 9245-9253 (1992), both of which are incorporated herein by reference. Such fragments may be fused to carrier molecules such as immunoglobulins for many purposes, including increasing the valency of protein binding sites.

The present invention also provides both full-length and mature forms (for example,

30   without a signal sequence or precursor sequence) of the disclosed proteins. The protein coding sequence is identified in the sequence listing by translation of the disclosed nucleotide sequences. The mature form of such protein may be obtained by expression of a full-length polynucleotide in a suitable mammalian cell or other host cell. The sequence of the mature form of the protein is also determinable from the amino acid sequence of the full-length form.

Where proteins of the present invention are membrane bound, soluble forms of the proteins are also provided. In such forms, part or all of the regions causing the proteins to be membrane bound are deleted so that the proteins are fully secreted from the cell in which it is expressed.

Protein compositions of the present invention may further comprise an acceptable  
5 carrier, such as a hydrophilic, *e.g.*, pharmaceutically acceptable, carrier.

The present invention further provides isolated polypeptides encoded by the nucleic acid fragments of the present invention or by degenerate variants of the nucleic acid fragments of the present invention. By "degenerate variant" is intended nucleotide fragments which differ from a nucleic acid fragment of the present invention (*e.g.*, an ORF) by nucleotide sequence  
10 but, due to the degeneracy of the genetic code, encode an identical polypeptide sequence. Preferred nucleic acid fragments of the present invention are the ORFs that encode proteins.

A variety of methodologies known in the art can be utilized to obtain any one of the isolated polypeptides or proteins of the present invention. At the simplest level, the amino acid sequence can be synthesized using commercially available peptide synthesizers. The  
15 synthetically-constructed protein sequences, by virtue of sharing primary, secondary or tertiary structural and/or conformational characteristics with proteins may possess biological properties in common therewith, including protein activity. This technique is particularly useful in producing small peptides and fragments of larger polypeptides. Fragments are useful, for example, in generating antibodies against the native polypeptide. Thus, they may be employed  
20 as biologically active or immunological substitutes for natural, purified proteins in screening of therapeutic compounds and in immunological processes for the development of antibodies.

The polypeptides and proteins of the present invention can alternatively be purified from cells which have been altered to express the desired polypeptide or protein. As used herein, a cell is said to be altered to express a desired polypeptide or protein when the cell,  
25 through genetic manipulation, is made to produce a polypeptide or protein which it normally does not produce or which the cell normally produces at a lower level. One skilled in the art can readily adapt procedures for introducing and expressing either recombinant or synthetic sequences into eukaryotic or prokaryotic cells in order to generate a cell which produces one of the polypeptides or proteins of the present invention.

30 The invention also relates to methods for producing a polypeptide comprising growing a culture of host cells of the invention in a suitable culture medium, and purifying the protein from the cells or the culture in which the cells are grown. For example, the methods of the invention include a process for producing a polypeptide in which a host cell containing a suitable expression vector that includes a polynucleotide of the invention is cultured under



conditions that allow expression of the encoded polypeptide. The polypeptide can be recovered from the culture, conveniently from the culture medium, or from a lysate prepared from the host cells and further purified. Preferred embodiments include those in which the protein produced by such process is a full length or mature form of the protein.

5 In an alternative method, the polypeptide or protein is purified from bacterial cells which naturally produce the polypeptide or protein. One skilled in the art can readily follow known methods for isolating polypeptides and proteins in order to obtain one of the isolated polypeptides or proteins of the present invention. These include, but are not limited to, immunochromatography, HPLC, size-exclusion chromatography, ion-exchange  
10 chromatography, and immuno-affinity chromatography. See, *e.g.*, Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag (1994); Sambrook, et al., in *Molecular Cloning: A Laboratory Manual*; Ausubel et al., *Current Protocols in Molecular Biology*. Polypeptide fragments that retain biological/immunological activity include fragments comprising greater than about 100 amino acids, or greater than about 200 amino acids, and  
15 fragments that encode specific protein domains.

The purified polypeptides can be used in *in vitro* binding assays which are well known in the art to identify molecules which bind to the polypeptides. These molecules include but are not limited to, for *e.g.*, small molecules, molecules from combinatorial libraries, antibodies, or other proteins. The molecules identified in the binding assay are then tested for antagonist  
20 or agonist activity in *in vivo* tissue culture or animal models that are well known in the art. In brief, the molecules are titrated into a plurality of cell cultures or animals and then tested for either cell/animal death or prolonged survival of the animal/cells.

In addition, the peptides of the invention or molecules capable of binding to the peptides may be complexed with toxins, *e.g.*, ricin or cholera, or with other compounds that  
25 are toxic to cells. The toxin-binding molecule complex is then targeted to a tumor or other cell by the specificity of the binding molecule for SEQ ID NO: 4-9, 14 or 16-18.

The protein of the invention may also be expressed as a product of transgenic animals, *e.g.*, as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein.

30 The proteins provided herein also include proteins characterized by amino acid sequences similar to those of purified proteins but into which modification are naturally provided or deliberately engineered. For example, modifications, in the peptide or DNA sequence, can be made by those skilled in the art using known techniques. Modifications of interest in the protein sequences may include the alteration, substitution, replacement, insertion

or deletion of a selected amino acid residue in the coding sequence. For example, one or more of the cysteine residues may be deleted or replaced with another amino acid to alter the conformation of the molecule. Techniques for such alteration, substitution, replacement, insertion or deletion are well known to those skilled in the art (see, *e.g.*, U.S. Pat. No. 4,518,584). Preferably, such alteration, substitution, replacement, insertion or deletion retains the desired activity of the protein. Regions of the protein that are important for the protein function can be determined by various methods known in the art including the alanine-scanning method which involved systematic substitution of single or strings of amino acids with alanine, followed by testing the resulting alanine-containing variant for biological activity. This type of analysis determines the importance of the substituted amino acid(s) in biological activity. Regions of the protein that are important for protein function can be determined by the eMATRIX program.

Other fragments and derivatives of the sequences of proteins which would be expected to retain protein activity in whole or in part and are useful for screening or other immunological methodologies may also be easily made by those skilled in the art given the disclosures herein. Such modifications are encompassed by the present invention.

The protein may also be produced by operably linking the isolated polynucleotide of the invention to suitable control sequences in one or more insect expression vectors, and employing an insect expression system. Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, *e.g.*, Invitrogen, San Diego, Calif., U.S.A. (the MaxBat™ kit), and such methods are well known in the art, as described in Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987), incorporated herein by reference. As used herein, an insect cell capable of expressing a polynucleotide of the present invention is "transformed."

The protein of the invention may be prepared by culturing transformed host cells under culture conditions suitable to express the recombinant protein. The resulting expressed protein may then be purified from such culture (*i.e.*, from culture medium or cell extracts) using known purification processes, such as gel filtration and ion exchange chromatography. The purification of the protein may also include an affinity column containing agents which will bind to the protein; one or more column steps over such affinity resins as concanavalin A-agarose, heparin-toyopearl™ or Cibacrom blue 3GA Sepharose™; one or more steps involving hydrophobic interaction chromatography using such resins as phenyl ether, butyl ether, or propyl ether; or immunoaffinity chromatography.

Alternatively, the protein of the invention may also be expressed in a form which will facilitate purification. For example, it may be expressed as a fusion protein, such as those of maltose binding protein (MBP), glutathione-S-transferase (GST) or thioredoxin (TRX), or as a His tag. Kits for expression and purification of such fusion proteins are commercially  
5 available from New England BioLab (Beverly, Mass.), Pharmacia (Piscataway, N.J.) and Invitrogen, respectively. The protein can also be tagged with an epitope and subsequently purified by using a specific antibody directed to such epitope. One such epitope ("FLAG®") is commercially available from Kodak (New Haven, Conn.).

Finally, one or more reverse-phase high performance liquid chromatography (RP-  
10 HPLC) steps employing hydrophobic RP-HPLC media, *e.g.*, silica gel having pendant methyl or other aliphatic groups, can be employed to further purify the protein. Some or all of the foregoing purification steps, in various combinations, can also be employed to provide a substantially homogeneous isolated recombinant protein. The protein thus purified is substantially free of other mammalian proteins and is defined in accordance with the present  
15 invention as an "isolated protein."

The polypeptides of the invention include analogs (variants). The polypeptides of the invention include prothrombinase-like analogs. This embraces fragments of prothrombinase-like polypeptide of the invention, as well prothrombinase-like polypeptides which comprise one or more amino acids deleted, inserted, or substituted. Also, analogs of the prothrombinase-  
20 like polypeptide of the invention embrace fusions of the prothrombinase-like polypeptides or modifications of the prothrombinase-like polypeptides, wherein the prothrombinase-like polypeptide or analog is fused to another moiety or moieties, *e.g.*, targeting moiety or another therapeutic agent. Such analogs may exhibit improved properties such as activity and/or stability. Examples of moieties which may be fused to the prothrombinase-like polypeptide or  
25 an analog include, for example, targeting moieties which provide for the delivery of polypeptide to neurons, *e.g.*, antibodies to central nervous system, or antibodies to receptor and ligands expressed on neuronal cells. Other moieties which may be fused to prothrombinase-like polypeptide include therapeutic agents which are used for treatment, for example anti-depressant drugs or other medications for neurological disorders. Also,  
30 prothrombinase-like polypeptides may be fused to neuron growth modulators, and other chemokines for targeted delivery.

### 5.6.1 DETERMINING POLYPEPTIDE AND POLYNUCLEOTIDE IDENTITY AND SIMILARITY

Preferred identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in computer programs including, but are not limited to, the GCG program package, including GAP (Devereux, J., et al., Nucleic Acids Research 12(1):387 (1984); Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, BLASTX, FASTA (Altschul, S.F. et al., J. Molec. Biol. 215:403-410 (1990), PSI-BLAST (Altschul S.F. et al., Nucleic Acids Res. vol. 25, pp. 3389-3402, herein incorporated by reference), eMatrix software (Wu et al., J. Comp. Biol., vol. 6, pp. 219-235 (1999), herein incorporated by reference), eMotif software (Nevill-Manning et al, ISMB-97, vol 4, pp. 202-209, herein incorporated by reference) and the Kyte-Doolittle hydrophobicity prediction algorithm (J. Mol Biol, 157, pp. 105-31 (1982), incorporated herein by reference). The BLAST programs are publicly available from the National Center for Biotechnology Information (NCBI) and other sources (BLAST Manual, Altschul, S., et al. NCB NLM NIH Bethesda, MD 20894; Altschul, S., et al., J. Mol. Biol. 215:403-410 (1990).

### 5.7 CHIMERIC AND FUSION PROTEINS

The invention also provides chimeric or fusion proteins. As used herein, a "chimeric protein" or "fusion protein" comprises a polypeptide of the invention operatively linked to another polypeptide. Within a fusion protein the polypeptide according to the invention can correspond to all or a portion of a protein according to the invention. In one embodiment, a fusion protein comprises at least one biologically active portion of a protein according to the invention. In another embodiment, a fusion protein comprises at least two biologically active portions of a protein according to the invention. Within the fusion protein, the term "operatively linked" is intended to indicate that the polypeptide(s) according to the invention and the other polypeptide(s) are fused in-frame to each other. The polypeptide can be fused to the N-terminus or C-terminus or in the middle.

For example, in one embodiment a fusion protein comprises a polypeptide according to the invention operably linked to the extracellular domain of a second protein.

In another embodiment, the fusion protein is a GST-fusion protein in which the polypeptide sequences of the invention are fused to the C-terminus of the GST (i.e., glutathione S-transferase) sequences.

In another embodiment, the fusion protein is an immunoglobulin fusion protein in which the polypeptide sequences according to the invention comprise one or more domains

fused to sequences derived from a member of the immunoglobulin protein family. The immunoglobulin fusion proteins of the invention can be incorporated into pharmaceutical compositions and administered to a subject to inhibit an interaction between a ligand and a protein of the invention on the surface of a cell, to thereby suppress signal transduction *in vivo*.

5 The immunoglobulin fusion proteins can be used to affect the bioavailability of a cognate ligand. Inhibition of the ligand/protein interaction may be useful therapeutically for both the treatment of proliferative and differentiative disorders, *e.g.*, cancer as well as modulating (*e.g.*, promoting or inhibiting) cell survival. Moreover, the immunoglobulin fusion proteins of the invention can be used to bind and to dimerize 2 receptors and thereby transduce an  
10 intracellular signal. The immunoglobulin fusion proteins may also be used as immunogens to produce antibodies in a subject, to purify ligands, and in screening assays to identify molecules that inhibit the interaction of a polypeptide of the invention with a ligand.

A chimeric or fusion protein of the invention can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide  
15 sequences are ligated together in-frame in accordance with conventional techniques, *e.g.*, by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA  
20 synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers that give rise to complementary overhangs between two consecutive gene fragments that can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, Ausubel et al. (eds.) CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, 1992). Moreover, many expression vectors are commercially  
25 available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A nucleic acid encoding a polypeptide of the invention can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the protein of the invention.

## 5.8 GENE THERAPY

30 Mutations in the polynucleotides of the invention gene may result in loss of normal function of the encoded protein. The invention thus provides gene therapy to restore normal activity of the polypeptides of the invention; or to treat disease states involving polypeptides of the invention. Delivery of a functional gene encoding polypeptides of the invention to appropriate cells is effected *ex vivo*, *in situ*, or *in vivo* by use of vectors, and more particularly

viral vectors (e.g., adenovirus, adeno-associated virus, or a retrovirus), or *ex vivo* by use of physical DNA transfer methods (e.g., liposomes or chemical treatments). See, for example, Anderson, *Nature*, supplement to vol. 392, no. 6679, pp.25-20 (1998). For additional reviews of gene therapy technology see Friedmann, *Science*, 244: 1275-1281 (1989); Verma, *Scientific American*: 68-84 (1990); and Miller, *Nature*, 357: 455-460 (1992). Introduction of any one of the nucleotides of the present invention or a gene encoding the polypeptides of the present invention can also be accomplished with extrachromosomal substrates (transient expression) or artificial chromosomes (stable expression). Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes. Alternatively, it is contemplated that in other human disease states, preventing the expression of or inhibiting the activity of polypeptides of the invention will be useful in treating the disease states. It is contemplated that antisense therapy or gene therapy could be applied to negatively regulate the expression of polypeptides of the invention.

Other methods inhibiting expression of a protein include the introduction of antisense molecules to the nucleic acids of the present invention, their complements, or their translated RNA sequences, by methods known in the art. Further, the polypeptides of the present invention can be inhibited by using targeted deletion methods, or the insertion of a negative regulatory element such as a silencer, which is tissue specific.

The present invention still further provides cells genetically engineered *in vivo* to express the polynucleotides of the invention, wherein such polynucleotides are in operative association with a regulatory sequence heterologous to the host cell which drives expression of the polynucleotides in the cell. These methods can be used to increase or decrease the expression of the polynucleotides of the present invention.

Knowledge of DNA sequences provided by the invention allows for modification of cells to permit, increase, or decrease, expression of endogenous polypeptide. Cells can be modified (e.g., by homologous recombination) to provide increased polypeptide expression by replacing, in whole or in part, the naturally occurring promoter with all or part of a heterologous promoter so that the cells express the protein at higher levels. The heterologous promoter is inserted in such a manner that it is operatively linked to the desired protein encoding sequences. See, for example, PCT International Publication No. WO 94/12650, PCT International Publication No. WO 92/20808, and PCT International Publication No. WO 91/09955. It is also contemplated that, in addition to heterologous promoter DNA, amplifiable marker DNA (e.g., *ada*, *dhfr*, and the multifunctional CAD gene which encodes carbamyl phosphate synthase, aspartate

transcarbamylase, and dihydroorotase) and/or intron DNA may be inserted along with the heterologous promoter DNA. If linked to the desired protein coding sequence, amplification of the marker DNA by standard selection methods results in co-amplification of the desired protein coding sequences in the cells.

5 In another embodiment of the present invention, cells and tissues may be engineered to express an endogenous gene comprising the polynucleotides of the invention under the control of inducible regulatory elements, in which case the regulatory sequences of the endogenous gene may be replaced by homologous recombination. As described herein, gene targeting can be used to replace a gene's existing regulatory region with a regulatory sequence isolated from a different  
10 gene or a novel regulatory sequence synthesized by genetic engineering methods. Such regulatory sequences may be comprised of promoters, enhancers, scaffold-attachment regions, negative regulatory elements, transcriptional initiation sites, regulatory protein binding sites or combinations of said sequences. Alternatively, sequences which affect the structure or stability of the RNA or protein produced may be replaced, removed, added, or otherwise modified by  
15 targeting. These sequences include polyadenylation signals, mRNA stability elements, splice sites, leader sequences for enhancing or modifying transport or secretion properties of the protein, or other sequences which alter or improve the function or stability of protein or RNA molecules.

The targeting event may be a simple insertion of the regulatory sequence, placing the gene under the control of the new regulatory sequence, *e.g.*, inserting a new promoter or enhancer or  
20 both upstream of a gene. Alternatively, the targeting event may be a simple deletion of a regulatory element, such as the deletion of a tissue-specific negative regulatory element. Alternatively, the targeting event may replace an existing element; for example, a tissue-specific enhancer can be replaced by an enhancer that has broader or different cell-type specificity than the naturally occurring elements. Here, the naturally occurring sequences are deleted and new  
25 sequences are added. In all cases, the identification of the targeting event may be facilitated by the use of one or more selectable marker genes that are contiguous with the targeting DNA, allowing for the selection of cells in which the exogenous DNA has integrated into the cell genome. The identification of the targeting event may also be facilitated by the use of one or more marker genes exhibiting the property of negative selection, such that the negatively  
30 selectable marker is linked to the exogenous DNA, but configured such that the negatively selectable marker flanks the targeting sequence, and such that a correct homologous recombination event with sequences in the host cell genome does not result in the stable integration of the negatively selectable marker. Markers useful for this purpose include the

Herpes Simplex Virus thymidine kinase (TK) gene or the bacterial xanthine-guanine phosphoribosyl-transferase (gpt) gene.

The gene targeting or gene activation techniques which can be used in accordance with this aspect of the invention are more particularly described in U.S. Patent No. 5,272,071 to Chappel; U.S. Patent No. 5,578,461 to Sherwin et al.; International Application No. PCT/US92/09627 (WO93/09222) by Selden et al.; and International Application No. PCT/US90/06436 (WO91/06667) by Skoultchi et al., each of which is incorporated by reference herein in its entirety.

## 10 5.9 TRANSGENIC ANIMALS

In preferred methods to determine biological functions of the polypeptides of the invention *in vivo*, one or more genes provided by the invention are either over expressed or inactivated in the germ line of animals using homologous recombination [Capecchi, Science 244:1288-1292 (1989)]. Animals in which the gene is over expressed, under the regulatory control of exogenous or endogenous promoter elements, are known as transgenic animals. Animals in which an endogenous gene has been inactivated by homologous recombination are referred to as "knockout" animals. Knockout animals, preferably non-human mammals, can be prepared as described in U.S. Patent No. 5,557,032, incorporated herein by reference. Transgenic animals are useful to determine the roles polypeptides of the invention play in biological processes, and preferably in disease states. Transgenic animals are useful as model systems to identify compounds that modulate lipid metabolism. Transgenic animals, preferably non-human mammals, are produced using methods as described in U.S. Patent No 5,489,743 and PCT Publication No. WO94/28122, incorporated herein by reference.

Transgenic animals can be prepared wherein all or part of a promoter of the polynucleotides of the invention is either activated or inactivated to alter the level of expression of the polypeptides of the invention. Inactivation can be carried out using homologous recombination methods described above. Activation can be achieved by supplementing or even replacing the homologous promoter to provide for increased protein expression. The homologous promoter can be supplemented by insertion of one or more heterologous enhancer elements known to confer promoter activation in a particular tissue.

The polynucleotides of the present invention also make possible the development, through, e.g., homologous recombination or knock out strategies, of animals that fail to express functional prothrombinase-like polypeptide or that express a variant of prothrombinase-like polypeptide. Such animals are useful as models for studying the *in vivo* activities of



prothrombinase-like polypeptide as well as for studying modulators of the prothrombinase-like polypeptide.

In preferred methods to determine biological functions of the polypeptides of the invention *in vivo*, one or more genes provided by the invention are either over expressed or  
5 inactivated in the germ line of animals using homologous recombination [Capecchi, Science 244:1288-1292 (1989)]. Animals in which the gene is over expressed, under the regulatory control of exogenous or endogenous promoter elements, are known as transgenic animals. Animals in which an endogenous gene has been inactivated by homologous recombination are referred to as "knockout" animals. Knockout animals, preferably non-human mammals, can be  
10 prepared as described in U.S. Patent No. 5,557,032, incorporated herein by reference. Transgenic animals are useful to determine the roles polypeptides of the invention play in biological processes, and preferably in disease states. Transgenic animals are useful as model systems to identify compounds that modulate lipid metabolism. Transgenic animals, preferably non-human mammals, are produced using methods as described in U.S. Patent No 5,489,743  
15 and PCT Publication No. WO94/28122, incorporated herein by reference.

Transgenic animals can be prepared wherein all or part of the polynucleotides of the invention promoter is either activated or inactivated to alter the level of expression of the polypeptides of the invention. Inactivation can be carried out using homologous recombination  
20 methods described above. Activation can be achieved by supplementing or even replacing the homologous promoter to provide for increased protein expression. The homologous promoter can be supplemented by insertion of one or more heterologous enhancer elements known to confer promoter activation in a particular tissue.

## 5.10 USES AND BIOLOGICAL ACTIVITY OF PROTHROMBINASE-LIKE 25 POLYPEPTIDE

The polynucleotides and proteins of the present invention are expected to exhibit one or more of the uses or biological activities (including those associated with assays cited herein) identified herein. Uses or activities described for proteins of the present invention may be provided by administration or use of such proteins or of polynucleotides encoding such  
30 proteins (such as, for example, in gene therapies or vectors suitable for introduction of DNA). The mechanism underlying the particular condition or pathology will dictate whether the polypeptides of the invention, the polynucleotides of the invention or modulators (activators or inhibitors) thereof would be beneficial to the subject in need of treatment. Thus, "therapeutic compositions of the invention" include compositions comprising isolated polynucleotides

(including recombinant DNA molecules, cloned genes and degenerate variants thereof) or polypeptides of the invention (including full length protein, mature protein and truncations or domains thereof), or compounds and other substances that modulate the overall activity of the target gene products, either at the level of target gene/protein expression or target protein activity. Such modulators include polypeptides, analogs, (variants), including fragments and fusion proteins, antibodies and other binding proteins; chemical compounds that directly or indirectly activate or inhibit the polypeptides of the invention (identified, e.g., via drug screening assays as described herein); antisense polynucleotides and polynucleotides suitable for triple helix formation; and in particular antibodies or other binding partners that specifically recognize one or more epitopes of the polypeptides of the invention.

The polypeptides of the present invention may likewise be involved in cellular activation or in one of the other physiological pathways described herein.

#### 5.10.1 RESEARCH USES AND UTILITIES

The polynucleotides provided by the present invention can be used by the research community for various purposes. The polynucleotides can be used to express recombinant protein for analysis, characterization or therapeutic use; as markers for tissues in which the corresponding protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in disease states); as molecular weight markers on gels; as chromosome markers or tags (when labeled) to identify chromosomes or to map related gene positions; to compare with endogenous DNA sequences in patients to identify potential genetic disorders; as probes to hybridize and thus discover novel, related DNA sequences; as a source of information to derive PCR primers for genetic fingerprinting; as a probe to "subtract-out" known sequences in the process of discovering other novel polynucleotides; for selecting and making oligomers for attachment to a "gene chip" or other support, including for examination of expression patterns; to raise anti-protein antibodies using DNA immunization techniques; and as an antigen to raise anti-DNA antibodies or elicit another immune response. Where the polynucleotide encodes a protein which binds or potentially binds to another protein (such as, for example, in a receptor-ligand interaction), the polynucleotide can also be used in interaction trap assays (such as, for example, that described in Gyuris et al., Cell 75:791-803 (1993)) to identify polynucleotides encoding the other protein with which binding occurs or to identify inhibitors of the binding interaction.

The polypeptides provided by the present invention can similarly be used in assays to determine biological activity, including in a panel of multiple proteins for high-throughput

screening; to raise antibodies or to elicit another immune response; as a reagent (including the labeled reagent) in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids; as markers for tissues in which the corresponding polypeptide is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state); and, of course, to isolate correlative receptors or ligands. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

The polypeptides of the invention are also useful for making antibody substances that are specifically immunoreactive with prothrombinase-like proteins. Antibodies and portions thereof (e.g., Fab fragments) which bind to the polypeptides of the invention can be used to identify the presence of such polypeptides in a sample. Such determinations are carried out using any suitable immunoassay format, and any polypeptide of the invention that is specifically bound by the antibody can be employed as a positive control.

Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products.

Methods for performing the uses listed above are well known to those skilled in the art. References disclosing such methods include without limitation "Molecular Cloning: A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory Press, Sambrook, J., E. F. Fritsch and T. Maniatis eds., 1989, and "Methods in Enzymology: Guide to Molecular Cloning Techniques", Academic Press, Berger, S. L. and A. R. Kimmel eds., 1987.

### 5.10.2 NUTRITIONAL USES

Polynucleotides and polypeptides of the present invention can also be used as nutritional sources or supplements. Such uses include without limitation use as a protein or amino acid supplement, use as a carbon source, use as a nitrogen source and use as a source of carbohydrate. In such cases the polypeptide or polynucleotide of the invention can be added to the feed of a particular organism or can be administered as a separate solid or liquid preparation, such as in the form of powder, pills, solutions, suspensions or capsules. In the case of microorganisms, the polypeptide or polynucleotide of the invention can be added to the medium in or on which the microorganism is cultured.

Additionally, the polypeptides of the invention can be used as molecular weight markers, and as a food supplement. A polypeptide consisting of SEQ ID NO: 4, for example, has a molecular mass of approximately 43 kDa in its unprocessed and unglycosylated state. Protein

food supplements are well known and the formulation of suitable food supplements including polypeptides of the invention is within the level of skill in the food preparation art.

### 5.10.3 CYTOKINE AND CELL PROLIFERATION/DIFFERENTIATION

#### 5 ACTIVITY

A polypeptide of the present invention may exhibit activity relating to cytokine, cell proliferation (either inducing or inhibiting) or cell differentiation (either inducing or inhibiting) activity or may induce production of other cytokines in certain cell populations. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays, and hence the assays serve as a convenient confirmation of cytokine activity. The activity of therapeutic compositions of the present invention is evidenced by any one of a number of routine factor dependent cell proliferation assays for cell lines including, without limitation, 32D, DA2, DA1G, T10, B9, B9/11, BaF3, MC9/G, M+(preB M+), 2E8, RB5, DA1, 123, T1165, HT2, CTLL2, TF-1, Mo7e, CMK, HUVEC, and Caco. Therapeutic compositions of the invention can be used in the following:

Assays for T-cell or thymocyte proliferation include without limitation those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Bertagnolli et al., J. Immunol. 145:1706-1712, 1990; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Bertagnolli, et al., I. Immunol. 149:3778-3783, 1992; Bowman et al., I. Immunol. 152:1756-1761, 1994.

Assays for cytokine production and/or proliferation of spleen cells, lymph node cells or thymocytes include, without limitation, those described in: Polyclonal T cell stimulation, Kruisbeek, A. M. and Shevach, E. M. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 3.12.1-3.12.14, John Wiley and Sons, Toronto. 1994; and Measurement of mouse and human interleukin- $\gamma$ , Schreiber, R. D. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.8.1-6.8.8, John Wiley and Sons, Toronto. 1994.

Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include, without limitation, those described in: Measurement of Human and Murine Interleukin 2 and Interleukin 4, Bottomly, K., Davis, L. S. and Lipsky, P. E. In Current Protocols in

- Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 6.3.1-6.3.12, John Wiley and Sons, Toronto. 1991; deVries et al., J. Exp. Med. 173:1205-1211, 1991; Moreau et al., Nature 336:690-692, 1988; Greenberger et al., Proc. Natl. Acad. Sci. U.S.A. 80:2931-2938, 1983; Measurement of mouse and human interleukin 6--Nordan, R. In Current Protocols in Immunology. J. E. Coligan eds. Vol 1 pp. 6.6.1-6.6.5, John Wiley and Sons, Toronto. 1991; Smith et al., Proc. Natl. Acad. Sci. U.S.A. 83:1857-1861, 1986; Measurement of human Interleukin 11--Bennett, F., Giannotti, J., Clark, S. C. and Turner, K. J. In Current Protocols in Immunology. J. E. Coligan eds. Vol 1 pp. 6.15.1 John Wiley and Sons, Toronto. 1991; Measurement of mouse and human Interleukin 9-Ciarletta, A., Giannotti, J., Clark, S. C. and Turner, K. J. In Current Protocols in Immunology. J. E. Coligan eds. Vol 1 pp. 6.13.1, John Wiley and Sons, Toronto. 1991.

- Assays for T-cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T-cell effects by measuring proliferation and cytokine production) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, *In Vitro* assays for Mouse Lymphocyte Function; Chapter 6, Cytokines and their cellular receptors; Chapter 7, Immunologic studies in Humans); Weinberger et al., Proc. Natl. Acad. Sci. USA 77:6091-6095, 1980; Weinberger et al., Eur. J. Immun. 11:405-411, 1981; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988.

#### 5.10.4 STEM CELL GROWTH FACTOR ACTIVITY

- A polypeptide of the present invention may exhibit stem cell growth factor activity and be involved in the proliferation, differentiation and survival of pluripotent and totipotent stem cells including primordial germ cells, embryonic stem cells, hematopoietic stem cells and/or germ line stem cells. Administration of the polypeptide of the invention to stem cells *in vivo* or *ex vivo* may maintain and expand cell populations in a totipotential or pluripotential state which would be useful for re-engineering damaged or diseased tissues, transplantation, manufacture of bio-pharmaceuticals and the development of bio-sensors. The ability to produce large quantities of human cells has important working applications for the production of human proteins which currently must be obtained from non-human sources or donors, implantation of cells to treat diseases such as Parkinson's, Alzheimer's and other neurodegenerative diseases; tissues for grafting such as bone marrow, skin, cartilage, tendons, bone, muscle (including cardiac muscle), blood vessels, cornea, neural cells, gastrointestinal

cells and others; and organs for transplantation such as kidney, liver, pancreas (including islet cells), heart and lung.

It is contemplated that multiple different exogenous growth factors and/or cytokines may be administered in combination with the polypeptide of the invention to achieve the  
5 desired effect, including any of the growth factors listed herein, other stem cell maintenance factors, and specifically including stem cell factor (SCF), leukemia inhibitory factor (LIF), Flt-3 ligand (Flt-3L), any of the interleukins, recombinant soluble IL-6 receptor fused to IL-6, macrophage inflammatory protein 1-alpha (MIP-1-alpha), G-CSF, GM-CSF, thrombopoietin (TPO), platelet factor 4 (PF-4), platelet-derived growth factor (PDGF), neural growth factors  
10 and basic fibroblast growth factor (bFGF).

Since totipotent stem cells can give rise to virtually any mature cell type, expansion of these cells in culture will facilitate the production of large quantities of mature cells. Techniques for culturing stem cells are known in the art and administration of polypeptides of the invention, optionally with other growth factors and/or cytokines, is expected to enhance the  
15 survival and proliferation of the stem cell populations. This can be accomplished by direct administration of the polypeptide of the invention to the culture medium. Alternatively, stroma cells transfected with a polynucleotide that encodes for the polypeptide of the invention can be used as a feeder layer for the stem cell populations in culture or in vivo. Stromal support cells for feeder layers may include embryonic bone marrow fibroblasts, bone marrow stromal cells,  
20 fetal liver cells, or cultured embryonic fibroblasts (see U.S. Patent No. 5,690,926).

Stem cells themselves can be transfected with a polynucleotide of the invention to induce autocrine expression of the polypeptide of the invention. This will allow for generation of undifferentiated totipotent/pluripotent stem cell lines that are useful as is or that can then be differentiated into the desired mature cell types. These stable cell lines can also serve  
25 as a source of undifferentiated totipotent/pluripotent mRNA to create cDNA libraries and templates for polymerase chain reaction experiments. These studies would allow for the isolation and identification of differentially expressed genes in stem cell populations that regulate stem cell proliferation and/or maintenance.

Expansion and maintenance of totipotent stem cell populations will be useful in the  
30 treatment of many pathological conditions. For example, polypeptides of the present invention may be used to manipulate stem cells in culture to give rise to neuroepithelial cells that can be used to augment or replace cells damaged by illness, autoimmune disease, accidental damage or genetic disorders. The polypeptide of the invention may be useful for inducing the proliferation of neural cells and for the regeneration of nerve and brain tissue, i.e. for the

treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders which involve degeneration, death or trauma to neural cells or nerve tissue. In addition, the expanded stem cell populations can also be genetically altered for gene therapy purposes and to decrease host rejection of replacement tissues after grafting or  
5 implantation.

Expression of the polypeptide of the invention and its effect on stem cells can also be manipulated to achieve controlled differentiation of the stem cells into more differentiated cell types. A broadly applicable method of obtaining pure populations of a specific differentiated cell type from undifferentiated stem cell populations involves the use of a cell-type specific  
10 promoter driving a selectable marker. The selectable marker allows only cells of the desired type to survive. For example, stem cells can be induced to differentiate into cardiomyocytes (Wobus et al., Differentiation, 48: 173-182, (1991); Klug et al., J. Clin. Invest., 98(1): 216-224, (1998)) or skeletal muscle cells (Browder, L. W. In: *Principles of Tissue Engineering* eds. Lanza et al., Academic Press (1997)). Alternatively, directed differentiation of stem cells  
15 can be accomplished by culturing the stem cells in the presence of a differentiation factor such as retinoic acid and an antagonist of the polypeptide of the invention which would inhibit the effects of endogenous stem cell factor activity and allow differentiation to proceed.

*In vitro* cultures of stem cells can be used to determine if the polypeptide of the invention exhibits stem cell growth factor activity. Stem cells are isolated from any one of  
20 various cell sources (including hematopoietic stem cells and embryonic stem cells) and cultured on a feeder layer, as described by Thompson et al. Proc. Natl. Acad. Sci, U.S.A., 92: 7844-7848 (1995), in the presence of the polypeptide of the invention alone or in combination with other growth factors or cytokines. The ability of the polypeptide of the invention to induce stem cells proliferation is determined by colony formation on semi-solid support e.g. as  
25 described by Bernstein et al., Blood, 77: 2316-2321 (1991).

#### 5.10.5 HEMATOPOIESIS REGULATING ACTIVITY

A polypeptide of the present invention may be involved in regulation of hematopoiesis and, consequently, in the treatment of myeloid or lymphoid cell disorders. Even marginal  
30 biological activity in support of colony forming cells or of factor-dependent cell lines indicates involvement in regulating hematopoiesis, e.g. in supporting the growth and proliferation of erythroid progenitor cells alone or in combination with other cytokines, thereby indicating utility, for example, in treating various anemias or for use in conjunction with irradiation/chemotherapy to stimulate the production of erythroid precursors and/or erythroid

cells; in supporting the growth and proliferation of myeloid cells such as granulocytes and monocytes/macrophages (i.e., traditional CSF activity) useful, for example, in conjunction with chemotherapy to prevent or treat consequent myelo-suppression; in supporting the growth and proliferation of megakaryocytes and consequently of platelets thereby allowing prevention  
5 or treatment of various platelet disorders such as thrombocytopenia, and generally for use in place of or complimentary to platelet transfusions; and/or in supporting the growth and proliferation of hematopoietic stem cells which are capable of maturing to any and all of the above-mentioned hematopoietic cells and therefore find therapeutic utility in various stem cell disorders (such as those usually treated with transplantation, including, without limitation,  
10 aplastic anemia and paroxysmal nocturnal hemoglobinuria), as well as in repopulating the stem cell compartment post irradiation/chemotherapy, either *in-vivo* or *ex-vivo* (i.e., in conjunction with bone marrow transplantation or with peripheral progenitor cell transplantation (homologous or heterologous)) as normal cells or genetically manipulated for gene therapy.

Therapeutic compositions of the invention can be used in the following:

15 Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above.

Assays for embryonic stem cell differentiation (which will identify, among others, proteins that influence embryonic differentiation hematopoiesis) include, without limitation, those described in: Johansson et al. Cellular Biology 15:141-151, 1995; Keller et al.,  
20 Molecular and Cellular Biology 13:473-486, 1993; McClanahan et al., Blood 81:2903-2915, 1993.

Assays for stem cell survival and differentiation (which will identify, among others, proteins that regulate lympho-hematopoiesis) include, without limitation, those described in: Methylcellulose colony forming assays, Freshney, M. G. In Culture of Hematopoietic Cells.  
25 R. I. Freshney, et al. eds. Vol pp. 265-268, Wiley-Liss, Inc., New York, N.Y. 1994; Hirayama et al., Proc. Natl. Acad. Sci. USA 89:5907-5911, 1992; Primitive hematopoietic colony forming cells with high proliferative potential, McNiece, I. K. and Briddell, R. A. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 23-39, Wiley-Liss, Inc., New York, N.Y. 1994; Neben et al., Experimental Hematology 22:353-359, 1994;  
30 Cobblestone area forming cell assay, Ploemacher, R. E. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 1-21, Wiley-Liss, Inc., New York, N.Y. 1994; Long term bone marrow cultures in the presence of stromal cells, Spooncer, E., Dexter, M. and Allen, T. In Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 163-179, Wiley-Liss, Inc., New York, N.Y. 1994; Long term culture initiating cell assay, Sutherland, H. J. In



Culture of Hematopoietic Cells. R. I. Freshney, et al. eds. Vol pp. 139-162, Wiley-Liss, Inc., New York, N.Y. 1994.

#### 5.10.6 TISSUE GROWTH ACTIVITY

5 A polypeptide of the present invention also may be involved in bone, cartilage, tendon, ligament and/or nerve tissue growth or regeneration, as well as in wound healing and tissue repair and replacement, and in healing of burns, incisions and ulcers.

A polypeptide of the present invention which induces cartilage and/or bone growth in circumstances where bone is not normally formed, has application in the healing of bone  
10 fractures and cartilage damage or defects in humans and other animals. Compositions of a polypeptide, antibody, binding partner, or other modulator of the invention may have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. De novo bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma induced, or oncologic resection induced craniofacial defects, and  
15 also is useful in cosmetic plastic surgery.

A polypeptide of this invention may also be involved in attracting bone-forming cells, stimulating growth of bone-forming cells, or inducing differentiation of progenitors of bone-forming cells. Treatment of osteoporosis, osteoarthritis, bone degenerative disorders, or  
periodontal disease, such as through stimulation of bone and/or cartilage repair or by blocking  
20 inflammation or processes of tissue destruction (collagenase activity, osteoclast activity, etc.) mediated by inflammatory processes may also be possible using the composition of the invention.

Another category of tissue regeneration activity that may involve the polypeptide of the present invention is tendon/ligament formation. Induction of tendon/ligament-like tissue or  
25 other tissue formation in circumstances where such tissue is not normally formed, has application in the healing of tendon or ligament tears, deformities and other tendon or ligament defects in humans and other animals. Such a preparation employing a tendon/ligament-like tissue inducing protein may have prophylactic use in preventing damage to tendon or ligament tissue, as well as use in the improved fixation of tendon or ligament to bone or other tissues,  
30 and in repairing defects to tendon or ligament tissue. De novo tendon/ligament-like tissue formation induced by a composition of the present invention contributes to the repair of congenital, trauma induced, or other tendon or ligament defects of other origin, and is also useful in cosmetic plastic surgery for attachment or repair of tendons or ligaments. The compositions of the present invention may provide environment to attract tendon- or ligament-

forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo* for return *in vivo* to effect tissue repair. The compositions of the invention may also be useful in the treatment of tendinitis, carpal tunnel syndrome and other tendon or  
5 ligament defects. The compositions may also include an appropriate matrix and/or sequestering agent as a carrier as is well known in the art.

The compositions of the present invention may also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, i.e. for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic  
10 disorders, which involve degeneration, death or trauma to neural cells or nerve tissue. More specifically, a composition may be used in the treatment of diseases of the peripheral nervous system, such as peripheral nerve injuries, peripheral neuropathy and localized neuropathies, and central nervous system diseases, such as Alzheimer's, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Further conditions which  
15 may be treated in accordance with the present invention include mechanical and traumatic disorders, such as spinal cord disorders, head trauma and cerebrovascular diseases such as stroke. Peripheral neuropathies resulting from chemotherapy or other medical therapies may also be treatable using a composition of the invention.

Compositions of the invention may also be useful to promote better or faster closure of  
20 non-healing wounds, including without limitation pressure ulcers, ulcers associated with vascular insufficiency, surgical and traumatic wounds, and the like.

Compositions of the present invention may also be involved in the generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac) and vascular (including  
25 vascular endothelium) tissue, or for promoting the growth of cells comprising such tissues. Part of the desired effects may be by inhibition or modulation of fibrotic scarring may allow normal tissue to regenerate. A polypeptide of the present invention may also exhibit angiogenic activity.

A composition of the present invention may also be useful for gut protection or  
30 regeneration and treatment of lung or liver fibrosis, reperfusion injury in various tissues, and conditions resulting from systemic cytokine damage.

A composition of the present invention may also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells; or for inhibiting the growth of tissues described above.

Therapeutic compositions of the invention can be used in the following:

Assays for tissue generation activity include, without limitation, those described in: International Patent Publication No. WO95/16035 (bone, cartilage, tendon); International Patent Publication No. WO95/05846 (nerve, neuronal); International Patent Publication No. WO91/07491 (skin, endothelium).

Assays for wound healing activity include, without limitation, those described in: Winter, Epidermal Wound Healing, pp. 71-112 (Maibach, H. I. and Rovee, D. T., eds.), Year Book Medical Publishers, Inc., Chicago, as modified by Eaglstein and Mertz, J. Invest. Dermatol 71:382-84 (1978).

#### 5.10.7 IMMUNE STIMULATING OR SUPPRESSING ACTIVITY

A polypeptide of the present invention may also exhibit immune stimulating or immune suppressing activity, including without limitation the activities for which assays are described herein. A polynucleotide of the invention can encode a polypeptide exhibiting such activities.

A protein may be useful in the treatment of various immune deficiencies and disorders (including severe combined immunodeficiency (SCID)), e.g., in regulating (up or down) growth and proliferation of T and/or B lymphocytes, as well as effecting the cytolytic activity of NK cells and other cell populations. These immune deficiencies may be genetic or be caused by viral (e.g., HIV) as well as bacterial or fungal infections, or may result from autoimmune disorders. More specifically, infectious diseases caused by viral, bacterial, fungal or other infection may be treatable using a protein of the present invention, including infections by HIV, hepatitis viruses, herpes viruses, mycobacteria, Leishmania spp., malaria spp. and various fungal infections such as candidiasis. Of course, in this regard, proteins of the present invention may also be useful where a boost to the immune system generally may be desirable, i.e., in the treatment of cancer.

Autoimmune disorders which may be treated using a protein of the present invention include, for example, connective tissue disease, multiple sclerosis, systemic lupus erythematosus, rheumatoid arthritis, autoimmune pulmonary inflammation, Guillain-Barre syndrome, autoimmune thyroiditis, insulin dependent diabetes mellitus, myasthenia gravis, graft-versus-host disease and autoimmune inflammatory eye disease. Such a protein (or antagonists thereof, including antibodies) of the present invention may also be useful in the treatment of allergic reactions and conditions (e.g., anaphylaxis, serum sickness, drug reactions, food allergies, insect venom allergies, mastocytosis, allergic rhinitis, hypersensitivity pneumonitis, urticaria, angioedema, eczema, atopic dermatitis, allergic contact

dermatitis, erythema multiforme, Stevens-Johnson syndrome, allergic conjunctivitis, atopic keratoconjunctivitis, venereal keratoconjunctivitis, giant papillary conjunctivitis and contact allergies), such as asthma (particularly allergic asthma) or other respiratory problems. Other conditions, in which immune suppression is desired (including, for example, organ  
5 transplantation), may also be treatable using a protein (or antagonists thereof) of the present invention. The therapeutic effects of the polypeptides or antagonists thereof on allergic reactions can be evaluated by in vivo animals models such as the cumulative contact enhancement test (Lastbom et al., Toxicology 125: 59-66, 1998), skin prick test (Hoffmann et al., Allergy 54: 446-54, 1999), guinea pig skin sensitization test (Vohr et al., Arch. Toxicol.  
10 73: 501-9), and murine local lymph node assay (Kimber et al., J. Toxicol. Environ. Health 53: 563-79).

Using the proteins of the invention it may also be possible to modulate immune responses, in a number of ways. Down regulation may be in the form of inhibiting or blocking an immune response already in progress or may involve preventing the induction of an immune  
15 response. The functions of activated T cells may be inhibited by suppressing T cell responses or by inducing specific tolerance in T cells, or both. Immunosuppression of T cell responses is generally an active, non-antigen-specific, process which requires continuous exposure of the T cells to the suppressive agent. Tolerance, which involves inducing non-responsiveness or  
20 anergy in T cells, is distinguishable from immunosuppression in that it is generally antigen-specific and persists after exposure to the tolerizing agent has ceased. Operationally, tolerance can be demonstrated by the lack of a T cell response upon reexposure to specific antigen in the absence of the tolerizing agent.

Down regulating or preventing one or more antigen functions (including without limitation B lymphocyte antigen functions (such as, for example, B7)), e.g., preventing high  
25 level lymphokine synthesis by activated T cells, will be useful in situations of tissue, skin and organ transplantation and in graft-versus-host disease (GVHD). For example, blockage of T cell function should result in reduced tissue destruction in tissue transplantation. Typically, in tissue transplants, rejection of the transplant is initiated through its recognition as foreign by T cells, followed by an immune reaction that destroys the transplant. The administration of a  
30 therapeutic composition of the invention may prevent cytokine synthesis by immune cells, such as T cells, and thus acts as an immunosuppressant. Moreover, a lack of costimulation may also be sufficient to anergize the T cells, thereby inducing tolerance in a subject. Induction of long-term tolerance by B lymphocyte antigen-blocking reagents may avoid the necessity of repeated administration of these blocking reagents. To achieve sufficient immunosuppression or

tolerance in a subject, it may also be necessary to block the function of a combination of B lymphocyte antigens.

The efficacy of particular therapeutic compositions in preventing organ transplant rejection or GVHD can be assessed using animal models that are predictive of efficacy in humans. Examples of appropriate systems which can be used include allogeneic cardiac grafts in rats and xenogeneic pancreatic islet cell grafts in mice, both of which have been used to examine the immunosuppressive effects of CTLA4Ig fusion proteins in vivo as described in Lenschow et al., *Science* 257:789-792 (1992) and Turka et al., *Proc. Natl. Acad. Sci USA*, 89:11102-11105 (1992). In addition, murine models of GVHD (see Paul ed., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 846-847) can be used to determine the effect of therapeutic compositions of the invention on the development of that disease.

Blocking antigen function may also be therapeutically useful for treating autoimmune diseases. Many autoimmune disorders are the result of inappropriate activation of T cells that are reactive against self tissue and which promote the production of cytokines and autoantibodies involved in the pathology of the diseases. Preventing the activation of autoreactive T cells may reduce or eliminate disease symptoms. Administration of reagents which block stimulation of T cells can be used to inhibit T cell activation and prevent production of autoantibodies or T cell-derived cytokines which may be involved in the disease process. Additionally, blocking reagents may induce antigen-specific tolerance of autoreactive T cells which could lead to long-term relief from the disease. The efficacy of blocking reagents in preventing or alleviating autoimmune disorders can be determined using a number of well-characterized animal models of human autoimmune diseases. Examples include murine experimental autoimmune encephalitis, systemic lupus erythematosus in MRL/lpr/lpr mice or NZB hybrid mice, murine autoimmune collagen arthritis, diabetes mellitus in NOD mice and BB rats, and murine experimental myasthenia gravis (see Paul ed., *Fundamental Immunology*, Raven Press, New York, 1989, pp. 840-856).

Upregulation of an antigen function (e.g., a B lymphocyte antigen function), as a means of up regulating immune responses, may also be useful in therapy. Upregulation of immune responses may be in the form of enhancing an existing immune response or eliciting an initial immune response. For example, enhancing an immune response may be useful in cases of viral infection, including systemic viral diseases such as influenza, the common cold, and encephalitis.

Alternatively, anti-viral immune responses may be enhanced in an infected patient by removing T cells from the patient, costimulating the T cells in vitro with viral antigen-pulsed

APCs either expressing a peptide of the present invention or together with a stimulatory form of a soluble peptide of the present invention and reintroducing the in vitro activated T cells into the patient. Another method of enhancing anti-viral immune responses would be to isolate infected cells from a patient, transfect them with a nucleic acid encoding a protein of the present invention as described herein such that the cells express all or a portion of the protein on their surface, and reintroduce the transfected cells into the patient. The infected cells would now be capable of delivering a costimulatory signal to, and thereby activate, T cells in vivo.

A polypeptide of the present invention may provide the necessary stimulation signal to T cells to induce a T cell mediated immune response against the transfected tumor cells. In addition, tumor cells which lack MHC class I or MHC class II molecules, or which fail to reexpress sufficient mounts of MHC class I or MHC class II molecules, can be transfected with nucleic acid encoding all or a portion of (e.g., a cytoplasmic-domain truncated portion) of an MHC class I alpha chain protein and  $\beta_2$  microglobulin protein or an MHC class II alpha chain protein and an MHC class II beta chain protein to thereby express MHC class I or MHC class II proteins on the cell surface. Expression of the appropriate class I or class II MHC in conjunction with a peptide having the activity of a B lymphocyte antigen (e.g., B7-1, B7-2, B7-3) induces a T cell mediated immune response against the transfected tumor cell. Optionally, a gene encoding an antisense construct which blocks expression of an MHC class II associated protein, such as the invariant chain, can also be cotransfected with a DNA encoding a peptide having the activity of a B lymphocyte antigen to promote presentation of tumor associated antigens and induce tumor specific immunity. Thus, the induction of a T cell mediated immune response in a human subject may be sufficient to overcome tumor-specific tolerance in the subject.

The activity of a protein of the invention may, among other means, be measured by the following methods:

Suitable assays for thymocyte or splenocyte cytotoxicity include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Herrmann et al., Proc. Natl. Acad. Sci. USA 78:2488-2492, 1981; Herrmann et al., J. Immunol. 128:1968-1974, 1982; Handa et al., J. Immunol. 135:1564-1572, 1985; Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J.

Immunol. 140:508-512, 1988; Bowman et al., J. Virology 61:1992-1998; Bertagnolli et al., Cellular Immunology 133:327-341, 1991; Brown et al., J. Immunol. 153:3079-3092, 1994.

Assays for T-cell-dependent immunoglobulin responses and isotype switching (which will identify, among others, proteins that modulate T-cell dependent antibody responses and that affect Th1/Th2 profiles) include, without limitation, those described in: Maliszewski, J. Immunol. 144:3028-3033, 1990; and Assays for B cell function: In vitro antibody production, Mond, J. J. and Brunswick, M. In Current Protocols in Immunology. J. E. e.a. Coligan eds. Vol 1 pp. 3.8.1-3.8.16, John Wiley and Sons, Toronto. 1994.

Mixed lymphocyte reaction (MLR) assays (which will identify, among others, proteins that generate predominantly Th1 and CTL responses) include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 3, In Vitro assays for Mouse Lymphocyte Function 3.1-3.19; Chapter 7, Immunologic studies in Humans); Takai et al., J. Immunol. 137:3494-3500, 1986; Takai et al., J. Immunol. 140:508-512, 1988; Bertagnolli et al., J. Immunol. 149:3778-3783, 1992.

Dendritic cell-dependent assays (which will identify, among others, proteins expressed by dendritic cells that activate naive T-cells) include, without limitation, those described in: Guery et al., J. Immunol. 134:536-544, 1995; Inaba et al., Journal of Experimental Medicine 173:549-559, 1991; Macatonia et al., Journal of Immunology 154:5071-5079, 1995; Porgador et al., Journal of Experimental Medicine 182:255-260, 1995; Nair et al., Journal of Virology 67:4062-4069, 1993; Huang et al., Science 264:961-965, 1994; Macatonia et al., Journal of Experimental Medicine 169:1255-1264, 1989; Bhardwaj et al., Journal of Clinical Investigation 94:797-807, 1994; and Inaba et al., Journal of Experimental Medicine 172:631-640, 1990.

Assays for lymphocyte survival/apoptosis (which will identify, among others, proteins that prevent apoptosis after superantigen induction and proteins that regulate lymphocyte homeostasis) include, without limitation, those described in: Darzynkiewicz et al., Cytometry 13:795-808, 1992; Gorczyca et al., Leukemia 7:659-670, 1993; Gorczyca et al., Cancer Research 53:1945-1951, 1993; Itoh et al., Cell 66:233-243, 1991; Zacharchuk, Journal of Immunology 145:4037-4045, 1990; Zamai et al., Cytometry 14:891-897, 1993; Gorczyca et al., International Journal of Oncology 1:639-648, 1992.

Assays for proteins that influence early steps of T-cell commitment and development include, without limitation, those described in: Antica et al., Blood 84:111-117, 1994; Fine et

al., Cellular Immunology 155:111-122, 1994; Galy et al., Blood 85:2770-2778, 1995; Toki et al., Proc. Nat. Acad. Sci. USA 88:7548-7551, 1991.

#### 5.10.8 ACTIVIN/INHIBIN ACTIVITY

5 A polypeptide of the present invention may also exhibit activin- or inhibin-related activities. A polynucleotide of the invention may encode a polypeptide exhibiting such characteristics. Inhibins are characterized by their ability to inhibit the release of follicle stimulating hormone (FSH), while activins are characterized by their ability to stimulate the release of follicle stimulating hormone (FSH). Thus, a polypeptide of the present  
10 invention, alone or in heterodimers with a member of the inhibin family, may be useful as a contraceptive based on the ability of inhibins to decrease fertility in female mammals and decrease spermatogenesis in male mammals. Administration of sufficient amounts of other inhibins can induce infertility in these mammals. Alternatively, the polypeptide of the invention, as a homodimer or as a heterodimer with other protein subunits of the inhibin  
15 group, may be useful as a fertility inducing therapeutic, based upon the ability of activin molecules in stimulating FSH release from cells of the anterior pituitary. See, for example, U.S. Pat. No. 4,798,885. A polypeptide of the invention may also be useful for advancement of the onset of fertility in sexually immature mammals, so as to increase the lifetime reproductive performance of domestic animals such as, but not limited to, cows, sheep and  
20 pigs.

The activity of a polypeptide of the invention may, among other means, be measured by the following methods.

Assays for activin/inhibin activity include, without limitation, those described in: Vale et al., Endocrinology 91:562-572, 1972; Ling et al., Nature 321:779-782, 1986; Vale et al.,  
25 Nature 321:776-779, 1986; Mason et al., Nature 318:659-663, 1985; Forage et al., Proc. Natl. Acad. Sci. USA 83:3091-3095, 1986.

#### 5.10.9 CHEMOTACTIC/CHEMOKINETIC ACTIVITY

A polypeptide of the present invention may be involved in chemotactic or chemokinetic  
30 activity for mammalian cells, including, for example, monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or endothelial cells. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Chemotactic and chemokinetic receptor activation can be used to mobilize or attract a desired cell population to a desired site of action. Chemotactic or chemokinetic compositions (e.g. proteins, antibodies, binding



partners, or modulators of the invention) provide particular advantages in treatment of wounds and other trauma to tissues, as well as in treatment of localized infections. For example, attraction of lymphocytes, monocytes or neutrophils to tumors or sites of infection may result in improved immune responses against the tumor or infecting agent.

5           A protein or peptide has chemotactic activity for a particular cell population if it can stimulate, directly or indirectly, the directed orientation or movement of such cell population. Preferably, the protein or peptide has the ability to directly stimulate directed movement of cells. Whether a particular protein has chemotactic activity for a population of cells can be readily determined by employing such protein or peptide in any known assay for cell  
10 chemotaxis.

Therapeutic compositions of the invention can be used in the following:

Assays for chemotactic activity (which will identify proteins that induce or prevent chemotaxis) consist of assays that measure the ability of a protein to induce the migration of cells across a membrane as well as the ability of a protein to induce the adhesion of one cell  
15 population to another cell population. Suitable assays for movement and adhesion include, without limitation, those described in: Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Marguiles, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley-Interscience (Chapter 6.12, Measurement of alpha and beta Chemokines 6.12.1-6.12.28; Taub et al. J. Clin. Invest. 95:1370-1376, 1995; Lind et al. APMIS 103:140-  
20 146, 1995; Muller et al Eur. J. Immunol. 25:1744-1748; Gruber et al. J. of Immunol. 152:5860-5867, 1994; Johnston et al. J. of Immunol. 153:1762-1768, 1994.

#### 5.10.10           HEMOSTATIC AND THROMBOLYTIC ACTIVITY

A polypeptide of the invention may also be involved in hemostasis or thrombolysis or  
25 thrombosis. A polynucleotide of the invention can encode a polypeptide exhibiting such attributes. Compositions may be useful in treatment of various coagulation disorders (including hereditary disorders, such as hemophilias) or to enhance coagulation and other hemostatic events in treating wounds resulting from trauma, surgery or other causes. A composition of the invention may also be useful for dissolving or inhibiting formation of  
30 thromboses and for treatment and prevention of conditions resulting therefrom (such as, for example, infarction of cardiac and central nervous system vessels (e.g., stroke).

Therapeutic compositions of the invention can be used in the following:

Assay for hemostatic and thrombolytic activity include, without limitation, those described in: Linet et al., J. Clin. Pharmacol. 26:131-140, 1986; Burdick et al., Thrombosis

Rés. 45:413-419, 1987; Humphrey et al., Fibrinolysis 5:71-79 (1991); Schaub, Prostaglandins 35:467-474, 1988.

#### 5.10.11 CANCER DIAGNOSIS AND THERAPY

5 Polypeptides of the invention may be involved in cancer cell generation, proliferation or metastasis. Detection of the presence or amount of polynucleotides or polypeptides of the invention may be useful for the diagnosis and/or prognosis of one or more types of cancer. For example, the presence or increased expression of a polynucleotide/polypeptide of the invention may indicate a hereditary risk of cancer, a precancerous condition, or an ongoing  
10 malignancy. Conversely, a defect in the gene or absence of the polypeptide may be associated with a cancer condition. Identification of single nucleotide polymorphisms associated with cancer or a predisposition to cancer may also be useful for diagnosis or prognosis.

Cancer treatments promote tumor regression by inhibiting tumor cell proliferation, inhibiting angiogenesis (growth of new blood vessels that is necessary to support tumor  
15 growth) and/or prohibiting metastasis by reducing tumor cell motility or invasiveness. Therapeutic compositions of the invention may be effective in adult and pediatric oncology including in solid phase tumors/malignancies, locally advanced tumors, human soft tissue sarcomas, metastatic cancer, including lymphatic metastases, blood cell malignancies including multiple myeloma, acute and chronic leukemias, and lymphomas, head and neck cancers  
20 including mouth cancer, larynx cancer and thyroid cancer, lung cancers including small cell carcinoma and non-small cell cancers, breast cancers including small cell carcinoma and ductal carcinoma, gastrointestinal cancers including esophageal cancer, stomach cancer, colon cancer, colorectal cancer and polyps associated with colorectal neoplasia, pancreatic cancers, liver cancer, urologic cancers including bladder cancer and prostate cancer, malignancies of the  
25 female genital tract including ovarian carcinoma, uterine (including endometrial) cancers, and solid tumor in the ovarian follicle, kidney cancers including renal cell carcinoma, brain cancers including intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, metastatic tumor cell invasion in the central nervous system, bone cancers including osteomas, skin cancers including malignant melanoma, tumor progression of human skin keratinocytes,  
30 squamous cell carcinoma, basal cell carcinoma, hemangiopericytoma and Kaposi's sarcoma.

Polypeptides, polynucleotides, or modulators of polypeptides of the invention (including inhibitors and stimulators of the biological activity of the polypeptide of the invention) may be administered to treat cancer. Therapeutic compositions can be administered in therapeutically effective dosages alone or in combination with adjuvant cancer therapy such as surgery,

chemotherapy, radiotherapy, thermotherapy, and laser therapy, and may provide a beneficial effect, e.g. reducing tumor size, slowing rate of tumor growth, inhibiting metastasis, or otherwise improving overall clinical condition, without necessarily eradicating the cancer.

The composition can also be administered in therapeutically effective amounts as a  
5 portion of an anti-cancer cocktail. An anti-cancer cocktail is a mixture of the polypeptide or modulator of the invention with one or more anti-cancer drugs in addition to a pharmaceutically acceptable carrier for delivery. The use of anti-cancer cocktails as a cancer treatment is routine. Anti-cancer drugs that are well known in the art and can be used as a treatment in combination with the polypeptide or modulator of the invention include:  
10 Actinomycin D, Aminoglutethimide, Asparaginase, Bleomycin, Busulfan, Carboplatin, Carmustine, Chlorambucil, Cisplatin (cis-DDP), Cyclophosphamide, Cytarabine HCl (Cytosine arabinoside), Dacarbazine, Dactinomycin, Daunorubicin HCl, Doxorubicin HCl, Estramustine phosphate sodium, Etoposide (V16-213), Floxuridine, 5-Fluorouracil (5-Fu), Flutamide, Hydroxyurea (hydroxycarbamide), Ifosfamide, Interferon Alpha-2a, Interferon  
15 Alpha-2b, Leuprolide acetate (LHRH-releasing factor analog), Lomustine, Mechlorethamine HCl (nitrogen mustard), Melphalan, Mercaptopurine, Mesna, Methotrexate (MTX), Mitomycin, Mitoxantrone HCl, Octreotide, Plicamycin, Procarbazine HCl, Streptozocin, Tamoxifen citrate, Thioguanine, Thiotepa, Vinblastine sulfate, Vincristine sulfate, Amsacrine, Azacitidine, Hexamethylmelamine, Interleukin-2, Mitoguazone, Pentostatin, Semustine,  
20 Teniposide, and Vindesine sulfate.

In addition, therapeutic compositions of the invention may be used for prophylactic treatment of cancer. There are hereditary conditions and/or environmental situations (e.g. exposure to carcinogens) known in the art that predispose an individual to developing cancers. Under these circumstances, it may be beneficial to treat these individuals with therapeutically  
25 effective doses of the polypeptide of the invention to reduce the risk of developing cancers.

*In vitro* models can be used to determine the effective doses of the polypeptide of the invention as a potential cancer treatment. These *in vitro* models include proliferation assays of cultured tumor cells, growth of cultured tumor cells in soft agar (see Freshney, (1987) Culture of Animal Cells: A Manual of Basic Technique, Wiley-Liss, New York, NY Ch 18 and Ch 21),  
30 tumor systems in nude mice as described in Giovanella et al., J. Natl. Can. Inst., 52: 921-30 (1974), mobility and invasive potential of tumor cells in Boyden Chamber assays as described in Pilkington et al., Anticancer Res., 17: 4107-9 (1997), and angiogenesis assays such as induction of vascularization of the chick chorioallantoic membrane or induction of vascular endothelial cell migration as described in Ribatta et al., Intl. J. Dev. Biol., 40: 1189-97 (1999)

and Li et al., Clin. Exp. Metastasis, 17:423-9 (1999), respectively. Suitable tumor cells lines are available, e.g. from American Type Tissue Culture Collection catalogs.

#### 5.10.12 RECEPTOR/LIGAND ACTIVITY

5 A polypeptide of the present invention may also demonstrate activity as receptor, receptor ligand or inhibitor or agonist of receptor/ligand interactions. A polynucleotide of the invention can encode a polypeptide exhibiting such characteristics. Examples of such receptors and ligands include, without limitation, cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell  
10 interactions and their ligands (including without limitation, cellular adhesion molecules (such as selectins, integrins and their ligands) and receptor/ligand pairs involved in antigen presentation, antigen recognition and development of cellular and humoral immune responses. Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the present invention  
15 (including, without limitation, fragments of receptors and ligands) may themselves be useful as inhibitors of receptor/ligand interactions.

The activity of a polypeptide of the invention may, among other means, be measured by the following methods:

Suitable assays for receptor-ligand activity include without limitation those described in:  
20 Current Protocols in Immunology, Ed by J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, W. Strober, Pub. Greene Publishing Associates and Wiley- Interscience (Chapter 7.28, Measurement of Cellular Adhesion under static conditions 7.28.1- 7.28.22), Takai et al., Proc. Natl. Acad. Sci. USA 84:6864-6868, 1987; Bierer et al., J. Exp. Med. 168:1145-1156, 1988; Rosenstein et al., J. Exp. Med. 169:149-160 1989; Stoltenborg et al., J.  
25 Immunol. Methods 175:59-68, 1994; Stitt et al., Cell 80:661-670, 1995.

By way of example, the polypeptides of the invention may be used as a receptor for a ligand(s) thereby transmitting the biological activity of that ligand(s). Ligands may be identified through binding assays, affinity chromatography, dihybrid screening assays, BIAcore assays, gel overlay assays, or other methods known in the art.

30 Studies characterizing drugs or proteins as agonist or antagonist or partial agonists or a partial antagonist require the use of other proteins as competing ligands. The polypeptides of the present invention or ligand(s) thereof may be labeled by being coupled to radioisotopes, colorimetric molecules or a toxin molecules by conventional methods. ("Guide to Protein Purification" Murray P. Deutscher (ed) Methods in Enzymology Vol. 182 (1990) Academic

Press, Inc. San Diego). Examples of radioisotopes include, but are not limited to, tritium and carbon-14. Examples of colorimetric molecules include, but are not limited to, fluorescent molecules such as fluorescamine, or rhodamine or other colorimetric molecules. Examples of toxins include, but are not limited, to ricin.

5

### 5.10.13 DRUG SCREENING

This invention is particularly useful for screening chemical compounds by using the novel polypeptides or binding fragments thereof in any of a variety of drug screening techniques. The polypeptides or fragments employed in such a test may either be free in  
10 solution, affixed to a solid support, borne on a cell surface or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or a fragment thereof. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for  
15 example, the formation of complexes between polypeptides of the invention or fragments and the agent being tested or examine the diminution in complex formation between the novel polypeptides and an appropriate cell line, which are well known in the art.

Sources for test compounds that may be screened for ability to bind to or modulate (i.e., increase or decrease) the activity of polypeptides of the invention include (1) inorganic  
20 and organic chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of either random or mimetic peptides, oligonucleotides or organic molecules.

Chemical libraries may be readily synthesized or purchased from a number of commercial sources, and may include structural analogs of known compounds or compounds that are identified as "hits" or "leads" via natural product screening.

25 The sources of natural product libraries are microorganisms (including bacteria and fungi), animals, plants or other vegetation, or marine organisms, and libraries of mixtures for screening may be created by: (1) fermentation and extraction of broths from soil, plant or marine microorganisms or (2) extraction of the organisms themselves. Natural product libraries include polyketides, non-ribosomal peptides, and (non-naturally occurring) variants  
30 thereof. For a review, see *Science* 282:63-68 (1998).

Combinatorial libraries are composed of large numbers of peptides, oligonucleotides or organic compounds and can be readily prepared by traditional automated synthesis methods, PCR, cloning or proprietary synthetic methods. Of particular interest are peptide and oligonucleotide combinatorial libraries. Still other libraries of interest include peptide, protein,

peptidomimetic, multiparallel synthetic collection, recombinatorial, and polypeptide libraries. For a review of combinatorial chemistry and libraries created therefrom, see Myers, *Curr. Opin. Biotechnol.* 8:701-707 (1997). For reviews and examples of peptidomimetic libraries, see Al-Obeidi et al., *Mol. Biotechnol.* 9(3):205-23 (1998); Hruby et al., *Curr Opin Chem Biol*, 1(1):114-19 (1997); Dorner et al., *Bioorg Med Chem*, 4(5):709-15 (1996) (alkylated dipeptides).

Identification of modulators through use of the various libraries described herein permits modification of the candidate "hit" (or "lead") to optimize the capacity of the "hit" to bind a polypeptide of the invention. The molecules identified in the binding assay are then  
10 tested for antagonist or agonist activity in *in vivo* tissue culture or animal models that are well known in the art. In brief, the molecules are titrated into a plurality of cell cultures or animals and then tested for either cell/animal death or prolonged survival of the animal/cells.

The binding molecules thus identified may be complexed with toxins, e.g., ricin or cholera, or with other compounds that are toxic to cells such as radioisotopes. The toxin-  
15 binding molecule complex is then targeted to a tumor or other cell by the specificity of the binding molecule for a polypeptide of the invention. Alternatively, the binding molecules may be complexed with imaging agents for targeting and imaging purposes.

#### 5.10.14 ASSAY FOR RECEPTOR ACTIVITY

20 The invention also provides methods to detect specific binding of a polypeptide e.g. a ligand or a receptor. The art provides numerous assays particularly useful for identifying previously unknown binding partners for receptor polypeptides of the invention. For example, expression cloning using mammalian or bacterial cells, or dihybrid screening assays can be used to identify polynucleotides encoding binding partners. As another example, affinity  
25 chromatography with the appropriate immobilized polypeptide of the invention can be used to isolate polypeptides that recognize and bind polypeptides of the invention. There are a number of different libraries used for the identification of compounds, and in particular small molecule, that modulate (*i.e.*, increase or decrease) biological activity of a polypeptide of the invention. Ligands for receptor polypeptides of the invention can also be identified by adding  
30 exogenous ligands, or cocktails of ligands to two cells populations that are genetically identical except for the expression of the receptor of the invention: one cell population expresses the receptor of the invention whereas the other does not. The response of the two cell populations to the addition of ligands(s) are then compared. Alternatively, an expression library can be co-expressed with the polypeptide of the invention in cells and assayed for an autocrine response

to identify potential ligand(s). As still another example, BIAcore assays, gel overlay assays, or other methods known in the art can be used to identify binding partner polypeptides, including, (1) organic and inorganic chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of random peptides, oligonucleotides or organic molecules.

5           The role of downstream intracellular signaling molecules in the signaling cascade of the polypeptide of the invention can be determined. For example, a chimeric protein in which the cytoplasmic domain of the polypeptide of the invention is fused to the extracellular portion of a protein, whose ligand has been identified, is produced in a host cell. The cell is then incubated with the ligand specific for the extracellular portion of the chimeric protein, thereby activating  
10       the chimeric receptor. Known downstream proteins involved in intracellular signaling can then be assayed for expected modifications i.e. phosphorylation. Other methods known to those in the art can also be used to identify signaling molecules involved in receptor activity.

#### 5.10.15           ANTI-INFLAMMATORY ACTIVITY

15           Compositions of the present invention may also exhibit anti-inflammatory activity. The anti-inflammatory activity may be achieved by providing a stimulus to cells involved in the inflammatory response, by inhibiting or promoting cell-cell interactions (such as, for example, cell adhesion), by inhibiting or promoting chemotaxis of cells involved in the inflammatory process, inhibiting or promoting cell extravasation, or by stimulating or suppressing production  
20       of other factors which more directly inhibit or promote an inflammatory response.

Compositions with such activities can be used to treat inflammatory conditions including chronic or acute conditions, including without limitation intimation associated with infection (such as septic shock, sepsis or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection,  
25       nephritis, cytokine or chemokine-induced lung injury, inflammatory bowel disease, Crohn's disease or resulting from over production of cytokines such as TNF or IL-1. Compositions of the invention may also be useful to treat anaphylaxis and hypersensitivity to an antigenic substance or material. Compositions of this invention may be utilized to prevent or treat conditions such as, but not limited to, sepsis, acute pancreatitis, endotoxin shock, cytokine  
30       induced shock, rheumatoid arthritis, chronic inflammatory arthritis, pancreatic cell damage from diabetes mellitus type 1, graft versus host disease, inflammatory bowel disease, inflammation associated with pulmonary disease, other autoimmune disease or inflammatory disease, an antiproliferative agent such as for acute or chronic myelogenous leukemia or in the prevention of premature labor secondary to intrauterine infections.

#### 5.10.16 LEUKEMIAS

Leukemias and related disorders may be treated or prevented by administration of a therapeutic that promotes or inhibits function of the polynucleotides and/or polypeptides of the invention. Such leukemias and related disorders include but are not limited to acute leukemia, acute lymphocytic leukemia, acute myelocytic leukemia, myeloblastic, promyelocytic, myelomonocytic, monocytic, erythroleukemia, chronic leukemia, chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia (for a review of such disorders, see Fishman et al., 1985, Medicine, 2d Ed., J.B. Lippincott Co., Philadelphia).

#### 5.10.17 NERVOUS SYSTEM DISORDERS

Nervous system disorders, involving cell types which can be tested for efficacy of intervention with compounds that modulate the activity of the polynucleotides and/or polypeptides of the invention, and which can be treated upon thus observing an indication of therapeutic utility, include but are not limited to nervous system injuries, and diseases or disorders which result in either a disconnection of axons, a diminution or degeneration of neurons, or demyelination. Nervous system lesions which may be treated in a patient (including human and non-human mammalian patients) according to the invention include but are not limited to the following lesions of either the central (including spinal cord, brain) or peripheral nervous systems:

(i) traumatic lesions, including lesions caused by physical injury or associated with surgery, for example, lesions which sever a portion of the nervous system, or compression injuries;

(ii) ischemic lesions, in which a lack of oxygen in a portion of the nervous system results in neuronal injury or death, including cerebral infarction or ischemia, or spinal cord infarction or ischemia;

(iii) infectious lesions, in which a portion of the nervous system is destroyed or injured as a result of infection, for example, by an abscess or associated with infection by human immunodeficiency virus, herpes zoster, or herpes simplex virus or with Lyme disease, tuberculosis, syphilis;

(iv) degenerative lesions, in which a portion of the nervous system is destroyed or injured as a result of a degenerative process including but not limited to degeneration associated with Parkinson's disease, Alzheimer's disease, Huntington's chorea, or amyotrophic lateral sclerosis;



(v) lesions associated with nutritional diseases or disorders, in which a portion of the nervous system is destroyed or injured by a nutritional disorder or disorder of metabolism including but not limited to, vitamin B12 deficiency, folic acid deficiency, Wernicke disease, tobacco-alcohol amblyopia, Marchiafava-Bignami disease (primary degeneration of the corpus callosum), and alcoholic cerebellar degeneration;

(vi) neurological lesions associated with systemic diseases including but not limited to diabetes (diabetic neuropathy, Bell's palsy), systemic lupus erythematosus, carcinoma, or sarcoidosis;

(vii) lesions caused by toxic substances including alcohol, lead, or particular neurotoxins; and

(viii) demyelinated lesions in which a portion of the nervous system is destroyed or injured by a demyelinating disease including but not limited to multiple sclerosis, human immunodeficiency virus-associated myelopathy, transverse myelopathy or various etiologies, progressive multifocal leukoencephalopathy, and central pontine myelinolysis.

Therapeutics which are useful according to the invention for treatment of a nervous system disorder may be selected by testing for biological activity in promoting the survival or differentiation of neurons. For example, and not by way of limitation, therapeutics which elicit any of the following effects may be useful according to the invention:

- (i) increased survival time of neurons in culture;
- (ii) increased sprouting of neurons in culture or *in vivo*;
- (iii) increased production of a neuron-associated molecule in culture or *in vivo*, *e.g.*, choline acetyltransferase or acetylcholinesterase with respect to motor neurons; or
- (iv) decreased symptoms of neuron dysfunction *in vivo*.

Such effects may be measured by any method known in the art. In preferred, non-limiting embodiments, increased survival of neurons may be measured by the method set forth in Arakawa et al. (1990, J. Neurosci. 10:3507-3515); increased sprouting of neurons may be detected by methods set forth in Pestronk et al. (1980, Exp. Neurol. 70:65-82) or Brown et al. (1981, Ann. Rev. Neurosci. 4:17-42); increased production of neuron-associated molecules may be measured by bioassay, enzymatic assay, antibody binding, Northern blot assay, *etc.*, depending on the molecule to be measured; and motor neuron dysfunction may be measured by assessing the physical manifestation of motor neuron disorder, *e.g.*, weakness, motor neuron conduction velocity, or functional disability.

In specific embodiments, motor neuron disorders that may be treated according to the invention include but are not limited to disorders such as infarction, infection, exposure to

toxin, trauma, surgical damage, degenerative disease or malignancy that may affect motor neurons as well as other components of the nervous system, as well as disorders that selectively affect neurons such as amyotrophic lateral sclerosis, and including but not limited to progressive spinal muscular atrophy, progressive bulbar palsy, primary lateral sclerosis, infantile and juvenile muscular atrophy, progressive bulbar paralysis of childhood (Fazio-Londe syndrome), poliomyelitis and the post polio syndrome, and Hereditary Motor-sensory Neuropathy (Charcot-Marie-Tooth Disease).

#### 5.10.18 OTHER ACTIVITIES

A polypeptide of the invention may also exhibit one or more of the following additional activities or effects: inhibiting the growth, infection or function of, or killing, infectious agents, including, without limitation, bacteria, viruses, fungi and other parasites; effecting (suppressing or enhancing) bodily characteristics, including, without limitation, height, weight, hair color, eye color, skin, fat to lean ratio or other tissue pigmentation, or organ or body part size or shape (such as, for example, breast augmentation or diminution, change in bone form or shape); effecting biorhythms or circadian cycles or rhythms; effecting the fertility of male or female subjects; effecting the metabolism, catabolism, anabolism, processing, utilization, storage or elimination of dietary fat, lipid, protein, carbohydrate, vitamins, minerals, co-factors or other nutritional factors or component(s); effecting behavioral characteristics, including, without limitation, appetite, libido, stress, cognition (including cognitive disorders), depression (including depressive disorders) and violent behaviors; providing analgesic effects or other pain reducing effects; promoting differentiation and growth of embryonic stem cells in lineages other than hematopoietic lineages; hormonal or endocrine activity; in the case of enzymes, correcting deficiencies of the enzyme and treating deficiency-related diseases; treatment of hyperproliferative disorders (such as, for example, psoriasis); immunoglobulin-like activity (such as, for example, the ability to bind antigens or complement); and the ability to act as an antigen in a vaccine composition to raise an immune response against such protein or another material or entity which is cross-reactive with such protein.

#### 5.10.19 IDENTIFICATION OF POLYMORPHISMS

The demonstration of polymorphisms makes possible the identification of such polymorphisms in human subjects and the pharmacogenetic use of this information for diagnosis and treatment. Such polymorphisms may be associated with, e.g., differential predisposition or susceptibility to various disease states (such as disorders involving

inflammation or immune response) or a differential response to drug administration, and this genetic information can be used to tailor preventive or therapeutic treatment appropriately. For example, the existence of a polymorphism associated with a predisposition to inflammation or autoimmune disease makes possible the diagnosis of this condition in humans by identifying the presence of the polymorphism.

Polymorphisms can be identified in a variety of ways known in the art which all generally involve obtaining a sample from a patient, analyzing DNA from the sample, optionally involving isolation or amplification of the DNA, and identifying the presence of the polymorphism in the DNA. For example, PCR may be used to amplify an appropriate fragment of genomic DNA which may then be sequenced. Alternatively, the DNA may be subjected to allele-specific oligonucleotide hybridization (in which appropriate oligonucleotides are hybridized to the DNA under conditions permitting detection of a single base mismatch) or to a single nucleotide extension assay (in which an oligonucleotide that hybridizes immediately adjacent to the position of the polymorphism is extended with one or more labeled nucleotides). In addition, traditional restriction fragment length polymorphism analysis (using restriction enzymes that provide differential digestion of the genomic DNA depending on the presence or absence of the polymorphism) may be performed. Arrays with nucleotide sequences of the present invention can be used to detect polymorphisms. The array can comprise modified nucleotide sequences of the present invention in order to detect the nucleotide sequences of the present invention. In the alternative, any one of the nucleotide sequences of the present invention can be placed on the array to detect changes from those sequences.

Alternatively a polymorphism resulting in a change in the amino acid sequence could also be detected by detecting a corresponding change in amino acid sequence of the protein, e.g., by an antibody specific to the variant sequence.

#### **5.10.20 ARTHRITIS AND INFLAMMATION**

The immunosuppressive effects of the compositions of the invention against rheumatoid arthritis is determined in an experimental animal model system. The experimental model system is adjuvant induced arthritis in rats, and the protocol is described by J. Holoshitz, et al., 1983, Science, 219:56, or by B. Waksman et al., 1963, Int. Arch. Allergy Appl. Immunol., 23:129. Induction of the disease can be caused by a single injection, generally intradermally, of a suspension of killed Mycobacterium tuberculosis in complete Freund's adjuvant (CFA). The route of injection can vary, but rats may be injected at the base of the tail

with an adjuvant mixture. The polypeptide is administered in phosphate buffered solution (PBS) at a dose of about 1-5 mg/kg. The control consists of administering PBS only.

The procedure for testing the effects of the test compound would consist of intradermally injecting killed *Mycobacterium tuberculosis* in CFA followed by immediately  
5 administering the test compound and subsequent treatment every other day until day 24. At 14, 15, 18, 20, 22, and 24 days after injection of *Mycobacterium* CFA, an overall arthritis score may be obtained as described by J. Holoskitz above. An analysis of the data would reveal that the test compound would have a dramatic affect on the swelling of the joints as measured by a decrease of the arthritis score.

10

## 5.11 THERAPEUTIC METHODS

The compositions (including polypeptide fragments, analogs, variants and antibodies or other binding partners or modulators including antisense polynucleotides) of the invention have  
15 numerous applications in a variety of therapeutic methods. Examples of therapeutic applications include, but are not limited to, those exemplified herein.

### 5.11.1 EXAMPLE

One embodiment of the invention is the administration of an effective amount of the  
20 prothrombinase-like polypeptides or other composition of the invention to individuals affected by a disease or disorder that can be modulated by regulating the soluble prothrombinase-like receptor of the invention. While the mode of administration is not particularly important, parenteral administration is preferred. An exemplary mode of administration is to deliver an intravenous bolus. The dosage of prothrombinase-like polypeptides or other composition of  
25 the invention will normally be determined by the prescribing physician. It is to be expected that the dosage will vary according to the age, weight, condition and response of the individual patient. Typically, the amount of polypeptide administered per dose will be in the range of about 0.01µg/kg to 100 mg/kg of body weight, with the preferred dose being about 0.1µg/kg to 10 mg/kg of patient body weight. For parenteral administration, prothrombinase-like  
30 polypeptides of the invention will be formulated in an injectable form combined with a pharmaceutically acceptable parenteral vehicle. Such vehicles are well known in the art and examples include water, saline, Ringer's solution, dextrose solution, and solutions consisting of small amounts of the human serum albumin. The vehicle may contain minor amounts of

additives that maintain the isotonicity and stability of the polypeptide or other active ingredient. The preparation of such solutions is within the skill of the art.

## 5 5.12 PHARMACEUTICAL FORMULATIONS AND ROUTES OF ADMINISTRATION

A protein or other composition of the present invention (from whatever source derived, including without limitation from recombinant and non-recombinant sources and including antibodies and other binding partners of the polypeptides of the invention) may be administered  
10 to a patient in need, by itself, or in pharmaceutical compositions where it is mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a variety of disorders. Such a composition may optionally contain (in addition to protein or other active ingredient and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term "pharmaceutically acceptable" means a non-toxic material that does not  
15 interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF,  
20 thrombopoietin, stem cell factor, and erythropoietin. In further compositions, proteins of the invention may be combined with other agents beneficial to the treatment of the disease or disorder in question. These agents include various growth factors such as epidermal growth factor (EGF), platelet-derived growth factor (PDGF), transforming growth factors (TGF- $\alpha$  and TGF- $\beta$ ), insulin-like growth factor (IGF), as well as cytokines described herein.

25 The pharmaceutical composition may further contain other agents which either enhance the activity of the protein or other active ingredient or complement its activity or use in treatment. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with protein or other active ingredient of the invention, or to minimize side effects. Conversely, protein or other active ingredient of the  
30 present invention may be included in formulations of the particular clotting factor, cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent to minimize side effects of the clotting factor, cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, or anti-inflammatory agent (such

as IL-1Ra, IL-1 Hy1, IL-1 Hy2, anti-TNF, corticosteroids, immunosuppressive agents). A protein of the present invention may be active in multimers (e.g., heterodimers or homodimers) or complexes with itself or other proteins. As a result, pharmaceutical compositions of the invention may comprise a protein of the invention in such multimeric or complexed form.

As an alternative to being included in a pharmaceutical composition of the invention including a first protein, a second protein or a therapeutic agent may be concurrently administered with the first protein (e.g., at the same time, or at differing times provided that therapeutic concentrations of the combination of agents is achieved at the treatment site).

Techniques for formulation and administration of the compounds of the instant application may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, latest edition. A therapeutically effective dose further refers to that amount of the compound sufficient to result in amelioration of symptoms, *e.g.*, treatment, healing, prevention or amelioration of the relevant medical condition, or an increase in rate of treatment, healing, prevention or amelioration of such conditions. When applied to an individual active ingredient, administered alone, a therapeutically effective dose refers to that ingredient alone. When applied to a combination, a therapeutically effective dose refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of protein or other active ingredient of the present invention is administered to a mammal having a condition to be treated. Protein or other active ingredient of the present invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines or other hematopoietic factors. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, protein or other active ingredient of the present invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering protein or other active ingredient of the present invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), thrombolytic or anti-thrombotic factors.

### 5.12.1 ROUTES OF ADMINISTRATION

Suitable routes of administration may, for example, include oral, rectal, transmucosal, or intestinal administration; parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections. Administration of protein or other active ingredient of the present invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, topical application or cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. Intravenous administration to the patient is preferred.

Alternately, one may administer the compound in a local rather than systemic manner, for example, via injection of the compound directly into a arthritic joints or in fibrotic tissue, often in a depot or sustained release formulation. In order to prevent the scarring process frequently occurring as complication of glaucoma surgery, the compounds may be administered topically, for example, as eye drops. Furthermore, one may administer the drug in a targeted drug delivery system, for example, in a liposome coated with a specific antibody, targeting, for example, arthritic or fibrotic tissue. The liposomes will be targeted to and taken up selectively by the afflicted tissue.

The polypeptides of the invention are administered by any route that delivers an effective dosage to the desired site of action. The determination of a suitable route of administration and an effective dosage for a particular indication is within the level of skill in the art. Preferably for wound treatment, one administers the therapeutic compound directly to the site. Suitable dosage ranges for the polypeptides of the invention can be extrapolated from these dosages or from similar studies in appropriate animal models. Dosages can then be adjusted as necessary by the clinician to provide maximal therapeutic benefit.

## 5.12.2 COMPOSITIONS/FORMULATIONS

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. These pharmaceutical compositions may be manufactured in a manner that is itself known, *e.g.*, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes. Proper formulation is dependent upon the route of administration chosen. When a therapeutically effective amount of protein or other active ingredient of the present invention is administered orally, protein or other active ingredient of the present

invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% protein or other active ingredient of the present invention, and preferably from about 25 to 90% protein or other active ingredient of the present invention. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5 to 90% by weight of protein or other active ingredient of the present invention, and preferably from about 1 to 50% protein or other active ingredient of the present invention.

When a therapeutically effective amount of protein or other active ingredient of the present invention is administered by intravenous, cutaneous or subcutaneous injection, protein or other active ingredient of the present invention will be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such parenterally acceptable protein or other active ingredient solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or subcutaneous injection should contain, in addition to protein or other active ingredient of the present invention, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art. For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

For oral administration, the compounds can be formulated readily by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained solid excipient,



optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, 5 gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, 10 talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active compound doses.

Pharmaceutical preparations which can be used orally include push-fit capsules made of 15 gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In 20 addition, stabilizers may be added. All formulations for oral administration should be in dosages suitable for such administration. For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from 25 pressurized packs or a nebuliser, with the use of a suitable propellant, *e.g.*, dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, *e.g.*, gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound 30 and a suitable powder base such as lactose or starch. The compounds may be formulated for parenteral administration by injection, *e.g.*, by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, *e.g.*, in ampules or in multi-dose containers, with an added preservative. The compositions may take such forms as

suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active  
5 compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers  
10 or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, *e.g.*, sterile pyrogen-free water, before use.

The compounds may also be formulated in rectal compositions such as suppositories or retention enemas, *e.g.*, containing conventional suppository bases such as cocoa butter or other  
15 glycerides. In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as  
20 sparingly soluble derivatives, for example, as a sparingly soluble salt.

A pharmaceutical carrier for the hydrophobic compounds of the invention is a co-solvent system comprising benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. The co-solvent system may be the VPD co-solvent system. VPD is a solution of 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant polysorbate  
25 80, and 65% w/v polyethylene glycol 300, made up to volume in absolute ethanol. The VPD co-solvent system (VPD:5W) consists of VPD diluted 1:1 with a 5% dextrose in water solution. This co-solvent system dissolves hydrophobic compounds well, and itself produces low toxicity upon systemic administration. Naturally, the proportions of a co-solvent system may be varied considerably without destroying its solubility and toxicity characteristics.  
30 Furthermore, the identity of the co-solvent components may be varied: for example, other low-toxicity nonpolar surfactants may be used instead of polysorbate 80; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, *e.g.* polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose. Alternatively, other delivery systems for hydrophobic pharmaceutical compounds

may be employed. Liposomes and emulsions are well known examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed, although usually at the cost of greater toxicity. Additionally, the compounds may be delivered using a sustained-release system, such as semipermeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various types of sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein or other active ingredient stabilization may be employed.

The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols. Many of the active ingredients of the invention may be provided as salts with pharmaceutically compatible counter ions. Such pharmaceutically acceptable base addition salts are those salts which retain the biological effectiveness and properties of the free acids and which are obtained by reaction with inorganic or organic bases such as sodium hydroxide, magnesium hydroxide, ammonia, trialkylamine, dialkylamine, monoalkylamine, dibasic amino acids, sodium acetate, potassium benzoate, triethanol amine and the like.

The pharmaceutical composition of the invention may be in the form of a complex of the protein(s) or other active ingredient of present invention along with protein or peptide antigens. The protein and/or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC and structurally related proteins including those encoded by class I and class II MHC genes on host cells will serve to present the peptide antigen(s) to T lymphocytes. The antigen components could also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules that can directly signal T cells. Alternatively antibodies able to bind surface immunoglobulin and other molecules on B cells as well as antibodies able to bind the TCR and other molecules on T cells can be combined with the pharmaceutical composition of the invention.

The pharmaceutical composition of the invention may be in the form of a liposome in which protein of the present invention is combined, in addition to other pharmaceutically

acceptable carriers, with amphipathic agents such as lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include, without limitation, monoglycerides, diglycerides, sulfatides, lysolecithins, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. Patent Nos. 4,235,871; 4,501,728; 4,837,028; and 4,737,323, all of which are incorporated herein by reference.

The amount of protein or other active ingredient of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of protein or other active ingredient of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein or other active ingredient of the present invention and observe the patient's response. Larger doses of protein or other active ingredient of the present invention may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01  $\mu\text{g}$  to about 100 mg (preferably about 0.1  $\mu\text{g}$  to about 10 mg, more preferably about 0.1  $\mu\text{g}$  to about 1 mg) of protein or other active ingredient of the present invention per kg body weight. For compositions of the present invention which are useful for bone, cartilage, tendon or ligament regeneration, the therapeutic method includes administering the composition topically, systematically, or locally as an implant or device. When administered, the therapeutic composition for use in this invention is, of course, in a pyrogen-free, physiologically acceptable form. Further, the composition may desirably be encapsulated or injected in a viscous form for delivery to the site of bone, cartilage or tissue damage. Topical administration may be suitable for wound healing and tissue repair. Therapeutically useful agents other than a protein or other active ingredient of the invention which may also optionally be included in the composition as described above, may alternatively or additionally, be administered simultaneously or sequentially with the composition in the methods of the invention. Preferably for bone and/or cartilage formation, the composition would include a matrix capable of delivering the protein-containing or other active ingredient-containing composition to the site of bone and/or cartilage damage, providing a structure for the developing bone and cartilage and optimally capable of being resorbed into

the body. Such matrices may be formed of materials presently in use for other implanted medical applications.

The choice of matrix material is based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance and interface properties. The particular application of the compositions will define the appropriate formulation. Potential matrices for the compositions may be biodegradable and chemically defined calcium sulfate, tricalcium phosphate, hydroxyapatite, polylactic acid, polyglycolic acid and polyanhydrides. Other potential materials are biodegradable and biologically well-defined, such as bone or dermal collagen. Further matrices are comprised of pure proteins or extracellular matrix components. Other potential matrices are nonbiodegradable and chemically defined, such as sintered hydroxyapatite, bioglass, aluminates, or other ceramics. Matrices may be comprised of combinations of any of the above mentioned types of material, such as polylactic acid and hydroxyapatite or collagen and tricalcium phosphate. The bioceramics may be altered in composition, such as in calcium-aluminate-phosphate and processing to alter pore size, particle size, particle shape, and biodegradability. Presently preferred is a 50:50 (mole weight) copolymer of lactic acid and glycolic acid in the form of porous particles having diameters ranging from 150 to 800 microns. In some applications, it will be useful to utilize a sequestering agent, such as carboxymethyl cellulose or autologous blood clot, to prevent the protein compositions from disassociating from the matrix.

A preferred family of sequestering agents is cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, poly(ethylene glycol), polyoxyethylene oxide, carboxyvinyl polymer and poly(vinyl alcohol). The amount of sequestering agent useful herein is 0.5-20 wt %, preferably 1-10 wt % based on total formulation weight, which represents the amount necessary to prevent desorption of the protein from the polymer matrix and to provide appropriate handling of the composition, yet not so much that the progenitor cells are prevented from infiltrating the matrix, thereby providing the protein the opportunity to assist the osteogenic activity of the progenitor cells. In further compositions, proteins or other active ingredient of the invention may be combined with other agents beneficial to the treatment of the bone and/or cartilage defect, wound, or tissue in question. These agents include various growth factors such as epidermal growth

factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- $\alpha$  and TGF- $\beta$ ), and insulin-like growth factor (IGF).

The therapeutic compositions are also presently valuable for veterinary applications. Particularly domestic animals and thoroughbred horses, in addition to humans, are desired patients for such treatment with proteins or other active ingredient of the present invention. The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, *e.g.*, amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (*e.g.*, bone), the patient's age, sex, and diet, the severity of any infection, time of administration and other clinical factors. The dosage may vary with the type of matrix used in the reconstitution and with inclusion of other proteins in the pharmaceutical composition. For example, the addition of other known growth factors, such as IGF I (insulin like growth factor I), to the final composition, may also effect the dosage. Progress can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, X-rays, histomorphometric determinations and tetracycline labeling.

Polynucleotides of the present invention can also be used for gene therapy. Such polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a mammalian subject. Polynucleotides of the invention may also be administered by other known methods for introduction of nucleic acid into a cell or organism (including, without limitation, in the form of viral vectors or naked DNA). Cells may also be cultured *ex vivo* in the presence of proteins of the present invention in order to proliferate or to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes.

### 5.12.3 EFFECTIVE DOSAGE

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. More specifically, a therapeutically effective amount means an amount effective to prevent development of or to alleviate the existing symptoms of the subject being treated. Determination of the effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from appropriate *in vitro* assays. For example, a dose can be formulated in animal models to achieve

a circulating concentration range that can be used to more accurately determine useful doses in humans. For example, a dose can be formulated in animal models to achieve a circulating concentration range that includes the  $IC_{50}$  as determined in cell culture (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of the protein's biological activity). Such information can be used to more accurately determine useful doses in humans.

A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms or a prolongation of survival in a patient. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the  $LD_{50}$  (the dose lethal to 50% of the population) and the  $ED_{50}$  (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between  $LD_{50}$  and  $ED_{50}$ . Compounds which exhibit high therapeutic indices are preferred. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. See, *e.g.*, Fingl et al., 1975, in "The Pharmacological Basis of Therapeutics", Ch. 1 p.1. Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the desired effects, or minimal effective concentration (MEC). The MEC will vary for each compound but can be estimated from *in vitro* data. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, HPLC assays or bioassays can be used to determine plasma concentrations.

Dosage intervals can also be determined using MEC value. Compounds should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90%. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

An exemplary dosage regimen for polypeptides or other compositions of the invention will be in the range of about 0.01  $\mu\text{g/kg}$  to 100  $\text{mg/kg}$  of body weight daily, with the preferred dose being about 0.1  $\mu\text{g/kg}$  to 25  $\text{mg/kg}$  of patient body weight daily, varying in adults and

children. Dosing may be once daily, or equivalent doses may be delivered at longer or shorter intervals.

The amount of composition administered will, of course, be dependent on the subject being treated, on the subject's age and weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

#### 5.12.4 PACKAGING

The compositions may, if desired, be presented in a pack or dispenser device which may contain one or more unit dosage forms containing the active ingredient. The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising a compound of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition.

#### 5.13 ANTIBODIES

Also included in the invention are antibodies to proteins, or fragments of proteins of the invention. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, i.e., molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. Such antibodies include, but are not limited to, polyclonal, monoclonal, chimeric, single chain,  $F_{ab}$ ,  $F_{ab}'$  and  $F_{(ab)2}$  fragments, and an  $F_{ab}$  expression library. In general, an antibody molecule obtained from humans relates to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in the molecule. Certain classes have subclasses as well, such as IgG<sub>1</sub>, IgG<sub>2</sub>, and others. Furthermore, in humans, the light chain may be a kappa chain or a lambda chain. Reference herein to antibodies includes a reference to all such classes, subclasses and types of human antibody species.

An isolated related protein of the invention may be intended to serve as an antigen, or a portion or fragment thereof, and additionally can be used as an immunogen to generate antibodies that immunospecifically bind the antigen, using standard techniques for polyclonal and monoclonal antibody preparation. The full-length protein can be used or, alternatively, the invention provides antigenic peptide fragments of the antigen for use as immunogens. An antigenic peptide fragment comprises at least 6 amino acid residues of the amino acid sequence of the full length protein, such as an amino acid sequence shown in SEQ ID NO: 4, and



encompasses an epitope thereof such that an antibody raised against the peptide forms a specific immune complex with the full length protein or with any fragment that contains the epitope. Preferably, the antigenic peptide comprises at least 10 amino acid residues, or at least 15 amino acid residues, or at least 20 amino acid residues, or at least 30 amino acid residues.

- 5 Preferred epitopes encompassed by the antigenic peptide are regions of the protein that are located on its surface; commonly these are hydrophilic regions.

In certain embodiments of the invention, at least one epitope encompassed by the antigenic peptide is a region of -related protein that is located on the surface of the protein, *e.g.*, a hydrophilic region. A hydrophobicity analysis of the human related protein sequence  
10 will indicate which regions of a related protein are particularly hydrophilic and, therefore, are likely to encode surface residues useful for targeting antibody production. As a means for targeting antibody production, hydropathy plots showing regions of hydrophilicity and hydrophobicity may be generated by any method well known in the art, including, for example, the Kyte Doolittle or the Hopp Woods methods, either with or without Fourier  
15 transformation. See, *e.g.*, Hopp and Woods, 1981, *Proc. Nat. Acad. Sci. USA* 78: 3824-3828; Kyte and Doolittle 1982, *J. Mol. Biol.* 157: 105-142, each of which is incorporated herein by reference in its entirety. Antibodies that are specific for one or more domains within an antigenic protein, or derivatives, fragments, analogs or homologs thereof, are also provided herein.

- 20 A protein of the invention, or a derivative, fragment, analog, homolog or ortholog thereof, may be utilized as an immunogen in the generation of antibodies that immunospecifically bind these protein components.

Various procedures known within the art may be used for the production of polyclonal or monoclonal antibodies directed against a protein of the invention, or against derivatives,  
25 fragments, analogs homologs or orthologs thereof (see, for example, *Antibodies: A Laboratory Manual*, Harlow E, and Lane D, 1988, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, incorporated herein by reference). Some of these antibodies are discussed below.

#### 5.13.1 Polyclonal Antibodies

- 30 For the production of polyclonal antibodies, various suitable host animals (*e.g.*, rabbit, goat, mouse or other mammal) may be immunized by one or more injections with the native protein, a synthetic variant thereof, or a derivative of the foregoing. An appropriate immunogenic preparation can contain, for example, the naturally occurring immunogenic protein, a chemically synthesized polypeptide representing the immunogenic protein, or a

recombinantly expressed immunogenic protein. Furthermore, the protein may be conjugated to a second protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. The preparation can further  
5 include an adjuvant. Various adjuvants used to increase the immunological response include, but are not limited to, Freund's (complete and incomplete), mineral gels (e.g., aluminum hydroxide), surface active substances (e.g., lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, dinitrophenol, etc.), adjuvants usable in humans such as Bacille Calmette-Guerin and *Corynebacterium parvum*, or similar immunostimulatory agents.  
10 Additional examples of adjuvants which can be employed include MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate).

The polyclonal antibody molecules directed against the immunogenic protein can be isolated from the mammal (e.g., from the blood) and further purified by well known techniques, such as affinity chromatography using protein A or protein G, which provide  
15 primarily the IgG fraction of immune serum. Subsequently, or alternatively, the specific antigen which is the target of the immunoglobulin sought, or an epitope thereof, may be immobilized on a column to purify the immune specific antibody by immunoaffinity chromatography. Purification of immunoglobulins is discussed, for example, by D. Wilkinson (The Scientist, published by The Scientist, Inc., Philadelphia PA, Vol. 14, No. 8 (April 17,  
20 2000), pp. 25-28).

### 5.13.2 Monoclonal Antibodies

The term "monoclonal antibody" (MAb) or "monoclonal antibody composition", as used herein, refers to a population of antibody molecules that contain only one molecular  
25 species of antibody molecule consisting of a unique light chain gene product and a unique heavy chain gene product. In particular, the complementarity determining regions (CDRs) of the monoclonal antibody are identical in all the molecules of the population. MAbs thus contain an antigen binding site capable of immunoreacting with a particular epitope of the antigen characterized by a unique binding affinity for it.

30 Monoclonal antibodies can be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes can be immunized in vitro.

The immunizing agent will typically include the protein antigen, a fragment thereof or a fusion protein thereof. Generally, either peripheral blood lymphocytes are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell (Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103). Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells can be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies (Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63).

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against the antigen. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980). Preferably, antibodies having a high degree of specificity and a high binding affinity for the target antigen are isolated.

After the desired hybridoma cells are identified, the clones can be subcloned by limiting dilution procedures and grown by standard methods. Suitable culture media for this purpose

include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells can be grown in vivo as ascites in a mammal.

The monoclonal antibodies secreted by the subclones can be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel  
5 electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies can also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by  
10 using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA can be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to  
15 obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also can be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Patent No. 4,816,567; Morrison, Nature **368**, 812-13 (1994)) or by covalently joining to the immunoglobulin coding  
sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a  
20 non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

### 5.13.2 Humanized Antibodies

25 The antibodies directed against the protein antigens of the invention can further comprise humanized antibodies or human antibodies. These antibodies are suitable for administration to humans without engendering an immune response by the human against the administered immunoglobulin. Humanized forms of antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-  
30 binding subsequences of antibodies) that are principally comprised of the sequence of a human immunoglobulin, and contain minimal sequence derived from a non-human immunoglobulin. Humanization can be performed following the method of Winter and co-workers (Jones et al., Nature, **321**:522-525 (1986); Riechmann et al., Nature, **332**:323-327 (1988); Verhoeyen et al., Science, **239**:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the

corresponding sequences of a human antibody. (See also U.S. Patent No. 5,225,539.) In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies can also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences.

5 In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the framework regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a  
10 human immunoglobulin (Jones et al., 1986; Riechmann et al., 1988; and Presta, Curr. Op. Struct. Biol., 2:593-596 (1992)).

### 5.13.3 Human Antibodies

Fully human antibodies relate to antibody molecules in which essentially the entire  
15 sequences of both the light chain and the heavy chain, including the CDRs, arise from human genes. Such antibodies are termed "human antibodies", or "fully human antibodies" herein. Human monoclonal antibodies can be prepared by the trioma technique; the human B-cell hybridoma technique (see Kozbor, et al., 1983 *Immunol Today* 4: 72) and the EBV hybridoma technique to produce human monoclonal antibodies (see Cole, et al., 1985 In: *MONOCLONAL*  
20 *ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96). Human monoclonal antibodies may be utilized in the practice of the present invention and may be produced by using human hybridomas (see Cote, et al., 1983. *Proc Natl Acad Sci USA* 80: 2026-2030) or by transforming human B-cells with Epstein Barr Virus in vitro (see Cole, et al., 1985 In: *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, Alan R. Liss, Inc., pp. 77-96).

25 In addition, human antibodies can also be produced using additional techniques, including phage display libraries (Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)). Similarly, human antibodies can be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon  
30 challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in Marks et al. (Bio/Technology 10, 779-783 (1992)); Lonberg et al. (Nature 368 856-859 (1994)); Morrison (Nature 368, 812-13 (1994)); Fishwild

et al, (Nature Biotechnology 14, 845-51 (1996)); Neuberger (Nature Biotechnology 14, 826 (1996)); and Lonberg and Huszar (Intern. Rev. Immunol. 13 65-93 (1995)).

Human antibodies may additionally be produced using transgenic nonhuman animals which are modified so as to produce fully human antibodies rather than the animal's  
5 endogenous antibodies in response to challenge by an antigen. (See PCT publication WO94/02602). The endogenous genes encoding the heavy and light immunoglobulin chains in the nonhuman host have been incapacitated, and active loci encoding human heavy and light chain immunoglobulins are inserted into the host's genome. The human genes are incorporated, for example, using yeast artificial chromosomes containing the requisite human  
10 DNA segments. An animal which provides all the desired modifications is then obtained as progeny by crossbreeding intermediate transgenic animals containing fewer than the full complement of the modifications. The preferred embodiment of such a nonhuman animal is a mouse, and is termed the Xenomouse™ as disclosed in PCT publications WO 96/33735 and WO 96/34096. This animal produces B cells which secrete fully human immunoglobulins.  
15 The antibodies can be obtained directly from the animal after immunization with an immunogen of interest, as, for example, a preparation of a polyclonal antibody, or alternatively from immortalized B cells derived from the animal, such as hybridomas producing monoclonal antibodies. Additionally, the genes encoding the immunoglobulins with human variable regions can be recovered and expressed to obtain the antibodies directly, or  
20 can be further modified to obtain analogs of antibodies such as, for example, single chain Fv molecules.

An example of a method of producing a nonhuman host, exemplified as a mouse, lacking expression of an endogenous immunoglobulin heavy chain is disclosed in U.S. Patent No. 5,939,598. It can be obtained by a method including deleting the J segment genes from at  
25 least one endogenous heavy chain locus in an embryonic stem cell to prevent rearrangement of the locus and to prevent formation of a transcript of a rearranged immunoglobulin heavy chain locus, the deletion being effected by a targeting vector containing a gene encoding a selectable marker; and producing from the embryonic stem cell a transgenic mouse whose somatic and germ cells contain the gene encoding the selectable marker.

30 A method for producing an antibody of interest, such as a human antibody, is disclosed in U.S. Patent No. 5,916,771. It includes introducing an expression vector that contains a nucleotide sequence encoding a heavy chain into one mammalian host cell in culture, introducing an expression vector containing a nucleotide sequence encoding a light chain into

another mammalian host cell, and fusing the two cells to form a hybrid cell. The hybrid cell expresses an antibody containing the heavy chain and the light chain.

In a further improvement on this procedure, a method for identifying a clinically relevant epitope on an immunogen, and a correlative method for selecting an antibody that  
5 binds immunospecifically to the relevant epitope with high affinity, are disclosed in PCT publication WO 99/53049.

#### 5.13.4 F<sub>ab</sub> Fragments and Single Chain Antibodies

According to the invention, techniques can be adapted for the production of  
10 single-chain antibodies specific to an antigenic protein of the invention (see e.g., U.S. Patent No. 4,946,778). In addition, methods can be adapted for the construction of F<sub>ab</sub> expression libraries (see e.g., Huse, et al., 1989 Science 246: 1275-1281) to allow rapid and effective identification of monoclonal F<sub>ab</sub> fragments with the desired specificity for a protein or derivatives, fragments, analogs or homologs thereof. Antibody fragments that contain the  
15 idiotypes to a protein antigen may be produced by techniques known in the art including, but not limited to: (i) an F<sub>(ab')<sub>2</sub></sub> fragment produced by pepsin digestion of an antibody molecule; (ii) an F<sub>ab</sub> fragment generated by reducing the disulfide bridges of an F<sub>(ab')<sub>2</sub></sub> fragment; (iii) an F<sub>ab</sub> fragment generated by the treatment of the antibody molecule with papain and a reducing agent  
and (iv) F<sub>v</sub> fragments.

20

#### 5.13.5 Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for an antigenic protein of the invention. The second binding target is  
25 any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities (Milstein and Cuello, Nature, 305:537-539 (1983)). Because of the random  
30 assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published 13 May 1993, and in Traunecker *et al.*, 1991 *EMBO J.*, 10:3655-3659.

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology, 121:210 (1986).

According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')<sub>2</sub> bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared using chemical linkage. Brennan et al., Science 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')<sub>2</sub> fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Additionally, Fab' fragments can be directly recovered from E. coli and chemically coupled to form bispecific antibodies. Shalaby et al., J. Exp. Med. 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')<sub>2</sub> molecule. Each Fab'



fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

5           Various techniques for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., J. Immunol. 148(5):1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge  
10           region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., Proc. Natl. Acad. Sci. USA 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain ( $V_H$ ) connected to a light-chain variable  
15           domain ( $V_L$ ) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the  $V_H$  and  $V_L$  domains of one fragment are forced to pair with the complementary  $V_L$  and  $V_H$  domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., J. Immunol. 152:5368 (1994).

20           Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., J. Immunol. 147:60 (1991). Exemplary bispecific antibodies can bind to two different epitopes, at least one of which originates in the protein antigen of the invention. Alternatively, an anti-antigenic arm of an immunoglobulin molecule can be combined with an arm which binds to a triggering molecule  
25           on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (Fc  $\gamma$ R), such as Fc  $\gamma$ RI (CD64), Fc  $\gamma$ RII (CD32) and Fc  $\gamma$ RIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular antigen. Bispecific antibodies can also be used to direct cytotoxic agents to cells which express a particular antigen. These antibodies possess an antigen-binding arm and an arm which binds a cytotoxic  
30           agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the protein antigen described herein and further binds tissue factor (TF).

### 5.13.6 Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention.

Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (WO 91/00360; WO 92/200373; EP 03089). It is contemplated that the antibodies can be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins can be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Patent No. 4,676,980.

### 5.13.7 Effector Function Engineering

It can be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) can be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated can have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., J. Exp Med., 176: 1191-1195 (1992) and Shopes, J. Immunol., 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity can also be prepared using heterobifunctional cross-linkers as described in Wolff et al. Cancer Research, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and can thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., Anti-Cancer Drug Design, 3: 219-230 (1989).

### 5.13.8 Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolaca americana proteins (PAPI, PAPII, and

PAP-S), momordica charantia inhibitor, curcin, croton, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include  $^{212}\text{Bi}$ ,  $^{131}\text{I}$ ,  $^{131}\text{In}$ ,  $^{90}\text{Y}$ , and  $^{186}\text{Re}$ .

5 Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimide HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives  
10 (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., Science, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of  
15 radionucleotide to the antibody. See WO94/11026.

In another embodiment, the antibody can be conjugated to a "receptor" (such as streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is in turn  
20 conjugated to a cytotoxic agent.

#### 5.14 COMPUTER READABLE SEQUENCES

In one application of this embodiment, a nucleotide sequence of the present invention can be recorded on computer readable media. As used herein, "computer readable media"  
25 refers to any medium which can be read and accessed directly by a computer. Such media include, but are not limited to: magnetic storage media, such as floppy discs, hard disc storage medium, and magnetic tape; optical storage media such as CD-ROM; electrical storage media such as RAM and ROM; and hybrids of these categories such as magnetic/optical storage media. A skilled artisan can readily appreciate how any of the presently known computer  
30 readable mediums can be used to create a manufacture comprising computer readable medium having recorded thereon a nucleotide sequence of the present invention. As used herein, "recorded" refers to a process for storing information on computer readable medium. A skilled artisan can readily adopt any of the presently known methods for recording information

on computer readable medium to generate manufactures comprising the nucleotide sequence information of the present invention.

A variety of data storage structures are available to a skilled artisan for creating a computer readable medium having recorded thereon a nucleotide sequence of the present invention. The choice of the data storage structure will generally be based on the means chosen to access the stored information. In addition, a variety of data processor programs and formats can be used to store the nucleotide sequence information of the present invention on computer readable medium. The sequence information can be represented in a word processing text file, formatted in commercially-available software such as WordPerfect and Microsoft Word, or represented in the form of an ASCII file, stored in a database application, such as DB2, Sybase, Oracle, or the like. A skilled artisan can readily adapt any number of data processor structuring formats (*e.g.* text file or database) in order to obtain computer readable medium having recorded thereon the nucleotide sequence information of the present invention.

By providing any of the nucleotide sequences SEQ ID NO: 1-3, 10-13 or 15 or a representative fragment thereof; or a nucleotide sequence at least 95% identical to any of the nucleotide sequences of the SEQ ID NO:1-3, 10-13 or 15 in computer readable form, a skilled artisan can routinely access the sequence information for a variety of purposes. Computer software is publicly available which allows a skilled artisan to access sequence information provided in a computer readable medium. The examples which follow demonstrate how software which implements the BLAST (Altschul et al., J. Mol. Biol. 215:403-410 (1990)) and BLAZE (Brutlag et al., Comp. Chem. 17:203-207 (1993)) search algorithms on a Sybase system is used to identify open reading frames (ORFs) within a nucleic acid sequence. Such ORFs may be protein encoding fragments and may be useful in producing commercially important proteins such as enzymes used in fermentation reactions and in the production of commercially useful metabolites.

As used herein, "a computer-based system" refers to the hardware means, software means, and data storage means used to analyze the nucleotide sequence information of the present invention. The minimum hardware means of the computer-based systems of the present invention comprises a central processing unit (CPU), input means, output means, and data storage means. A skilled artisan can readily appreciate that any one of the currently available computer-based systems are suitable for use in the present invention. As stated above, the computer-based systems of the present invention comprise a data storage means having stored therein a nucleotide sequence of the present invention and the necessary hardware means

and software means for supporting and implementing a search means. As used herein, "data storage means" refers to memory which can store nucleotide sequence information of the present invention, or a memory access means which can access manufactures having recorded thereon the nucleotide sequence information of the present invention.

5 As used herein, "search means" refers to one or more programs which are implemented on the computer-based system to compare a target sequence or target structural motif with the sequence information stored within the data storage means. Search means are used to identify fragments or regions of a known sequence which match a particular target sequence or target motif. A variety of known algorithms are disclosed publicly and a variety of commercially  
10 available software for conducting search means are and can be used in the computer-based systems of the present invention. Examples of such software includes, but is not limited to, Smith-Waterman, MacPattern (EMBL), BLASTN and BLASTA (NPOLYPEPTIDEIA). A skilled artisan can readily recognize that any one of the available algorithms or implementing software packages for conducting homology searches can be adapted for use in the present  
15 computer-based systems. As used herein, a "target sequence" can be any nucleic acid or amino acid sequence of six or more nucleotides or two or more amino acids. A skilled artisan can readily recognize that the longer a target sequence is, the less likely a target sequence will be present as a random occurrence in the database. The most preferred sequence length of a target sequence is from about 10 to 100 amino acids, or from about 30 to 300 nucleotide  
20 residues. However, it is well recognized that searches for commercially important fragments, such as sequence fragments involved in gene expression and protein processing, may be of shorter length.

As used herein, "a target structural motif," or "target motif," refers to any rationally selected sequence or combination of sequences in which the sequence(s) are chosen based on a  
25 three-dimensional configuration which is formed upon the folding of the target motif. There are a variety of target motifs known in the art. Protein target motifs include, but are not limited to, enzyme active sites and signal sequences. Nucleic acid target motifs include, but are not limited to, promoter sequences, hairpin structures and inducible expression elements (protein binding sequences).

30

### 5.15 TRIPLE HELIX FORMATION

In addition, the fragments of the present invention, as broadly described, can be used to control gene expression through triple helix formation or antisense DNA or RNA, both of which methods are based on the binding of a polynucleotide sequence to DNA or RNA.

Polynucleotides suitable for use in these methods are usually 20 to 40 bases in length and are designed to be complementary to a region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 15241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Olmno, J. Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Triple helix-formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the present invention is necessary for the design of an antisense or triple helix oligonucleotide.

### 5.16 DIAGNOSTIC ASSAYS AND KITS

The present invention further provides methods to identify the presence or expression of one of the ORFs of the present invention, or homolog thereof, in a test sample, using a nucleic acid probe or antibodies of the present invention, optionally conjugated or otherwise associated with a suitable label.

In general, methods for detecting a polynucleotide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the polynucleotide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polynucleotide of the invention is detected in the sample. Such methods can also comprise contacting a sample under stringent hybridization conditions with nucleic acid primers that anneal to a polynucleotide of the invention under such conditions, and amplifying annealed polynucleotides, so that if a polynucleotide is amplified, a polynucleotide of the invention is detected in the sample.

In general, methods for detecting a polypeptide of the invention can comprise contacting a sample with a compound that binds to and forms a complex with the polypeptide for a period sufficient to form the complex, and detecting the complex, so that if a complex is detected, a polypeptide of the invention is detected in the sample.

In detail, such methods comprise incubating a test sample with one or more of the antibodies or one or more of the nucleic acid probes of the present invention and assaying for binding of the nucleic acid probes or antibodies to components within the test sample.

Conditions for incubating a nucleic acid probe or antibody with a test sample vary. Incubation conditions depend on the format employed in the assay, the detection methods employed, and the type and nature of the nucleic acid probe or antibody used in the assay.

One skilled in the art will recognize that any one of the commonly available hybridization, amplification or immunological assay formats can readily be adapted to employ the nucleic acid probes or antibodies of the present invention. Examples of such assays can be found in Chard, T., An Introduction to Radioimmunoassay and Related Techniques, Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock, G.R. et al., Techniques in Immunocytochemistry, Academic Press, Orlando, FL Vol. 1 (1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, P., Practice and Theory of immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands (1985). The test samples of the present invention include cells, protein or membrane extracts of cells, or biological fluids such as sputum, blood, serum, plasma, or urine. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing protein extracts or membrane extracts of cells are well known in the art and can be readily be adapted in order to obtain a sample which is compatible with the system utilized.

In another embodiment of the present invention, kits are provided which contain the necessary reagents to carry out the assays of the present invention. Specifically, the invention provides a compartment kit to receive, in close confinement, one or more containers which comprises: (a) a first container comprising one of the probes or antibodies of the present invention; and (b) one or more other containers comprising one or more of the following: wash reagents, reagents capable of detecting presence of a bound probe or antibody.

In detail, a compartment kit includes any kit in which reagents are contained in separate containers. Such containers include small glass containers, plastic containers or strips of plastic or paper. Such containers allows one to efficiently transfer reagents from one compartment to another compartment such that the samples and reagents are not cross-contaminated, and the agents or solutions of each container can be added in a quantitative fashion from one compartment to another. Such containers will include a container which will accept the test sample, a container which contains the antibodies used in the assay, containers which contain wash reagents (such as phosphate buffered saline, Tris-buffers, etc.), and containers which contain the reagents used to detect the bound antibody or probe. Types of detection reagents include labeled nucleic acid probes, labeled secondary antibodies, or in the alternative, if the primary antibody is labeled, the enzymatic, or antibody binding reagents which are capable of reacting with the labeled antibody. One skilled in the art will readily

recognize that the disclosed probes and antibodies of the present invention can be readily incorporated into one of the established kit formats which are well known in the art.

### 5.17 MEDICAL IMAGING

5           The novel polypeptides and binding partners of the invention are useful in medical imaging of sites expressing the molecules of the invention (e.g., where the polypeptide of the invention is involved in the immune response, for imaging sites of inflammation or infection). See, e.g., Kunkel et al., U.S. Pat. NO. 5,413,778. Such methods involve chemical attachment of a labeling or imaging agent, administration of the labeled polypeptide to a subject in a  
10   pharmaceutically acceptable carrier, and imaging the labeled polypeptide *in vivo* at the target site.

### 5.18 SCREENING ASSAYS

          Using the isolated proteins and polynucleotides of the invention, the present invention  
15   further provides methods of obtaining and identifying agents which bind to a polypeptide encoded by an ORF corresponding to any of the nucleotide sequences set forth in the SEQ ID NO: 1-3, 10-13 or 15, or bind to a specific domain of the polypeptide encoded by the nucleic acid. In detail, said method comprises the steps of:

- (a)   contacting an agent with an isolated protein encoded by an ORF of the present  
20   invention, or nucleic acid of the invention; and
- (b)   determining whether the agent binds to said protein or said nucleic acid.

          In general, therefore, such methods for identifying compounds that bind to a polynucleotide of the invention can comprise contacting a compound with a polynucleotide of the invention for a time sufficient to form a polynucleotide/compound complex, and detecting  
25   the complex, so that if a polynucleotide/compound complex is detected, a compound that binds to a polynucleotide of the invention is identified.

          Likewise, in general, therefore, such methods for identifying compounds that bind to a polypeptide of the invention can comprise contacting a compound with a polypeptide of the invention for a time sufficient to form a polypeptide/compound complex, and detecting the  
30   complex, so that if a polypeptide/compound complex is detected, a compound that binds to a polynucleotide of the invention is identified.

          Methods for identifying compounds that bind to a polypeptide of the invention can also comprise contacting a compound with a polypeptide of the invention in a cell for a time sufficient to form a polypeptide/compound complex, wherein the complex drives expression of



a receptor gene sequence in the cell, and detecting the complex by detecting reporter gene sequence expression, so that if a polypeptide/compound complex is detected, a compound that binds a polypeptide of the invention is identified.

Compounds identified via such methods can include compounds which modulate the activity of a polypeptide of the invention (that is, increase or decrease its activity, relative to activity observed in the absence of the compound). Alternatively, compounds identified via such methods can include compounds which modulate the expression of a polynucleotide of the invention (that is, increase or decrease expression relative to expression levels observed in the absence of the compound). Compounds, such as compounds identified via the methods of the invention, can be tested using standard assays well known to those of skill in the art for their ability to modulate activity/expression.

The agents screened in the above assay can be, but are not limited to, peptides, carbohydrates, vitamin derivatives, or other pharmaceutical agents. The agents can be selected and screened at random or rationally selected or designed using protein modeling techniques.

For random screening, agents such as peptides, carbohydrates, pharmaceutical agents and the like are selected at random and are assayed for their ability to bind to the protein encoded by the ORF of the present invention. Alternatively, agents may be rationally selected or designed. As used herein, an agent is said to be "rationally selected or designed" when the agent is chosen based on the configuration of the particular protein. For example, one skilled in the art can readily adapt currently available procedures to generate peptides, pharmaceutical agents and the like, capable of binding to a specific peptide sequence, in order to generate rationally designed antipeptide peptides, for example see Hurby et al., Application of Synthetic Peptides: Antisense Peptides," In Synthetic Peptides, A User's Guide, W.H. Freeman, NY (1992), pp. 289-307, and Kaspczak et al., Biochemistry 28:9230-8 (1989), or pharmaceutical agents, or the like.

In addition to the foregoing, one class of agents of the present invention, as broadly described, can be used to control gene expression through binding to one of the ORFs or EMFs of the present invention. As described above, such agents can be randomly screened or rationally designed/selected. Targeting the ORF or EMF allows a skilled artisan to design sequence specific or element specific agents, modulating the expression of either a single ORF or multiple ORFs which rely on the same EMF for expression control. One class of DNA binding agents are agents which contain base residues which hybridize or form a triple helix formation by binding to DNA or RNA. Such agents can be based on the classic

phosphodiester, ribonucleic acid backbone, or can be a variety of sulfhydryl or polymeric derivatives which have base attachment capacity.

Agents suitable for use in these methods usually contain 20 to 40 bases and are designed to be complementary to a region of the gene involved in transcription (triple helix -  
5 see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Triple helix-formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks  
10 translation of an mRNA molecule into polypeptide. Both techniques have been demonstrated to be effective in model systems. Information contained in the sequences of the present invention is necessary for the design of an antisense or triple helix oligonucleotide and other DNA binding agents.

Agents which bind to a protein encoded by one of the ORFs of the present invention  
15 can be used as a diagnostic agent. Agents which bind to a protein encoded by one of the ORFs of the present invention can be formulated using known techniques to generate a pharmaceutical composition.

#### 5.19 USE OF NUCLEIC ACIDS AS PROBES

20 Another aspect of the subject invention is to provide for polypeptide-specific nucleic acid hybridization probes capable of hybridizing with naturally occurring nucleotide sequences. The hybridization probes of the subject invention may be derived from any of the nucleotide sequences SEQ ID NO:1-3, 10-13 or 15. Because the corresponding gene is only expressed in a limited number of tissues, a hybridization probe derived from any of the nucleotide  
25 sequences SEQ ID NO: 1-3, 10-13 and 15 can be used as an indicator of the presence of RNA of cell type of such a tissue in a sample.

Any suitable hybridization technique can be employed, such as, for example, in situ hybridization. PCR as described in US Patents Nos. 4,683,195 and 4,965,188 provides additional uses for oligonucleotides based upon the nucleotide sequences. Such probes used in  
30 PCR may be of recombinant origin, may be chemically synthesized, or a mixture of both. The probe will comprise a discrete nucleotide sequence for the detection of identical sequences or a degenerate pool of possible sequences for identification of closely related genomic sequences.

Other means for producing specific hybridization probes for nucleic acids include the cloning of nucleic acid sequences into vectors for the production of mRNA probes. Such

vectors are known in the art and are commercially available and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerase as T7 or SP6 RNA polymerase and the appropriate radioactively labeled nucleotides. The nucleotide sequences may be used to construct hybridization probes for mapping their respective genomic sequences. The nucleotide sequence provided herein may be mapped to a chromosome or specific regions of a chromosome using well known genetic and/or chromosomal mapping techniques. These techniques include *in situ* hybridization, linkage analysis against known chromosomal markers, hybridization screening with libraries or flow-sorted chromosomal preparations specific to known chromosomes, and the like. The technique of fluorescent *in situ* hybridization of chromosome spreads has been described, among other places, in Verma et al (1988) Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York NY.

Fluorescent *in situ* hybridization of chromosomal preparations and other physical chromosome mapping techniques may be correlated with additional genetic map data. Examples of genetic map data can be found in the 1994 Genome Issue of Science (265:1981f). Correlation between the location of a nucleic acid on a physical chromosomal map and a specific disease (or predisposition to a specific disease) may help delimit the region of DNA associated with that genetic disease. The nucleotide sequences of the subject invention may be used to detect differences in gene sequences between normal, carrier or affected individuals.

## 5.20 PREPARATION OF SUPPORT BOUND OLIGONUCLEOTIDES

Oligonucleotides, i.e., small nucleic acid segments, may be readily prepared by, for example, directly synthesizing the oligonucleotide by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer.

Support bound oligonucleotides may be prepared by any of the methods known to those of skill in the art using any suitable support such as glass, polystyrene or Teflon. One strategy is to precisely spot oligonucleotides synthesized by standard synthesizers. Immobilization can be achieved using passive adsorption (Inouye & Hondo, 1990 J. Clin Microbiol 28(6) 1462-72); using UV light (Nagata *et al.*, 1985; Dahlen *et al.*, 1987; Morrissey & Collins, Mol. Cell Probes 1989 3(2) 189-207) or by covalent binding of base modified DNA (Keller *et al.*, 1988; 1989); all references being specifically incorporated herein.

Another strategy that may be employed is the use of the strong biotin-streptavidin interaction as a linker. For example, Broude *et al.* (1994) Proc. Natl. Acad. Sci USA 91(8) 3072-6 describe the use of biotinylated probes, although these are duplex probes, that are

immobilized on streptavidin-coated magnetic beads. Streptavidin-coated beads may be purchased from Dynal, Oslo. Of course, this same linking chemistry is applicable to coating any surface with streptavidin. Biotinylated probes may be purchased from various sources, such as, e.g., Operon Technologies (Alameda, CA).

5           Nunc Laboratories (Naperville, IL) is also selling suitable material that could be used. Nunc Laboratories have developed a method by which DNA can be covalently bound to the microwell surface termed CovaLink NH. CovaLink NH is a polystyrene surface grafted with secondary amino groups (>NH) that serve as bridge-heads for further covalent coupling. CovaLink Modules may be purchased from Nunc Laboratories. DNA molecules may be bound  
10 to CovaLink exclusively at the 5'-end by a phosphoramidate bond, allowing immobilization of more than 1 pmol of DNA (Rasmussen *et al.*, (1991) *Anal Biochem* 198(1) 138-42.

The use of CovaLink NH strips for covalent binding of DNA molecules at the 5'-end has been described (Rasmussen *et al.*, 1991). In this technology, a phosphoramidate bond is employed (Chu *et al.*, 1983 *Nucleic Acids* 11(18) 6513-29). This is beneficial as immobilization  
15 using only a single covalent bond is preferred. The phosphoramidate bond joins the DNA to the CovaLink NH secondary amino groups that are positioned at the end of spacer arms covalently grafted onto the polystyrene surface through a 2 nm long spacer arm. To link an oligonucleotide to CovaLink NH via an phosphoramidate bond, the oligonucleotide terminus must have a 5'-end phosphate group. It is, perhaps, even possible for biotin to be covalently bound to CovaLink and  
20 then streptavidin used to bind the probes.

More specifically, the linkage method includes dissolving DNA in water (7.5 ng/ul) and denaturing for 10 min. at 95°C and cooling on ice for 10 min. Ice-cold 0.1 M 1-methylimidazole, pH 7.0 (1-MeIm), is then added to a final concentration of 10 mM 1-MeIm. A ss DNA solution is then dispensed into CovaLink NH strips (75 ul/well) standing on ice.

25           Carbodiimide 0.2 M 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide(EDC), dissolved in 10 mM 1-MeIm, is made fresh and 25 ul added per well. The strips are incubated for 5 hours at 50°C. After incubation the strips are washed using, e.g., Nunc-Immuno Wash; first the wells are washed 3 times, then they are soaked with washing solution for 5 min., and finally they are washed 3 times (where in the washing solution is 0.4 N NaOH, 0.25% SDS heated to 50°C).

30           It is contemplated that a further suitable method for use with the present invention is that described in PCT Patent Application WO 90/03382 (Southern & Maskos), incorporated herein by reference. This method of preparing an oligonucleotide bound to a support involves attaching a nucleoside 3'-reagent through the phosphate group by a covalent phosphodiester link to aliphatic hydroxyl groups carried by the support. The oligonucleotide is then synthesized on the supported

nucleoside and protecting groups removed from the synthetic oligonucleotide chain under standard conditions that do not cleave the oligonucleotide from the support. Suitable reagents include nucleoside phosphoramidite and nucleoside hydrogen phosphate.

An on-chip strategy for the preparation of DNA probe for the preparation of DNA probe  
5 arrays may be employed. For example, addressable laser-activated photodeprotection may be employed in the chemical synthesis of oligonucleotides directly on a glass surface, as described by Fodor *et al.* (1991) *Science* 251(4995) 767-73, incorporated herein by reference. Probes may also be immobilized on nylon supports as described by Van Ness *et al.* (1991) *Nucleic Acids Res.* 19(12) 3345-50; or linked to Teflon using the method of Duncan & Cavalier (1988) *Anal*  
10 *Biochem* 169(1) 104-8; all references being specifically incorporated herein.

To link an oligonucleotide to a nylon support, as described by Van Ness *et al.* (1991), requires activation of the nylon surface via alkylation and selective activation of the 5'-amine of oligonucleotides with cyanuric chloride.

One particular way to prepare support bound oligonucleotides is to utilize the light-  
15 generated synthesis described by Pease *et al.*, (1994) *Proc. Natl. Acad. Sci USA* 91(11) 5022-6. These authors used current photolithographic techniques to generate arrays of immobilized oligonucleotide probes (DNA chips). These methods, in which light is used to direct the synthesis of oligonucleotide probes in high-density, miniaturized arrays, utilize photolabile 5'-protected *N*-acyl-deoxynucleoside phosphoramidites, surface linker chemistry and versatile combinatorial  
20 synthesis strategies. A matrix of 256 spatially defined oligonucleotide probes may be generated in this manner.

## 5.21 PREPARATION OF NUCLEIC ACID FRAGMENTS

The nucleic acids may be obtained from any appropriate source, such as cDNAs, genomic DNA, chromosomal DNA, microdissected chromosome bands, cosmid or YAC inserts, and  
25 RNA, including mRNA without any amplification steps. For example, Sambrook *et al.* (1989) describes three protocols for the isolation of high molecular weight DNA from mammalian cells (p. 9.14-9.23).

DNA fragments may be prepared as clones in M13, plasmid or lambda vectors and/or prepared directly from genomic DNA or cDNA by PCR or other amplification methods. Samples  
30 may be prepared or dispensed in multiwell plates. About 100-1000 ng of DNA samples may be prepared in 2-500 ml of final volume.

The nucleic acids would then be fragmented by any of the methods known to those of skill in the art including, for example, using restriction enzymes as described at 9.24-9.28 of Sambrook *et al.* (1989), shearing by ultrasound and NaOH treatment.

Low pressure shearing is also appropriate, as described by Schriefer *et al.* (1990) Nucleic  
5 Acids Res. 18(24) 7455-6. In this method, DNA samples are passed through a small French pressure cell at a variety of low to intermediate pressures. A lever device allows controlled application of low to intermediate pressures to the cell. The results of these studies indicate that low-pressure shearing is a useful alternative to sonic and enzymatic DNA fragmentation methods.

One particularly suitable way for fragmenting DNA is contemplated to be that using the  
10 two base recognition endonuclease, CviJI, described by Fitzgerald *et al.* (1992) Nucleic Acids Res. 20(14) 3753-62. These authors described an approach for the rapid fragmentation and fractionation of DNA into particular sizes that they contemplated to be suitable for shotgun cloning and sequencing.

The restriction endonuclease CviJI normally cleaves the recognition sequence PuGCPy  
15 between the G and C to leave blunt ends. Atypical reaction conditions, which alter the specificity of this enzyme (CviJI\*\*), yield a quasi-random distribution of DNA fragments from the small molecule pUC19 (2688 base pairs). Fitzgerald *et al.* (1992) quantitatively evaluated the randomness of this fragmentation strategy, using a CviJI\*\* digest of pUC19 that was size fractionated by a rapid gel filtration method and directly ligated, without end repair, to a lac Z  
20 minus M13 cloning vector. Sequence analysis of 76 clones showed that CviJI\*\* restricts pyGCPy and PuGCPu, in addition to PuGCPy sites, and that new sequence data is accumulated at a rate consistent with random fragmentation.

As reported in the literature, advantages of this approach compared to sonication and agarose gel fractionation include: smaller amounts of DNA are required (0.2-0.5 ug instead of 2-  
25 5 ug); and fewer steps are involved (no preligation, end repair, chemical extraction, or agarose gel electrophoresis and elution are needed).

Irrespective of the manner in which the nucleic acid fragments are obtained or prepared, it is important to denature the DNA to give single stranded pieces available for hybridization. This is achieved by incubating the DNA solution for 2-5 minutes at 80-90°C. The solution is then  
30 cooled quickly to 2°C to prevent renaturation of the DNA fragments before they are contacted with the chip. Phosphate groups must also be removed from genomic DNA by methods known in the art.

## 5.22 PREPARATION OF DNA ARRAYS

Arrays may be prepared by spotting DNA samples on a support such as a nylon membrane. Spotting may be performed by using arrays of metal pins (the positions of which correspond to an array of wells in a microtiter plate) to repeated by transfer of about 20 nl of a DNA solution to a nylon membrane. By offset printing, a density of dots higher than the density of the wells is achieved. One to 25 dots may be accommodated in 1 mm<sup>2</sup>, depending on the type of label used. By avoiding spotting in some preselected number of rows and columns, separate subsets (subarrays) may be formed. Samples in one subarray may be the same genomic segment of DNA (or the same gene) from different individuals, or may be different, overlapped genomic clones. Each of the subarrays may represent replica spotting of the same samples. In one example, a selected gene segment may be amplified from 64 patients. For each patient, the amplified gene segment may be in one 96-well plate (all 96 wells containing the same sample). A plate for each of the 64 patients is prepared. By using a 96-pin device, all samples may be spotted on one 8 x 12 cm membrane. Subarrays may contain 64 samples, one from each patient. Where the 96 subarrays are identical, the dot span may be 1 mm<sup>2</sup> and there may be a 1 mm space between subarrays.

Another approach is to use membranes or plates (available from NUNC, Naperville, Illinois) which may be partitioned by physical spacers e.g. a plastic grid molded over the membrane, the grid being similar to the sort of membrane applied to the bottom of multiwell plates, or hydrophobic strips. A fixed physical spacer is not preferred for imaging by exposure to flat phosphor-storage screens or x-ray films.

The present invention is illustrated in the following examples. Upon consideration of the present disclosure, one of skill in the art will appreciate that many other embodiments and variations may be made in the scope of the present invention. Accordingly, it is intended that the broader aspects of the present invention not be limited to the disclosure of the following examples. The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the invention, and compositions and methods which are functionally equivalent are within the scope of the invention. Indeed, numerous modifications and variations in the practice of the invention are expected to occur to those skilled in the art upon consideration of the present preferred embodiments. Consequently, the only limitations which should be placed upon the scope of the invention are those which appear in the appended claims.

All references cited within the body of the instant specification are hereby incorporated by reference in their entirety.

## 6.0 EXAMPLES

### EXAMPLE 1

#### Isolation of SEQ ID NO: 1 from a cDNA Library of Human Leukocytes

5 A plurality of novel nucleic acids were obtained from a cDNA library prepared from human ovary (Hyseq clone identification numbers 7735656 (SEQ ID NO: 1) and 7727065 (SEQ ID NO: 11)) using standard PCR, sequencing by hybridization sequence signature analysis, and Sanger sequencing techniques. The inserts of the library were amplified with PCR using primers specific for vector sequences flanking the inserts. These samples were  
10 spotted onto nylon membranes and interrogated with oligonucleotide probes to give sequence signatures. The clones were clustered into groups of similar or identical sequences, and single representative clones were selected from each group for gel sequencing. The 5' sequence of the amplified inserts was then deduced using the reverse M13 sequencing primer in a typical Sanger sequencing protocol. PCR products were purified and subjected to fluorescent dye  
15 terminator cycle sequencing. Single-pass gel sequencing was done using a 377 Applied Biosystems (ABI) sequencer. The insert was identified as a novel sequence not previously obtained from this library and not previously reported in public databases. The sequences were designated as SEQ ID NO: 1 and SEQ ID NO:11.

20

### EXAMPLE 2

#### ASSEMBLAGE OF SEQ ID NO: 2 AND SEQ ID NO: 12

The nucleic acids of the present invention, designated as SEQ ID NO: 2 and SEQ ID  
25 NO:12 were assembled using an EST sequence, SEQ ID NO:1 or SEQ ID NO:11, respectively, as a seed. Then a recursive algorithm was used to extend the seed into an extended assemblage, by pulling additional sequences from different databases (i.e., Hyseq's database containing EST sequences, dbEST version 114, gb pri 114, and UniGene version 101) that belong to this assemblage. The algorithm terminated when there was no additional sequences from the above  
30 databases that would extend the assemblage. Inclusion of component sequences into the assemblage was based on a BLASTN hit to the extending assemblage with BLAST score greater than 300 and percent identity greater than 95%.

The nearest neighbor result for the assembled contigs were obtained by a FASTA version 3 search against Genpept release 114, using FASTXY algorithm. FASTXY is an improved  
35 version of FASTA alignment which allows in-codon frame shifts. The nearest neighbor result



showed the closest homologue for each assemblage from Genpept (and contains the translated amino acid sequences for which the assemblage encodes). The nearest neighbor results is set forth below:

5

SEQ ID NO:	Accession No.	Description	Smith- Waterman Score	% Identity
2	Z36531	Homo sapiens fibrinogen-like protein	500	38.462
12	U44830	Rana sylvatica extended fibrinogen alpha chain	252	28.629

The nucleotide sequence within the assembled contigs that codes for signal peptide sequences and their cleavage sites can be determined from using Neural Network SignalP V1.1 program (from Center for Biological Sequence Analysis, The Technical University of Denmark). The process for identifying prokaryotic and eukaryotic signal peptides and their cleavage sites are also disclosed by Henrik Nielson, Jacob Engelbrecht, Soren Brunak, and Gunnar von Heijne in the publication "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites" Protein Engineering, vol. 10, no. 1, pp. 1-6 (1997), incorporated herein by reference. A maximum S score and a mean S score, as described in the Nielson et al reference, are obtained for each assembled contig. Starting from the first amino acid of the predicted signal sequence, a sequence of 45 amino acids is described. Not all forty-five amino acids in the sequence may comprise the signal peptide.

20 For SEQ ID NO: 12:

Beginning and end nucleotide location corresponding to amino acid segment	Maximum S score	mean S score	Amino acid segment containing signal peptide (A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop Codon,

			/=possible nucleotide deletion, \=possible nucleotide insertion)
54:189	0.922	0.765	MMSPSQASLLFLNVCIFICGEVVQGN CVHHSTDSSVVNIVEDGSN (SEQ ID NO:22)

The predicted amino acid sequence for SEQ ID NO:2 was obtained by using a software program called FASTY (available from <http://fasta.bioch.virginia.edu>) which selects a polypeptide based on a comparison of translated novel polynucleotide to known polynucleotides (W.R. Pearson, Methods in Enzymology, 183:63-98 (1990), incorporated herein by reference).

For SEQ ID NO: 2:

Predicted beginning nucleotide location corresponding to first amino acid residue of amino acid segment	Predicted end nucleotide location corresponding to last amino acid residue of amino acid segment	AMINO ACID SEGMENT (A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine, S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine, X=Unknown, *=Stop Codon, /=possible nucleotide deletion, \=possible nucleotide insertion)
3	717	DLKDTIGSVTKTPSGLYIIHPEGSSYPFEVMCDMDYR GGGWTVIQKRIDGIIDFQRLWCDYLDGFGDLLGEF WLGLKKIFYIVNQNTSFMLYVALESEDDTLAYAS YDNFWLEDETRFFKMHLGRYSGNAGDAFRGLKKE DNQNAMPFSTSDVDNDGCRPACLVNGQSVKSCSHL HNKTGWWFNECGLANLNGIHHFSGKLLATGIQWG TWTKNNSPVKIKSVSMKIRRMYPYKF (SEQ ID NO:21)

10

### EXAMPLE 3

#### ASSEMBLAGE OF SEQ ID NO: 3

Assembly of novel nucleotide sequence of SEQ ID NO: 3 was accomplished by using an EST sequence SEQ ID NO: 1 as a seed. The seed was extended by using software programs such as BLAST and Hyseq proprietary software to pull additional sequences from Hyseq's proprietary database containing EST sequences and by gel sequencing (377 Applied Biosystems (ABI) sequencer) using primers to extend the 3' end. Inclusion of component sequences into the assemblage was based BLAST scores greater than 1000 and a p-value of p-3

15

(depending on the length of homology). The 5' end was extended using RACE, as disclosed in Marathon-Ready™ cDNA User Manual (PT1156-1) (Clontech), herein incorporated by reference.

5

## EXAMPLE 4

ASSEMBLAGE OF SEQ ID NO: 10

Assembly of novel nucleotide sequence of SEQ ID NO: 10 was accomplished by using an EST sequence SEQ ID NO: 1 as a seed. The seed was further extended by gel sequencing (377 Applied Biosystems (ABI) sequencer) using primers to extend the 3' end (primer extension). The 5' end was extended using RACE, as disclosed in Marathon-Ready™ cDNA User Manual (PT1156-1) (Clontech), herein incorporated by reference.

15

## EXAMPLE 5

ASSEMBLAGE OF SEQ ID NO: 13

Using PHRAP (Univ. of Washington) or CAP4 (Paracel), a full length gene cDNA sequence and its corresponding protein sequence were generated from the assemblage of SEQ ID NO:12. Any frame shifts and incorrect stop codons were corrected by hand editing. During editing, the sequence was checked using FASTY and/or BLAST against Genbank (i.e. Genepept release 115). Other computer programs which may have been used in the editing process were phredPhrap and Consed (University of Washington) and ed\_ready, ed\_ext and cg\_zip-2 (Hyseq, Inc.). The full-length nucleotide sequence is shown as SEQ ID NO: 13.

A polypeptide (SEQ ID NO: 14) was predicted to be encoded by SEQ ID NO: 13 as set forth below. The polypeptide was predicted using software programs called BLASTX/FASTY which selects a polypeptide based on a comparison of translated novel polynucleotide to known polypeptides. The initial methionine ATG starts at position 54 of the nucleotide sequence and the putative stop codon, TAA, ends the coding region at position 1034.

30

## EXAMPLE 6

A. Expression of SEQ ID NO: 4-9, 14 and 16-18 in cells

Chinese Hamster Ovary (CHO) cells or other suitable cell types are grown in DMEM (ATCC) and 10% fetal bovine serum (FBS) (Gibco) to 70% confluence. Prior to transfection the media is changed to DMEM and 0.5% FCS. Cells are transfected with cDNAs for SEQ

35

ID NO: 2-3, 10, 12-13 or 15 or with pBGal vector by the FuGENE-6 transfection reagent (Boehringer). In summary, 4  $\mu$ l of FuGENE-6 is diluted in 100  $\mu$ l of DMEM and incubated for 5 minutes. Then, this is added to 1  $\mu$ g of DNA and incubated for 15 minutes before adding it to a 35 mm dish of CHO cells. The CHO cells are incubated at 37°C with 5% CO<sub>2</sub>. After  
5 24 hours, media and cell lysates are collected, centrifuged and dialyzed against assay buffer (15 mM Tris pH 7.6, 134 mM NaCl, 5 mM glucose, 3 mM CaCl<sub>2</sub> and MgCl<sub>2</sub>).

B. Expression Study Using SEQ ID NO: 2-3, 10, 12-13 or 15

The expression of SEQ ID NO: 2-3, 10, 12-13 or 15 in various tissues is analyzed  
10 using a semi-quantitative polymerase chain reaction-based technique. Human cDNA libraries are used as sources of expressed genes from tissues of interest (adult bladder, adult brain, adult heart, adult kidney, adult lymph node, adult liver, adult lung, adult ovary, adult placenta, adult rectum, adult spleen, adult testis, bone marrow, thymus, thyroid gland, fetal kidney, fetal liver, fetal liver-spleen, fetal skin, fetal brain, fetal leukocyte and macrophage). Gene-specific  
15 primers are used to amplify portions of the SEQ ID NO: 2-3, 10, 12-13 or 15 sequence from the samples. Amplified products are separated on an agarose gel, transferred and chemically linked to a nylon filter. The filter is then hybridized with a radioactively labeled (<sup>33</sup>P-dCTP) double-stranded probe generated from SEQ ID NO: 2-3, 10, 12-13 or 15 using a Klenow polymerase, random-prime method. The filters are washed (high stringency) and used to  
20 expose a phosphorimaging screen for several hours. Bands indicate the presence of cDNA including SEQ ID NO: 2-3, 10, 12-13 or 15 sequences in a specific library, and thus mRNA expression in the corresponding cell type or tissue.

## CLAIMS

## WE CLAIM:

1. An isolated polynucleotide comprising a nucleotide sequence selected from  
5 the group consisting of SEQ ID NO: 2-3, 10, 12-13 or 15, the translated protein coding portion thereof, the mature protein coding portion thereof, the extracellular portion thereof, or the active domain thereof.
2. An isolated polynucleotide encoding a polypeptide with biological activity,  
10 which polynucleotide hybridizes to the complement of a polynucleotide of claim 1 under stringent hybridization conditions.
3. An isolated polynucleotide encoding a polypeptide with biological activity,  
said polynucleotide having greater than about 90% sequence identity with the polynucleotide of  
15 claim 1.
4. The polynucleotide of claim 1 which is a DNA sequence.
5. An isolated polynucleotide which comprises the complement of the  
20 polynucleotide of claim 1.
6. A vector comprising the polynucleotide of claim 1.
7. An expression vector comprising the polynucleotide of claim 1.  
25
8. A host cell genetically engineered to express the polynucleotide of claim  
1.  
1.
9. The host cell of claim 8 wherein the polynucleotide is in operative  
30 association with a regulatory sequence that controls expression of the polynucleotide in the host cell.
10. An isolated polypeptide comprising an amino acid sequence which is at  
least 80% identical to the amino acid sequence selected from the group consisting of SEQ ID NO:

4-9, 14, or 16-18, the translated protein coding portion thereof, the mature protein coding portion thereof, the extracellular portion thereof, or the active domain thereof.

- 5                   11.     A composition comprising the polypeptide of claim 10 and a carrier.
12.     A polypeptide, having prothrombinase-like activity, comprising at least ten consecutive amino acids from the polypeptide sequences selected from the group consisting of SEQ ID NO: 4-9, 14, or 16-18.
- 10               13.     The polypeptide of claim 12, comprising at least five consecutive amino acids from the polypeptide sequences selected from the group consisting of SEQ ID NO. 4-9, 14, or 16-18.
14.     A polynucleotide encoding a polypeptide according to claim 12.
- 15               15.     A polynucleotide encoding a polypeptide according to claim 13.
16.     A polynucleotide encoding a polypeptide according to claim 10.
- 20               17.     An antibody specific for the polypeptide of claim 10.
18.     A method for detecting the polynucleotide of claim 1 in a sample, comprising:  
                  a)     contacting the sample with a compound that binds to and forms a complex  
25     with the polynucleotide of claim 1 for a period sufficient to form the complex; and  
                  b)     detecting the complex, so that if a complex is detected, the polynucleotide of claim 1 is detected.
19.     A method for detecting the polynucleotide of claim 1 in a sample, comprising:  
30     comprising:  
                  a)     contacting the sample under stringent hybridization conditions with nucleic acid primers that anneal to the polynucleotide of claim 1 under such conditions;

- b) amplifying a product comprising at least a portion of the polynucleotide of claim 1; and
- c) detecting said product and thereby the polynucleotide of claim 1 in the sample.

5

20. The method of claim 19, wherein the polynucleotide comprises an RNA molecule and the method further comprises reverse transcribing an annealed RNA molecule into a cDNA polynucleotide.

10

21. A method for detecting the polypeptide of claim 10 in a sample, comprising:

- a) contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex; and

15

- b) detecting formation of the complex, so that if a complex formation is detected, the polypeptide of claim 10 is detected.

22. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:

20

- a) contacting the compound with the polypeptide of claim 10 under conditions and for a time sufficient to form a polypeptide/compound complex; and
- b) detecting the complex, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

25

23. A method for identifying a compound that binds to the polypeptide of claim 10, comprising:

- a) contacting the compound with the polypeptide of claim 10, in a cell, for a time sufficient to form a polypeptide/compound complex, wherein the complex drives expression of a reporter gene sequence in the cell; and

30

- b) detecting the complex by detecting reporter gene sequence expression, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 10 is identified.

24. A method of producing a prothrombinase-like polypeptide, comprising,

a) culturing the host cell of claim 8 under conditions sufficient to express the polypeptide in said cell; and

b) isolating the polypeptide from the cell culture or cells of step (a).

5 25. A kit comprising the polypeptide of claim 10.

26. A nucleic acid array comprising the polynucleotide of claim 1 or a unique segment of the polynucleotide of claim 1 attached to a surface.

10 27. The array of claim 26, wherein the array detects full-matches to the polynucleotide or a unique segment of the polynucleotide of claim 1.

28. The array of claim 26, wherein the array detects mismatches to the polynucleotide or a unique segment of the polynucleotide of claim 1.

15

29. A method of treatment of a subject in need of enhanced activity or expression of prothrombinase-like polypeptide of claim 10 comprising administering to the subject a composition selected from the group consisting of:

20 a) a therapeutic amount of an agonist of said polypeptide;  
b) a therapeutic amount of the polypeptide; and  
c) a therapeutic amount of a polynucleotide encoding the polypeptide in a form and under conditions such that the polypeptide is produced, and a pharmaceutically acceptable carrier.

25 30. A method of treatment of a subject having need to inhibit activity or expression of prothrombinase-like polypeptide of claim 10 comprising administering to the subject a composition selected from the group consisting of:

a) a therapeutic amount of an antagonist to said polypeptide;  
b) a therapeutic amount of a polynucleotide that inhibits the expression of  
30 the nucleotide sequence encoding said polypeptide; and  
c) a therapeutic amount of a polypeptide that competes with the prothrombinase-like polypeptide for its ligand, and a pharmaceutically acceptable carrier.



31. An isolated polynucleotide with human prothrombinase activity comprising a nucleotide sequence of SEQ ID NO: 2 or 3, the mature protein coding portion thereof, or the active domain thereof.
- 5           32. An isolated polynucleotide encoding a polypeptide with biological activity, said polynucleotide which hybridizes to the complement of a polynucleotide of claim 31 under stringent hybridization conditions.
- 10           33. An isolated polynucleotide encoding a polypeptide with biological activity, said polynucleotide having greater than about 90% sequence identity with the polynucleotide of claim 31.
34. The polynucleotide of claim 31 which is a DNA sequence.
- 15           35. An isolated polynucleotide which comprises the complement of the polynucleotide of claim 31.
36. A vector comprising the polynucleotide of claim 31.
- 20           37. An expression vector comprising the polynucleotide of claim 31.
38. A host cell genetically engineered to express the polynucleotide of claim 31.
- 25           39. A host cell genetically engineered to contain the polynucleotide of claim 31 in operative association with a regulatory sequence that controls expression of the polynucleotide in the host cell.
- 30           40. An isolated polypeptide comprising an amino acid selected from the group consisting of SEQ ID NO: 4-9, the mature protein portion thereof, and active domain thereof.
41. A composition comprising the polypeptide of claim 40 and a carrier.
42. An antibody directed against the polypeptide of claim 40.

43. A method for detecting the polynucleotide of claim 31 in a sample, comprising:

- a) contacting the sample with a compound that binds to and forms a complex with the polynucleotide of claim 31 for a period sufficient to form the complex; and
- b) detecting the complex, so that if a complex is detected, the polynucleotide of claim 31 is detected.

44. A method for detecting the polynucleotide of claim 31 in a sample, comprising:

- a) contacting the sample under stringent hybridization conditions with nucleic acid primers that anneal to the polynucleotide of claim 31 under such conditions;
- b) amplifying a product comprising at least a portion of the polynucleotide of claim 31; and
- c) detecting said product and thereby the polynucleotide of claim 31 in the sample.

45. The method of claim 44, wherein the polynucleotide is an RNA molecule that encodes a polypeptide of claim 40, and the method further comprises reverse transcribing an annealed RNA molecule into a cDNA polynucleotide.

46. A method for detecting the polypeptide of claim 40 in a sample, comprising:

- a) contacting the sample with a compound that binds to and forms a complex with the polypeptide under conditions and for a period sufficient to form the complex; and
- b) detecting formation of the complex, so that if a complex formation is detected, the polypeptide of claim 40 is detected.

30

47. A method for identifying a compound that binds to the polypeptide of claim 40, comprising:

- a) contacting the compound with the polypeptide of claim 40 under conditions and for a time sufficient to form a polypeptide/compound complex; and

b) detecting the complex, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 40 is identified.

48. A method for identifying a compound that binds to the polypeptide of claim 40, comprising:

- a) contacting the compound with the polypeptide of claim 40, in a cell, for a time sufficient to form a polypeptide/compound complex, wherein the complex drives expression of a reporter gene sequence in the cell; and
- b) detecting the complex by detecting reporter gene sequence expression, so that if the polypeptide/compound complex is detected, a compound that binds to the polypeptide of claim 40 is identified.

49. A method of producing the polypeptide of claim 40, comprising,

- a) culturing the host cell of claim 38 for a period of time sufficient to express the polypeptide in said cell; and
- b) isolating the polypeptide from the cell culture or cells of step (a).

50. A kit comprising the polypeptide of claim 40.

51. A nucleic acid array comprising the polynucleotide of claim 31 or a unique segment of the polynucleotide of claim 31 attached to a surface.

52. The array of claim 51, wherein the array detects full-matches to the polynucleotide or a unique segment of the polynucleotide of claim 31.

53. The array of claim 51, wherein the array detects mismatches to the polynucleotide or a unique segment of the polynucleotide of claim 31.

54. A method of treatment comprising administering to a mammalian subject in need thereof a therapeutic amount of a composition comprising a polypeptide of claim 10 and a pharmaceutically acceptable carrier.

55. A polypeptide having prothrombinase activity comprising at least ten consecutive amino acids from the group consisting of SEQ ID NO: 4-9.

56. The polypeptide of claim 55, comprising at least five consecutive amino acids from the group consisting of SEQ ID NO: 4-9.

5 57. A polynucleotide encoding a polypeptide according to claim 55.

58. A polynucleotide encoding a polypeptide according to claim 56.

59. A polynucleotide encoding a polypeptide according to claim 40.

10

60. An isolated polypeptide with prothrombinase activity having at least an amino acid sequence of SEQ ID NO: 9, wherein A=Alanine, C=Cysteine, D=Aspartic Acid, E= Glutamic Acid, F=Phenylalanine, G=Glycine, H=Histidine, I=Isoleucine, K=Lysine, L=Leucine, M=Methionine, N=Asparagine, P=Proline, Q=Glutamine, R=Arginine,  
15 S=Serine, T=Threonine, V=Valine, W=Tryptophan, Y=Tyrosine.

# FIG. 1-1

BLASTX ALIGNMENT OF SEQ ID NO: 4 (PROTHROMBINASE-LIKE WITH HUMAN PROTHROMBINASE FG 12 PROTEIN (PROTHROMBINASE Fg1)

The Genbank database indicates that the fibrinogen-like protein is homologous (37% identity)

```
>gi |5730075 ref |NP_006673.1 |fibrinogen-like 2 precursor; fibroblast
>pir || S47273 pT49 protein-human >pir ||I37391 fibrinogen-like
protein expressed in T lymphocytes (pT49) -- human
>emb|CAA85298.1 |(Z36531) fibrinogen-like protein [Homo sapiens]
>gb|AAD10825.1 |(AF104015) fibrinogen-like protein 2 [Homo sapiens]
Length = 439
```

Score = 402 (141.5 bits), Expect = 5.7e-36, P = 5.7e-36  
Identities = 109/291 (37%), Positives = 155/291 (53%)

```
PROTHROMBINASE-LIKE: 98 L RNMMEQQASLDYLSNQVNELMNRVLLTTEVFRKQLDPFPHRPVQSHGL--DCTDIKD 155
+ N +D + A+L ++ VN L + ++ +Q+ RPVQ H + DC+D
PROTHROMBINASE FG12: 169 IENYVDSKVANLTFV----VNSLDGKCKSCPSQ----EQIQS----RPVQ--HL IYKDCSDYY- 217
```

TO FIG. 1-2

# FIG. 1-2

FROM FIG. 1-1

PROTHROMBINASE-LIKE: 156 TIGSVTKTPSGLYIHPGSSYPFEVMCDMDYRGGGWTVIQKRIDGIDFQRLWCDYLDG 215  
 IG K S Y + P+ + FEV CDM+ GGGWTV+Q R+DG +F R W DY G  
 PROTHROMBINASE FG12: 218 AIG---KRSSETYRVTDPKNSSEFVYCDMETMGGGWTVLQARLDGSTNFTRTWQDYKAG 274

PROTHROMBINASE-LIKE: 216 FGDLLGEFWLGLKKIFYIVNQNTSFMLYVALESEDDTLAYASYDNFWLEDETRFFKMHL 275  
 FG+L EFWLG KI + K +L + LE + YA YD F++ +E +++H+  
 PROTHROMBINASE FG12: 275 FGNLRREFWLGNDKIHLLTKSKE---MILRIDLEDFNGVELYALYDQFYVANEFLKYRLHV 332

PROTHROMBINASE-LIKE: 276 GRYSGNAGDAFRGLKKEDNQNAMPFSTSDVDNDGCRPACLVNGQSVKSCSHLHNKTGWWF 335  
 G Y+G AGDA R K N + F+T D DND P+ G +C L+ +GWWF  
 PROTHROMBINASE FG12: 333 GNYNGTAGDALP-FNKHYNHDLKFFTPDKDNDRY-PS-----G-----NCG-LYYSSGWWF 381

PROTHROMBINASE-LIKE: 336 NECGLANLNG-IHHFSGKLLATGIQWGTW---TKNNSPVKIKSVSMKIRRMYPN-YFK 388  
 + C ANLNG +H + + GI WGTW P KS + + M P +FK  
 PROTHROMBINASE FG12: 382 DACLSANLNGKYHQYRGVRNGIFWGTWPGVSEAHPPGGYKSSFKEAKMMIRPKHFK 438

# FIG. 2

BLASTP ALIGNMENT OF SEQ ID NO: 14 PROTHROMBINASE-LIKE POLYPEPTIDE (IDENTIFIED AS PROTHROMBINASE-LIKE) WITH HUMAN PROTHROMBINASE FGL2 PROTEIN SEQ ID NO: 19

Query: Prothrombinase-like polypeptide (SEQ ID NO: 14)  
 Subject: Human prothrombinase Fg12 protein (SEQ ID NO: 19)  
 sp | W88235 | W88235 Human prothrombinase Fg12 protein.  
 Length = 439

Score = 175 (61.6 bits), Expect = 4.0e-18, Sum P (2) = 4.0e-18  
 Identities = 49/132 (37%), Positives = 72/132 (54%)

Query: 345 LRMMDEQQASLDYLSNQVNELMNRVLLLTTEVFRKQLDPFPHRPVQSHGL--DCTDIKD 518  
 + N +D + A+L ++ VN L + ++ +Q+ RPVQ H + DC+D  
 Sbjct: 169 IENYVDSKVANLTFV---VNSLDGKCKSKCPSQ---EQIQS---RPVQ-HLIYKDCSDYY- 217

Query: 519 TIGSVTKTPSGLYIHPEGSSYPFEVMCDMDYRGGWTVIQKRIDGIDFQRLWCDYLDG 698  
 IG K S Y + P+ + FEV CDM+ GGGWTV+Q R+DG +F R W DY G  
 Sbjct: 218 AIG---KRSSETYRVTPDPKNSSFEVYCDMETMGGWTVLQARLDGGSTNFTRTWQDYKAG 274

Query: 699 FGDLLGDAFRGLKK 740  
 FG+L + + G K  
 Sbjct: 275 FGNLRREFWLGNDK 288

FIG.3

BLASTP ALIGNMENT OF SEQ ID NO: 14 PROTHROMBINASE--LIKE POLYPEPTIDE (IDENTIFIED AS PROTHROMBINASE--LIKE) WITH HUMAN PROTHROMBINASE FGL2 PROTEIN SEQ ID NO: 19

Query: Prothrombinase--like polypeptide (SEQ ID NO: 14)  
Subject: Human prothrombinase Fg12 protein (SEQ ID NO: 19)

sp | W88235 | W88235 Human prothrombinase Fg12 protein.  
Length = 439

Score = 129 (45.4 bits), Expect = 2.0e-18, Sum P (2) = 2.0e-18  
Identities = 40/115 (34%), Positives = 56/115 (48%)

Query: 687	YLDGFGDLLGDAFRGLKKEDNQNAMPFSTSDVDNDGCRPACLVNGQSVKSCSHLHNKTGW	866
	++ + GDA R K N + F+T D DND P+ G +C L+ +GW	
Sbjct: 331	HVGNYNGTAGDALR-FNKHYNHDLKFFTTDPKDNDRY-PS-----G-----NCG-LYYSSGW	379
Query: 867	WFNECGLANLG--IHHFSGKLLATGIQWGTW---TKNNSPVKIKSVSMKIRRMYPN-YFK	1031
	WF+ C ANLNG +H + + GI WGTW P KS + + M P +FK	
Sbjct: 380	WFDACLSANLNGKYYHQYRGVRNGIFWGTWPGVSEAHPPGGYKSSFKEAKMMIRPKHFK	438







## SEQUENCE LISTING

<110> HYSEQ, Inc.  
Haley, Dana A  
Boyle, Bryan J  
Ho, Alice S  
Zhou, Ping  
Mize, Nancy K  
Kuo, Chiauyun  
Arterburn, Matthew C  
Tang, Y. Tom  
Liu, Chenghua  
Drmanac, Radoje T

<120> Methods and Materials Relating to Prothrombinase-Like  
Polypeptides and Polynucleotides

<130> Prothrombinase-Like

<140> not yet assigned  
<141> 2000-12-22

<150> 09/643,313  
<151> 2000-08-21

<150> 09/596,196  
<151> 2000-06-17

<150> 09/522,317  
<151> 2000-04-25

<150> 09/491,404  
<151> 2000-01-25

<150> 09/488,725  
<151> 2000-01-21

<160> 22

<170> PatentIn Ver. 2.1

<210> 1  
<211> 725  
<212> DNA  
<213> Homo sapiens

<400> 1  
ctgatcttaa ggataccatt ggctctgtca ccaaaacacc gagtgggtta tacataattc 60  
accagaagg atctagctac ccatctgagg taatgtgtga catggattac agaggagggtg 120  
gatggactgt gatacagaaa agaattgatg ggataattga tttccagagg ttgtgggtgtg 180  
attatctgga tggatttggg gatcttctag gagaattttg gctaggactg aaaaagattt 240  
tttatatagt aaatcacaaa aataccagtt ttatgctgta tgtggctttg gaatctgaag 300  
atgacactct tgccttatgca tcatatgata atttttggct agaggatgaa acgagatttt 360  
ttaaaatgca cttaggacgg tattcaggaa atgctggtga tgcattccgg ggtctcaaaa 420  
aagaagataa tcaaaatgca atgcctttta gcacatcaga tgttgataat gatgggtgtc 480  
gccctgcatg cctgggtcaat ggtcaatctg tgaagagctg gagtcacctt cataacaaga 540  
cccggctgtg gtttaacgag tgtgggctaa ccaatctaaa tggcattcat cacttctctt 600  
ggaaaatggc ttgcaactgg aattcaatgg gggcacgtgg gaccacaaac cacttaacct 660  
gtccaaggaa ttaaatctgg gttccactgg aaaattagga agaaatgtgc caattccata 720  
ttttt 725

<210> 2.

&lt;211&gt; 1069

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

```

ctgatcttaa ggataccatt ggctctgtca ccaaaacacc gagtgggtta tacataattc 60
accagaagg atctagctac ccatttgagg taatgtgtga catggattac agaggaggtg 120
gatggactgt gatacagaaa agaattgatg ggataattga ttccagagg ttgtggtgtg 180
attatctgga tggatttggg gatcttctag gagaattttg gctaggactg aaaaagattt 240
tttatatagt aaatcagaaa aataccagtt ttatgctgta tgtggctttg gaatctgaag 300
atgacactct tgcctatgca tcatatgata atttttggct agaggatgaa acgagatttt 360
ttaaagtga cttaggacgg tattcaggaa atgctggtga tgcattccgg ggtctcaaaa 420
aagaagataa tcaaaatgca atgcctttta gcacatcaga tgttgataat gatgggtgtc 480
gccctgcatg cctgggtcaat ggtcagtctg tgaagagctg cagtcacctc cataacaaga 540
ccggctgggtg gtttaacgag tgtggtctag caaatotaaa tggcattcat cacttctctg 600
gaaaattgct tgcaactgga attcaatggg gcacgtggac caaaaacaac tcacctgtca 660
agattaaatc tgtttcaatg aaaattagaa gaatgtacaa tccatatttt aaataatctc 720
atttaacatt gtaatgcaag ttctacaatg ataatatatt aaagattttt aaaagtttat 780
cttttcactt agtgtttcaa acatattagg caaaatttaa ctgtagatgg catttagatg 840
ttatgagttt aattagaaaa ctccaatttt gtagtattct ataaaagaaa acatggctta 900
ttgtatgttt ttacttctga ctatattaac aatatacaat gaaatttggt tcaagtgaac 960
tacaacttgt ctctcctaaaa tttatagtga ttttaaagga ttttgctttt tctttgaagc 1020
atttttaaac cataatatgt tgtaaggaaa attgaaggga atattttac 1069

```

&lt;210&gt; 3

&lt;211&gt; 2013

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1)..(1167)

&lt;400&gt; 3

```

atg atg tct cca tcc caa gcc tca ctc tta ttc tta aat gta tgt att 48
Met Met Ser Pro Ser Gln Ala Ser Leu Leu Phe Leu Asn Val Cys Ile
1 5 10 15

ttt att tgt gga gaa gtt gta caa ggt aac tgt gta cat cat tct acg 96
Phe Ile Cys Gly Glu Val Val Gln Gly Asn Cys Val His His Ser Thr
20 25 30

gac tct tca gta gtt aac att gta gaa gat gga tct aat gca aaa gat 144
Asp Ser Ser Val Val Asn Ile Val Glu Asp Gly Ser Asn Ala Lys Asp
35 40 45

gaa agt aaa agt aat gat act gtt tgt aag gaa gac tgt gag gaa tca 192
Glu Ser Lys Ser Asn Asp Thr Val Cys Lys Glu Asp Cys Glu Glu Ser
50 55 60

tgt gat gtt aaa act aaa att aca cga gaa gaa aaa cat ttc atg tgt 240
Cys Asp Val Lys Thr Lys Ile Thr Arg Glu Glu Lys His Phe Met Cys
65 70 75 80

aga aat ttg caa aat tct att gtt tcc tac aca aga agt acc aaa aaa 288
Arg Asn Leu Gln Asn Ser Ile Val Ser Tyr Thr Arg Ser Thr Lys Lys
85 90 95

cta cta agg aat atg atg gat gag caa caa gct tcc ttg gat tat tta 336
Leu Leu Arg Asn Met Met Asp Glu Gln Gln Ala Ser Leu Asp Tyr Leu
100 105 110

```

tct aat cag gtt aac gag ctc atg aat aga gtt ctc ctt ttg act aca	384
Ser Asn Gln Val Asn Glu Leu Met Asn Arg Val Leu Leu Leu Thr Thr	
115 120 125	
gaa gtt ttt aga aaa cag ctg gat cct ttt cct cac aga cct gtt cag	432
Glu Val Phe Arg Lys Gln Leu Asp Pro Phe Pro His Arg Pro Val Gln	
130 135 140	
tca cat ggt tta gat tgc act gat att aag gat acc att ggc tct gtc	480
Ser His Gly Leu Asp Cys Thr Asp Ile Lys Asp Thr Ile Gly Ser Val	
145 150 155 160	
acc aaa aca ccg agt ggt tta tac ata att cac cca gaa gga tct agc	528
Thr Lys Thr Pro Ser Gly Leu Tyr Ile Ile His Pro Glu Gly Ser Ser	
165 170 175	
tac cca ttt gag gta atg tgt gac atg gat tac aga gga ggt gga tgg	576
Tyr Pro Phe Glu Val Met Cys Asp Met Asp Tyr Arg Gly Gly Gly Trp	
180 185 190	
act gtg ata cag aaa aga att gat ggg ata att gat ttc cag agg ttg	624
Thr Val Ile Gln Lys Arg Ile Asp Gly Ile Ile Asp Phe Gln Arg Leu	
195 200 205	
tgg tgt gat tat ctg gat gga ttt gga gat ctt cta gga gaa ttt tgg	672
Trp Cys Asp Tyr Leu Asp Gly Phe Gly Asp Leu Leu Gly Glu Phe Trp	
210 215 220	
cta gga ctg aaa aag att ttt tat ata gta aat cag aaa aat acc agt	720
Leu Gly Leu Lys Lys Ile Phe Tyr Ile Val Asn Gln Lys Asn Thr Ser	
225 230 235 240	
ttt atg ctg tat gtg gct ttg gaa tct gaa gat gac act ctt gct tat	768
Phe Met Leu Tyr Val Ala Leu Glu Ser Glu Asp Asp Thr Leu Ala Tyr	
245 250 255	
gca tca tat gat aat ttt tgg cta gag gat gaa acg aga ttt ttt aaa	816
Ala Ser Tyr Asp Asn Phe Trp Leu Glu Asp Glu Thr Arg Phe Phe Lys	
260 265 270	
atg cac tta gga cgg tat tca gga aat gct ggt gat gca ttc cgg ggt	864
Met His Leu Gly Arg Tyr Ser Gly Asn Ala Gly Asp Ala Phe Arg Gly	
275 280 285	
ctc aaa aaa gaa gat aat caa aat gca atg cct ttt agc aca tca gat	912
Leu Lys Lys Glu Asp Asn Gln Asn Ala Met Pro Phe Ser Thr Ser Asp	
290 295 300	
gtt gat aat gat ggg tgt cgc cct gca tgc ctg gtc aat ggt cag tct	960
Val Asp Asn Asp Gly Cys Arg Pro Ala Cys Leu Val Asn Gly Gln Ser	
305 310 315 320	
gtg aag agc tgc agt cac ctc cat aac aag acc ggc tgg tgg ttt aac	1008
Val Lys Ser Cys Ser His Leu His Asn Lys Thr Gly Trp Trp Phe Asn	
325 330 335	
gag tgt ggt cta gca aat cta aat ggc att cat cac ttc tct gga aaa	1056
Glu Cys Gly Leu Ala Asn Leu Asn Gly Ile His His Phe Ser Gly Lys	
340 345 350	
ttg ctt gca act gga att caa tgg ggc acg tgg acc aaa aac aac tca	1104
Leu Leu Ala Thr Gly Ile Gln Trp Gly Thr Trp Thr Lys Asn Asn Ser	
355 360 365	

cct gtc aag att aaa tct gtt tca atg aaa att aga aga atg tac aat 1152  
 Pro Val Lys Ile Lys Ser Val Ser Met Lys Ile Arg Arg Met Tyr Asn  
 370 375 380

cca tat ttt aaa taa tctcatttaa cattgtaatg caagttctac aatgataata 1207  
 Pro Tyr Phe Lys  
 385

tattaaagat ttttaaaagt ttatcttttc acttagtggt tcaaacatat taggcaaaat 1267

ttaactgtag atggcattta gatgttatga gttaattag aaaacttcaa tttttagta 1327

ttctataaaa gaaacatgg cttattgtat gtttttactt ctgactatat taacaatata 1387

caatgaaatt tgtttcaagt gaactacaac ttgtcttcct aaaatttata gtgattttta 1447

aggattttgc cttttctttg aagcattttt aaaccataat atgttgtaag gaaaattgaa 1507

gggaatattt tacttatttt tatactttat atgattatat aatctacaga taatttctac 1567

tgaagacagt tacaataaat aactttatgc agattaatat ataagctaca catgatgtaa 1627

aaaccttact atttctaggt gatgccatac catttttaaaa gtagtaagag tttgctgccc 1687

aaatagtttt tcttgttttc atatctaate atggttaact attttgttat tgtttgtaat 1747

aaatatatgt acttttatat cctgaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1807

aaaaaaaaagg ggcgcccctt taattttaaag gggcccttta aaccccggtt aaaccccggt 1867

aggggggcttt tatagggccc ccccttggtg tgtgcccccc ccccggggtt ttttttgacg 1927

ggggggggggc cccccccact tttttttttt taaaaaaaag agaaaattcc ccccttttgt 1987

gggaagggggg gttttttttt ttggggg 2013

<210> 4

<211> 388

<212> PRT

<213> Homo sapiens

<400> 4

Met	Met	Ser	Pro	Ser	Gln	Ala	Ser	Leu	Leu	Phe	Leu	Asn	Val	Cys	Ile
1				5				10					15		
Phe	Ile	Cys	Gly	Glu	Val	Val	Gln	Gly	Asn	Cys	Val	His	His	Ser	Thr
		20						25				30			
Asp	Ser	Ser	Val	Val	Asn	Ile	Val	Glu	Asp	Gly	Ser	Asn	Ala	Lys	Asp
		35					40					45			
Glu	Ser	Lys	Ser	Asn	Asp	Thr	Val	Cys	Lys	Glu	Asp	Cys	Glu	Glu	Ser
		50				55				60					
Cys	Asp	Val	Lys	Thr	Lys	Ile	Thr	Arg	Glu	Glu	Lys	His	Phe	Met	Cys
		65			70				75			80			
Arg	Asn	Leu	Gln	Asn	Ser	Ile	Val	Ser	Tyr	Thr	Arg	Ser	Thr	Lys	Lys
			85					90				95			
Leu	Leu	Arg	Asn	Met	Met	Asp	Glu	Gln	Gln	Ala	Ser	Leu	Asp	Tyr	Leu
			100				105					110			
Ser	Asn	Gln	Val	Asn	Glu	Leu	Met	Asn	Arg	Val	Leu	Leu	Leu	Thr	Thr
		115				120					125				
Glu	Val	Phe	Arg	Lys	Gln	Leu	Asp	Pro	Phe	Pro	His	Arg	Pro	Val	Gln
		130			135						140				
Ser	His	Gly	Leu	Asp	Cys	Thr	Asp	Ile	Lys	Asp	Thr	Ile	Gly	Ser	Val

```

145          150          155          160
Thr Lys Thr Pro Ser Gly Leu Tyr Ile Ile His Pro Glu Gly Ser Ser
          165          170          175
Tyr Pro Phe Glu Val Met Cys Asp Met Asp Tyr Arg Gly Gly Gly Trp
          180          185          190
Thr Val Ile Gln Lys Arg Ile Asp Gly Ile Ile Asp Phe Gln Arg Leu
          195          200          205
Trp Cys Asp Tyr Leu Asp Gly Phe Gly Asp Leu Leu Gly Glu Phe Trp
          210          215          220
Leu Gly Leu Lys Lys Ile Phe Tyr Ile Val Asn Gln Lys Asn Thr Ser
225          230          235          240
Phe Met Leu Tyr Val Ala Leu Glu Ser Glu Asp Asp Thr Leu Ala Tyr
          245          250          255
Ala Ser Tyr Asp Asn Phe Trp Leu Glu Asp Glu Thr Arg Phe Phe Lys
          260          265          270
Met His Leu Gly Arg Tyr Ser Gly Asn Ala Gly Asp Ala Phe Arg Gly
          275          280          285
Leu Lys Lys Glu Asp Asn Gln Asn Ala Met Pro Phe Ser Thr Ser Asp
290          295          300
Val Asp Asn Asp Gly Cys Arg Pro Ala Cys Leu Val Asn Gly Gln Ser
305          310          315          320
Val Lys Ser Cys Ser His Leu His Asn Lys Thr Gly Trp Trp Phe Asn
          325          330          335
Glu Cys Gly Leu Ala Asn Leu Asn Gly Ile His His Phe Ser Gly Lys
          340          345          350
Leu Leu Ala Thr Gly Ile Gln Trp Gly Thr Trp Thr Lys Asn Asn Ser
          355          360          365
Pro Val Lys Ile Lys Ser Val Ser Met Lys Ile Arg Arg Met Tyr Asn
          370          375          380
Pro Tyr Phe Lys
385

```

```

<210> 5
<211> 37
<212> PRT
<213> Homo sapiens

```

```

<400> 5
Val Met Cys Asp Met Asp Tyr Arg Gly Gly Gly Trp Thr Val Ile Gln
  1              5              10              15

Lys Arg Ile Asp Gly Ile Ile Asp Phe Gln Arg Leu Trp Cys Asp Tyr
      20              25              30

Leu Asp Gly Phe Gly
      35

```

```

<210> 6
<211> 30
<212> PRT
<213> Homo sapiens

```

```

<400> 6
Cys Ser His Leu His Asn Lys Thr Gly Trp Trp Phe Asn Glu Cys Gly
  1              5              10              15

Leu Ala Asn Leu Asn Gly Ile His His Phe Ser Gly Lys Leu
      20              25              30

```

<210> 7  
 <211> 15  
 <212> PRT  
 <213> Homo sapiens

<400> 7  
 Phe Lys Met His Leu Gly Arg Tyr Ser Gly Asn Ala Gly Asp Ala  
     1                    5                    10                    15

<210> 8  
 <211> 13  
 <212> PRT  
 <213> Homo sapiens

<400> 8  
 Glu Phe Trp Leu Gly Leu Lys Lys Ile Phe Tyr Ile Val  
     1                    5                    10

<210> 9  
 <211> 368  
 <212> PRT  
 <213> Homo sapiens

<400> 9  
 Glu Val Val Gln Gly Asn Cys Val His His Ser Thr Asp Ser Ser Val  
     1                    5                    10                    15  
 Val Asn Ile Val Glu Asp Gly Ser Asn Ala Lys Asp Glu Ser Lys Ser  
                     20                    25                    30  
 Asn Asp Thr Val Cys Lys Glu Asp Cys Glu Glu Ser Cys Asp Val Lys  
                     35                    40                    45  
 Thr Lys Ile Thr Arg Glu Glu Lys His Phe Met Cys Arg Asn Leu Gln  
                     50                    55                    60  
 Asn Ser Ile Val Ser Tyr Thr Arg Ser Thr Lys Lys Leu Leu Arg Asn  
     65                    70                    75                    80  
 Met Met Asp Glu Gln Gln Ala Ser Leu Asp Tyr Leu Ser Asn Gln Val  
                     85                    90                    95  
 Asn Glu Leu Met Asn Arg Val Leu Leu Leu Thr Thr Glu Val Phe Arg  
                     100                    105                    110  
 Lys Gln Leu Asp Pro Phe Pro His Arg Pro Val Gln Ser His Gly Leu  
                     115                    120                    125  
 Asp Cys Thr Asp Ile Lys Asp Thr Ile Gly Ser Val Thr Lys Thr Pro  
     130                    135                    140  
 Ser Gly Leu Tyr Ile Ile His Pro Glu Gly Ser Ser Tyr Pro Phe Glu  
     145                    150                    155                    160  
 Val Met Cys Asp Met Asp Tyr Arg Gly Gly Gly Trp Thr Val Ile Gln  
                     165                    170                    175  
 Lys Arg Ile Asp Gly Ile Ile Asp Phe Gln Arg Leu Trp Cys Asp Tyr  
                     180                    185                    190  
 Leu Asp Gly Phe Gly Asp Leu Leu Gly Glu Phe Trp Leu Gly Leu Lys



195	200	205
Lys Ile Phe Tyr Ile Val	Asn Gln Lys Asn Thr	Ser Phe Met Leu Tyr
210	215	220
Val Ala Leu Glu Ser Glu	Asp Asp Thr Leu Ala	Tyr Ala Ser Tyr Asp
225	230	235 240
Asn Phe Trp Leu Glu Asp	Glu Thr Arg Phe Phe	Lys Met His Leu Gly
	245 250	255
Arg Tyr Ser Gly Asn Ala	Gly Asp Ala Phe Arg	Gly Leu Lys Lys Glu
	260 265	270
Asp Asn Gln Asn Ala Met	Pro Phe Ser Thr Ser	Asp Val Asp Asn Asp
	275 280	285
Gly Cys Arg Pro Ala Cys	Leu Val Asn Gly Gln	Ser Val Lys Ser Cys
	290 295	300
Ser His Leu His Asn Lys	Thr Gly Trp Trp Phe	Asn Glu Cys Gly Leu
	310 315	320
Ala Asn Leu Asn Gly Ile	His His Phe Ser Gly	Lys Leu Leu Ala Thr
	325 330	335
Gly Ile Gln Trp Gly Thr	Trp Thr Lys Asn Asn	Ser Pro Val Lys Ile
	340 345	350
Lys Ser Val Ser Met Lys	Ile Arg Arg Met Tyr	Asn Pro Tyr Phe Lys
	355 360	365

<210> 10  
 <211> 2352  
 <212> DNA  
 <213> Homo sapiens

<400> 10  
 ccagacgagc tacctaagaa cagctaaaag agcacacccg tctatgtagc aaaatagtg 60  
 gaagatttat aggtagaggc gacaaaccta ccgagcctgg tgatagctgg ttgtccaaga 120  
 tagaatctta gttcaacttt aaatttgccc acagaaccct cttaatcccc ttgtaaattt 180  
 aactgttagt ccaagagga acagctcttt ggacactagg aaaaaacctt gtagagagag 240  
 taaaaaattt aacaccata gtaggcctaa aagcagccac caaaaaaaa ggggcctgac 300  
 tgatatttga agaagtgttt tcatctatcc aagaaaaata tgatgtctcc atccaagcc 360  
 tcactcttat tcttaaatgt atgtattttt atttgtggag aagttgtaca aggttaactgt 420  
 gtacatcatt ctacggactc ttcagtagtt aacattgtag aagatggatc taatgcaaaa 480  
 gatgaaagta aaagtaatga tactgtttgt aaggaagact gtgaggaatc atgtgatgtt 540  
 aaaactaaaa ttacacgaga agaaaaacat ttcattgtga gaaatttgca aaattctatt 600  
 gtttcctaca caagaagtac caaaaaacta ctaaggaata tgatggatga gcaacaagct 660  
 tccttggtgt attatctgga tcagggttaac gagctcatga atagagttct ccttttgact 720  
 acagaagttt ttagaaaaca gctggatcct tttcctcaca gacctgttca gtcacatggg 780  
 ttagattgca ctgatattaa ggataccatt ggctctgtca ccaaaccacc gagtggttta 840  
 tacataattc acccagaagg atctagctac ccatttgagg taatgtgtga catggattac 900  
 agaggaggtg gatggactgt gatacagaaa agaattgatg ggataattga tttccagagg 960  
 ttgtggtgtg attatctgga tggatttgga gatcttctag gagaattttg gctaggactg 1020  
 aaaaagattt tttatatagt aaatcagaaa aataccagtt ttatgctgta tgtggctttg 1080  
 gaatctgaag atgacactct tgcttatgca tcatatgata atttttggct agaggatgaa 1140  
 acgagatttt ttaaaatgca cttaggacgg tattcaggaa atgctggtga tgcattccgg 1200  
 ggtctcaaaa aagaagataa tcaaaatgca atgcctttta gcacatcaga tgttgataat 1260

```

gatgggtgtc gccctgcatg cctgggtcaat ggtcagtcctg tgaagagctg cagtcacctc 1320
cataacaaga ccggctgggtg gtttaacgag tgtgggtctag caaatctaaa tggcattcat 1380
cacttctctg gaaaattgct tgcaactgga attcaatggg gcacgtggac caaaaacaac 1440
tcacctgtca agattaaatc tgtttcaatg aaaattagaa gaatgtacaa tccatatttt 1500
aaataatctc atttaacatt gtaatgcaag ttctacaatg ataatatatt aaagattttt 1560
aaaagtttat cttttcactt agtggtttcaa acatattagg caaaatttaa ctgtagatgg 1620
catttagatg ttatgagttt aattagaaaa cttcaatttt gtagtattct ataaaagaaa 1680
acatggctta ttgtatgttt ttacttctga ctatattaac aatatacaat gaaatttggt 1740
tcaagtgaac tacaacttgt cttcctaaaa tttatagtga ttttaaagga ttttgcttt 1800
tctttgaagc atttttaaac cataatatgt tgtaaggaaa attgaaggga atattttact 1860
tatttttata ctttatatga ttatataato tacagataat ttctactgaa gacagtata 1920
ataaataact ttatgcagat taatatataa gctacacatg atgtaaaaac cttactattt 1980
ctaggtgatg ccataccatt ttaaaagtag taagagtttg ctgccccaat agtttttctt 2040
gttttcatat ctaatcatgg ttaactattt tgttattggt tgtaataaat atatgtactt 2100
ttatatcctg aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaggggcg 2160
ccctttaa cttaaaggggc cctttaaac ccgggtaaac ccccgaggg ggcttttata 2220
ggggccccc ttggtgtgtg ccccccccc ggggtttttt ttgacggggg gggggccccc 2280
cccactttt tttttttaa aaaaagagaa aattccccc tttgtggga aggggggttt 2340
tttttttgg gg 2352

```

<210> 11  
 <211> 453  
 <212> DNA  
 <213> Homo sapiens

```

<400> 11
taatcgang agcccgatct notgatattt gcngnccggt tttcatctat ccaagaaaaa 60
tatgatgtct ccattccctag ccttactctt attcttaaat gtatgtattt ttgtttgtgg 120
agaagctgta caaggtaact gtgtacatca ttctacggac tcttcagtag ttaacattgt 180
agaagatgga tctaatgcaa aagatgaaag taaaagtaat gatactgttt gtaaggaaga 240
ctgtgaggaa tcatgtgatg ttaaaactaa aattacacga gaagaaaaac attcatgtg 300
tagaaaatttg caaaattcta ttgtttccta cacaagaagt accaaaaaac tactaaggaa 360
tatgatggat gagcaacaag cttccttggg ttatttatct aatcaggtaa tgtgtgacat 420
ggattacaga ggaggtggat ggactgtgat aca 453

```

<210> 12  
 <211> 1648  
 <212> DNA  
 <213> Homo sapiens

```

<400> 12
ggtcgaccca cgcgtccgat ttgaagaagt gttttcatct atccaagaaa aatatgatgt 60
ctccatccca agcctcactc ttattcttaa atgtatgtat ttttatttgt ggagaagttg 120
tacaaggtaa ctgtgtacat cattctacgg actcttcagt agttaacatt gtagaagatg 180
gatctaattg aaaaagatgaa agtaaaagta atgatactgt ttgtaaggaa gactgtgagg 240
aatcatgtga tgttaaaact aaaattacac gagaagaaaa acatttcatg tgtagaattt 300
tgcaaaattc tattgtttcc tacacaagaa gtaccaaaaa actactaagg aatatgatgg 360
atgagcaaca agcttccttg gattattttat ctaatcagggt taacgagctc atgaatagag 420
ttctcctttt gactacagaa gtttttagaa aacagctgga tccttttcct cacagacctg 480
ttcagtcaca tgggttagat tgcaactgata ttaaggatac cattgggtct gtcacaaaaa 540
caccgagtggt tttatataca attcaccagc aaggatctag ctaccatttt gaggtaattg 600
gtgacatgga ttacagagga ggtggatgga ctgtgatata gaaaagaatt gatgggataa 660
ttgatttcca gaggttgtgg ttgtgattatc tggatggatt tggagatctt ctagggtgatg 720
cattccgggg tctcaaaaaa gaagataatc aaaatgcaat gccttttagc acatcagatg 780
ttgataatga tgggtgtcgc cctgcatgcc tgggtcaatgg tcagtctgtg aagagctgca 840
gtcacctcca taacaagacc ggctgggtggt ttaacgagtg tgggtctagca aatctaaatg 900
gcattcatca cttctctgga aaattgcttg caactgggat tcaatggggc acgtggacca 960
aaaacaactc acctgtcaag attaaacttg tttcaatgaa aattagaaga atgtacaatc 1020
catattttta ataactctat ttaacattgt aatgcaagtt ctacaatgat aatatattaa 1080
agatttttaa aagtttatct tttcacttag tgtttcaaac atattaggca aaatttaact 1140
gtagatggca ttttagatgt atgagtttaa ttagaaaaact tcaattttgt agtattctat 1200

```

```

aaaagaaaac atggccttatt gtatgttttt acttctgact atattaacaa tataacaatga 1260
aatttgtttc aagtgaacta caacttgtct tcctaaaatt tatagtgatt ttaaaggatt 1320
ttgccttttc tttgaagcat ttttaaacca taatatgttg taaggaaaat tgaaggggaat 1380
attttactta tttttatact ttatatgatt atataatcta cagataatct ctactgaaga 1440
cagttacaat aaataacttt atgcagatta atataatagc tacacatgat gtaaaaacct 1500
tactatttct aggtgatgcc ataccatttt aaaagtagta agagtttgct gcccaaatag 1560
tttttcttgt tttcatatct aatcatgggt aactattttg ttattgtttg taataaatat 1620
atgtactttt atatcctgaa aaaaaaaaa 1648

```

&lt;210&gt; 13

&lt;211&gt; 1651

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (54)..(1034)

&lt;400&gt; 13

```

ggctgaccca cgcgtccgat ttgaagaagt gttttcatct atccaagaaa aat atg      56
                                         Met
                                         1

atg tct cca tcc caa gcc tca ctc tta ttc tta aat gta tgt att ttt      104
Met Ser Pro Ser Gln Ala Ser Leu Leu Phe Leu Asn Val Cys Ile Phe
                    5                      10                      15

att tgt gga gaa gtt gta caa ggt aac tgt gta cat cat tct acg gac      152
Ile Cys Gly Glu Val Val Gln Gly Asn Cys Val His His Ser Thr Asp
                    20                      25                      30

tct tca gta gtt aac att gta gaa gat gga tct aat gca aaa gat gaa      200
Ser Ser Val Val Asn Ile Val Glu Asp Gly Ser Asn Ala Lys Asp Glu
                    35                      40                      45

agt aaa agt aat gat act gtt tgt aag gaa gac tgt gag gaa tca tgt      248
Ser Lys Ser Asn Asp Thr Val Cys Lys Glu Asp Cys Glu Glu Ser Cys
                    50                      55                      60                      65

gat gtt aaa act aaa att aca cga gaa gaa aaa cat ttc atg tgt aga      296
Asp Val Lys Thr Lys Ile Thr Arg Glu Glu Lys His Phe Met Cys Arg
                    70                      75                      80

aat ttg caa aat tct att gtt tcc tac aca aga agt acc aaa aaa cta      344
Asn Leu Gln Asn Ser Ile Val Ser Tyr Thr Arg Ser Thr Lys Lys Leu
                    85                      90                      95

cta agg aat atg atg gat gag caa caa gct tcc ttg gat tat tta tct      392
Leu Arg Asn Met Met Asp Glu Gln Gln Ala Ser Leu Asp Tyr Leu Ser
                    100                     105                     110

aat cag gtt aac gag ctc atg aat aga gtt ctc ctt ttg act aca gaa      440
Asn Gln Val Asn Glu Leu Met Asn Arg Val Leu Leu Leu Thr Thr Glu
                    115                     120                     125

gtt ttt aga aaa cag ctg gat cct ttt cct cac aga cct gtt cag tca      488
Val Phe Arg Lys Gln Leu Asp Pro Phe Pro His Arg Pro Val Gln Ser
                    130                     135                     140                     145

cat ggt tta gat tgc act gat att aag gat acc att ggc tct gtc acc      536
His Gly Leu Asp Cys Thr Asp Ile Lys Asp Thr Ile Gly Ser Val Thr
                    150                     155                     160

```

aaa aca ccg agt ggt tta tac ata att cac cca gaa gga tct agc tac 584  
 Lys Thr Pro Ser Gly Leu Tyr Ile Ile His Pro Glu Gly Ser Ser Tyr  
 165 170 175

cca ttt gag gta atg tgt gac atg gat tac aga gga ggt gga tgg act 632  
 Pro Phe Glu Val Met Cys Asp Met Asp Tyr Arg Gly Gly Gly Trp Thr  
 180 185 190

gtg ata cag aaa aga att gat ggg ata att gat ttc cag agg ttg tgg 680  
 Val Ile Gln Lys Arg Ile Asp Gly Ile Ile Asp Phe Gln Arg Leu Trp  
 195 200 205

tgt gat tat ctg gat gga ttt gga gat ctt cta ggt gat gca ttc cgg 728  
 Cys Asp Tyr Leu Asp Gly Phe Gly Asp Leu Leu Gly Asp Ala Phe Arg  
 210 215 220 225

ggt ctc aaa aaa gaa gat aat caa aat gca atg cct ttt agc aca tca 776  
 Gly Leu Lys Lys Glu Asp Asn Gln Asn Ala Met Pro Phe Ser Thr Ser  
 230 235 240

gat gtt gat aat gat ggg tgt cgc cct gca tgc ctg gtc aat ggt cag 824  
 Asp Val Asp Asn Asp Gly Cys Arg Pro Ala Cys Leu Val Asn Gly Gln  
 245 250 255

tct gtg aag agc tgc agt cac ctc cat aac aag acc ggc tgg tgg ttt 872  
 Ser Val Lys Ser Cys Ser His Leu His Asn Lys Thr Gly Trp Trp Phe  
 260 265 270

aac gag tgt ggt cta gca aat cta aat ggc att cat cac ttc tct gga 920  
 Asn Glu Cys Gly Leu Ala Asn Leu Asn Gly Ile His His Phe Ser Gly  
 275 280 285

aaa ttg ctt gca act gga att caa tgg ggc acg tgg acc aaa aac aac 968  
 Lys Leu Leu Ala Thr Gly Ile Gln Trp Gly Thr Trp Thr Lys Asn Asn  
 290 295 300 305

tca cct gtc aag att aaa tct gtt tca atg aaa att aga aga atg tac 1016  
 Ser Pro Val Lys Ile Lys Ser Val Ser Met Lys Ile Arg Arg Met Tyr  
 310 315 320

aat cca tat ttt aaa taa tctcatttaa cattgtaatg caagttctac 1064  
 Asn Pro Tyr Phe Lys  
 325

aatgataata tattaagat ttttaaaagt ttatcttttc acttagtggt tcaaacatat 1124  
 taggcaaaat ttaactgtag atggcattta gatggtatga gtttaattag aaaacttcaa 1184  
 ttttgtagta ttctataaaa gaaaacatgg cttattgtat gtttttactt ctgactatat 1244  
 taacaatata caatgaaatt tgtttcaagt gaactacaac ttgtcttcct aaaatttata 1304  
 gtgattttta aggattttgc cttttctttg aagcattttt aaaccataat atgttgtaag 1364  
 gaaaattgaa gggaatattt tacttatttt tatactttat atgattatat aatctacaga 1424  
 taattttctac tgaagacagt tacaataaat aactttatgc agattaatat ataagctaca 1484  
 catgatgtaa aaaccttact atttctaggt gatgccatac cattttaaaa gtagtaagag 1544  
 tttgctgccc aaatagtttt tcttgttttc atatctaate atggttaact attttggtat 1604

tgtttgtaat aaatatatgt actttttaaaa aaaacagaaa aaaaaaa

1651

<210> 14  
 <211> 326  
 <212> PRT  
 <213> Homo sapiens

<400> 14  
 Met Met Ser Pro Ser Gln Ala Ser Leu Leu Phe Leu Asn Val Cys Ile  
 1 5 10 15  
 Phe Ile Cys Gly Glu Val Val Gln Gly Asn Cys Val His His Ser Thr  
 20 25 30  
 Asp Ser Ser Val Val Asn Ile Val Glu Asp Gly Ser Asn Ala Lys Asp  
 35 40 45  
 Glu Ser Lys Ser Asn Asp Thr Val Cys Lys Glu Asp Cys Glu Glu Ser  
 50 55 60  
 Cys Asp Val Lys Thr Lys Ile Thr Arg Glu Glu Lys His Phe Met Cys  
 65 70 75 80  
 Arg Asn Leu Gln Asn Ser Ile Val Ser Tyr Thr Arg Ser Thr Lys Lys  
 85 90 95  
 Leu Leu Arg Asn Met Met Asp Glu Gln Gln Ala Ser Leu Asp Tyr Leu  
 100 105 110  
 Ser Asn Gln Val Asn Glu Leu Met Asn Arg Val Leu Leu Leu Thr Thr  
 115 120 125  
 Glu Val Phe Arg Lys Gln Leu Asp Pro Phe Pro His Arg Pro Val Gln  
 130 135 140  
 Ser His Gly Leu Asp Cys Thr Asp Ile Lys Asp Thr Ile Gly Ser Val  
 145 150 155 160  
 Thr Lys Thr Pro Ser Gly Leu Tyr Ile Ile His Pro Glu Gly Ser Ser  
 165 170 175  
 Tyr Pro Phe Glu Val Met Cys Asp Met Asp Tyr Arg Gly Gly Gly Trp  
 180 185 190  
 Thr Val Ile Gln Lys Arg Ile Asp Gly Ile Ile Asp Phe Gln Arg Leu  
 195 200 205  
 Trp Cys Asp Tyr Leu Asp Gly Phe Gly Asp Leu Leu Gly Asp Ala Phe  
 210 215 220  
 Arg Gly Leu Lys Lys Glu Asp Asn Gln Asn Ala Met Pro Phe Ser Thr  
 225 230 235 240  
 Ser Asp Val Asp Asn Asp Gly Cys Arg Pro Ala Cys Leu Val Asn Gly  
 245 250 255  
 Gln Ser Val Lys Ser Cys Ser His Leu His Asn Lys Thr Gly Trp Trp  
 260 265 270  
 Phe Asn Glu Cys Gly Leu Ala Asn Leu Asn Gly Ile His His Phe Ser  
 275 280 285  
 Gly Lys Leu Leu Ala Thr Gly Ile Gln Trp Gly Thr Trp Thr Lys Asn  
 290 295 300  
 Asn Ser Pro Val Lys Ile Lys Ser Val Ser Met Lys Ile Arg Arg Met  
 305 310 315 320  
 Tyr Asn Pro Tyr Phe Lys  
 325

<210> 15  
 <211> 981  
 <212> DNA  
 <213> Homo sapiens

<400> 15  
 atgatgtctc catcccaagc ctcaactotta ttcttaaatg tatgtatttt tatttggtgga 60  
 gaagttgtac aaggtaactg tgtacatcat tctacggact cttcagtagt taacattgta 120  
 gaagatggat ctaatgcaaa agatgaaagt aaaagtaatg atactgtttg taaggaagac 180

```

tgtgaggaat catgtgatgt taaaactaaa attacacgag aagaaaaaca tttcatgtgt 240
agaaatttgc aaaattctat tgtttctctac acaagaagta ccaaaaaact actaaggaaat 300
atgatggatg agcaacaagc ttccttggat tatttatcta atcagggttaa cgagctcatg 360
aatagagttc tccttttgac tacagaagtt tttagaaaac agctggatcc ttttctcac 420
agacctgttc agtcacatgg tttagattgc actgatatta aggataccat tggctctgtc 480
acaaaaacac cgagtggttt atacataatt caccagaag gatctagcta cccatttgag 540
gtaatgtgtg acatggatta cagaggaggt ggatggactg tgatacagaa aagaattgat 600
gggataattg atttccagag gttgtggtgt gattatctgg atggatttgg agatcttcta 660
ggtgatgcat tccggggtct caaaaaagaa gataatcaaa atgcaatgcc ttttagcaca 720
tcagatgttg ataattgatg gtgtcgccct gcatgcctgg tcaatggtca gtctgtgaag 780
agctgcagtc acctccataa caagaccggc tgggtggtta acgagtgtgg tctagcaa 840
ctaaatggca ttcacactt ctctggaaaa ttgcttgcaa ctggaattca atggggcacg 900
tggacaaaaa acaactcacc tgtcaagatt aaatctgtt caatgaaaat tagaagaatg 960
tacaatccat attttaata a 981

```

```

<210> 16
<211> 20
<212> PRT
<213> Homo sapiens

```

```

<400> 16
Met Met Ser Pro Ser Gln Ala Ser Leu Leu Phe Leu Asn Val Cys Ile
  1              5              10              15

Phe Ile Cys Gly
              20

```

```

<210> 17
<211> 240
<212> PRT
<213> Homo sapiens

```

```

<400> 17
Asn Leu Gln Asn Ser Ile Val Ser Tyr Thr Arg Ser Thr Lys Lys Leu
  1              5              10              15

Leu Arg Asn Met Met Asp Glu Gln Gln Ala Ser Leu Asp Tyr Leu Ser
              20              25              30

Asn Gln Val Asn Glu Leu Met Asn Arg Val Leu Leu Leu Thr Thr Glu
              35              40              45

Val Phe Arg Lys Gln Leu Asp Pro Phe Pro His Arg Pro Val Gln Ser
              50              55              60

His Gly Leu Asp Cys Thr Asp Ile Lys Asp Thr Ile Gly Ser Val Thr
              65              70              75              80

Lys Thr Pro Ser Gly Leu Tyr Ile Ile His Pro Glu Gly Ser Ser Tyr
              85              90              95

Pro Phe Glu Val Met Cys Asp Met Asp Tyr Arg Gly Gly Gly Trp Thr
              100              105              110

Val Ile Gln Lys Arg Ile Asp Gly Ile Ile Asp Phe Gln Arg Leu Trp
              115              120              125

Cys Asp Tyr Leu Asp Gly Phe Gly Asp Leu Leu Gly Asp Ala Phe Arg
              130              135              140

Gly Leu Lys Lys Glu Asp Asn Gln Asn Ala Met Pro Phe Ser Thr Ser

```

```

145              150              155              160
Asp Val Asp Asn Asp Gly Cys Arg Pro Ala Cys Leu Val Asn Gly Gln
      165              170              175

Ser Val Lys Ser Cys Ser His Leu His Asn Lys Thr Gly Trp Trp Phe
      180              185              190

Asn Glu Cys Gly Leu Ala Asn Leu Asn Gly Ile His His Phe Ser Gly
      195              200              205

Lys Leu Leu Ala Thr Gly Ile Gln Trp Gly Thr Trp Thr Lys Asn Asn
      210              215              220

Ser Pro Val Lys Ile Lys Ser Val Ser Met Lys Ile Arg Arg Met Tyr
      225              230              235              240

```

<210> 18  
 <211> 306  
 <212> PRT  
 <213> Homo sapiens

```

<400> 18
Glu Val Val Gln Gly Asn Cys Val His His Ser Thr Asp Ser Ser Val
  1              5              10              15

Val Asn Ile Val Glu Asp Gly Ser Asn Ala Lys Asp Glu Ser Lys Ser
      20              25              30

Asn Asp Thr Val Cys Lys Glu Asp Cys Glu Glu Ser Cys Asp Val Lys
      35              40              45

Thr Lys Ile Thr Arg Glu Glu Lys His Phe Met Cys Arg Asn Leu Gln
      50              55              60

Asn Ser Ile Val Ser Tyr Thr Arg Ser Thr Lys Lys Leu Leu Arg Asn
      65              70              75              80

Met Met Asp Glu Gln Gln Ala Ser Leu Asp Tyr Leu Ser Asn Gln Val
      85              90              95

Asn Glu Leu Met Asn Arg Val Leu Leu Leu Thr Thr Glu Val Phe Arg
      100              105              110

Lys Gln Leu Asp Pro Phe Pro His Arg Pro Val Gln Ser His Gly Leu
      115              120              125

Asp Cys Thr Asp Ile Lys Asp Thr Ile Gly Ser Val Thr Lys Thr Pro
      130              135              140

Ser Gly Leu Tyr Ile Ile His Pro Glu Gly Ser Ser Tyr Pro Phe Glu
      145              150              155              160

Val Met Cys Asp Met Asp Tyr Arg Gly Gly Gly Trp Thr Val Ile Gln
      165              170              175

Lys Arg Ile Asp Gly Ile Ile Asp Phe Gln Arg Leu Trp Cys Asp Tyr
      180              185              190

```

Leu Asp Gly Phe Gly Asp Leu Leu Gly Asp Ala Phe Arg Gly Leu Lys  
           195                                  200                                  205  
 Lys Glu Asp Asn Gln Asn Ala Met Pro Phe Ser Thr Ser Asp Val Asp  
           210                                  215                                  220  
 Asn Asp Gly Cys Arg Pro Ala Cys Leu Val Asn Gly Gln Ser Val Lys  
           225                                  230                                  235                                  240  
 Ser Cys Ser His Leu His Asn Lys Thr Gly Trp Trp Phe Asn Glu Cys  
                                   245                                  250                                  255  
 Gly Leu Ala Asn Leu Asn Gly Ile His His Phe Ser Gly Lys Leu Leu  
                                   260                                  265                                  270  
 Ala Thr Gly Ile Gln Trp Gly Thr Trp Thr Lys Asn Asn Ser Pro Val  
                                   275                                  280                                  285  
 Lys Ile Lys Ser Val Ser Met Lys Ile Arg Arg Met Tyr Asn Pro Tyr  
           290                                  295                                  300  
 Phe Lys  
 305

<210> 19  
 <211> 270  
 <212> PRT  
 <213> Homo sapiens

<400> 19  
 Ile Glu Asn Tyr Val Asp Ser Lys Val Ala Asn Leu Thr Phe Val Val  
   1                                  5                                  10                                  15  
 Asn Ser Leu Asp Gly Lys Cys Ser Lys Cys Pro Ser Gln Glu Gln Ile  
                                   20                                  25                                  30  
 Gln Ser Arg Pro Val Gln His Leu Ile Tyr Lys Asp Cys Ser Asp Tyr  
                                   35                                  40                                  45  
 Tyr Ala Ile Gly Lys Arg Ser Ser Glu Thr Tyr Arg Val Thr Pro Asp  
           50                                  55                                  60  
 Pro Lys Asn Ser Ser Phe Glu Val Tyr Cys Asp Met Glu Thr Met Gly  
           65                                  70                                  75                                  80  
 Gly Gly Trp Thr Val Leu Gln Ala Arg Leu Asp Gly Ser Thr Asn Phe  
                                   85                                  90                                  95  
 Thr Arg Thr Trp Gln Asp Tyr Lys Ala Gly Phe Gly Asn Leu Arg Arg  
                                   100                                  105                                  110  
 Glu Phe Trp Leu Gly Asn Asp Lys Ile His Leu Leu Thr Lys Ser Lys  
           115                                  120                                  125  
 Glu Met Ile Leu Arg Ile Asp Leu Glu Asp Phe Asn Gly Val Glu Leu  
           130                                  135                                  140  
 Tyr Ala Leu Tyr Asp Gln Phe Tyr Val Ala Asn Glu Phe Leu Lys Tyr  
           145                                  150                                  155                                  160  
 Arg Leu His Val Gly Asn Tyr Asn Gly Thr Ala Gly Asp Ala Leu Arg  
                                   165                                  170                                  175



Phe Asn Lys His Tyr Asn His Asp Leu Lys Phe Phe Thr Thr Pro Asp  
 180 185 190  
 Lys Asp Asn Asp Arg Tyr Pro Ser Gly Asn Cys Gly Leu Tyr Tyr Ser  
 195 200 205  
 Ser Gly Trp Trp Phe Asp Ala Cys Leu Ser Ala Asn Leu Asn Gly Lys  
 210 215 220  
 Tyr Tyr His Gln Lys Tyr Arg Gly Val Arg Asn Gly Ile Phe Trp Gly  
 225 230 235 240  
 Thr Trp Pro Gly Val Ser Glu Ala His Pro Gly Gly Tyr Lys Ser Ser  
 245 250 255  
 Phe Lys Glu Ala Lys Met Met Ile Arg Pro Lys His Phe Lys  
 260 265 270

<210> 20  
 <211> 307  
 <212> PRT  
 <213> Homo sapiens

<400> 20  
 Tyr Ile Asp Glu Thr Val Asn Ser Asn Ile Pro Thr Asn Leu Arg Val  
 1 5 10 15  
 Leu Arg Ser Ile Leu Glu Asn Leu Arg Ser Lys Ile Gln Lys Leu Glu  
 20 25 30  
 Ser Asp Val Ser Ala Gln Met Glu Tyr Cys Arg Thr Pro Cys Thr Val  
 35 40 45  
 Ser Cys Asn Ile Pro Val Val Ser Gly Lys Glu Cys Glu Glu Ile Ile  
 50 55 60  
 Arg Lys Gly Gly Glu Thr Ser Glu Met Tyr Leu Ile Gln Pro Asp Ser  
 65 70 75 80  
 Ser Val Lys Pro Tyr Arg Val Tyr Cys Asp Met Asn Thr Glu Asn Gly  
 85 90 95  
 Gly Trp Thr Val Ile Gln Asn Arg Gln Asp Gly Ser Val Asp Phe Gly  
 100 105 110  
 Arg Lys Trp Asp Pro Tyr Lys Gln Gly Phe Gly Asn Val Ala Thr Asn  
 115 120 125  
 Thr Asp Gly Lys Asn Tyr Cys Gly Leu Pro Gly Glu Tyr Trp Leu Gly  
 130 135 140  
 Asn Asp Lys Ile Ser Gln Leu Thr Arg Met Gly Pro Thr Glu Leu Leu  
 145 150 155 160  
 Ile Glu Met Glu Asp Trp Lys Gly Asp Lys Val Lys Ala His Tyr Gly  
 165 170 175  
 Gly Phe Thr Val Gln Asn Glu Ala Asn Lys Tyr Gln Ile Ser Val Asn  
 180 185 190  
 Lys Tyr Arg Gly Thr Ala Gly Asn Ala Leu Met Asp Gly Ala Ser Gln

195                      200                      205  
 Leu Met Gly Glu Asn Arg Thr Met Thr Ile His Asn Gly Met Phe Phe  
     210                      215                      220  
 Ser Thr Tyr Asp Arg Asp Asn Asp Gly Trp Leu Thr Ser Asp Pro Arg  
     225                      230                      235                      240  
 Lys Gln Cys Ser Lys Glu Asp Gly Gly Gly Trp Trp Tyr Asn Arg Cys  
     245                      250                      255  
 His Ala Ala Asn Pro Asn Gly Arg Tyr Tyr Trp Gly Gly Gln Tyr Thr  
     260                      265                      270  
 Trp Asp Met Ala Lys His Gly Thr Asp Asp Gly Val Val Trp Met Asn  
     275                      280                      285  
 Trp Lys Gly Ser Trp Tyr Ser Met Arg Lys Met Ser Met Lys Ile Arg  
     290                      295                      300  
 Pro Phe Phe  
 305

<210> 21  
 <211> 237  
 <212> PRT  
 <213> Homo sapiens

<400> 21  
 Asp Leu Lys Asp Thr Ile Gly Ser Val Thr Lys Thr Pro Ser Gly Leu  
     1                      5                      10                      15  
 Tyr Ile Ile His Pro Glu Gly Ser Ser Tyr Pro Phe Glu Val Met Cys  
     20                      25                      30  
 Asp Met Asp Tyr Arg Gly Gly Gly Trp Thr Val Ile Gln Lys Arg Ile  
     35                      40                      45  
 Asp Gly Ile Ile Asp Phe Gln Arg Leu Trp Cys Asp Tyr Leu Asp Gly  
     50                      55                      60  
 Phe Gly Asp Leu Leu Gly Glu Phe Trp Leu Gly Leu Lys Lys Ile Phe  
     65                      70                      75                      80  
 Tyr Ile Val Asn Gln Lys Asn Thr Ser Phe Met Leu Tyr Val Ala Leu  
     85                      90                      95  
 Glu Ser Glu Asp Asp Thr Leu Ala Tyr Ala Ser Tyr Asp Asn Phe Trp  
     100                      105                      110  
 Leu Glu Asp Glu Thr Arg Phe Phe Lys Met His Leu Gly Arg Tyr Ser  
     115                      120                      125  
 Gly Asn Ala Gly Asp Ala Phe Arg Gly Leu Lys Lys Glu Asp Asn Gln  
     130                      135                      140  
 Asn Ala Met Pro Phe Ser Thr Ser Asp Val Asp Asn Asp Gly Cys Arg  
     145                      150                      155                      160  
 Pro Ala Cys Leu Val Asn Gly Gln Ser Val Lys Ser Cys Ser His Leu  
     165                      170                      175

His Asn Lys Thr Gly Trp Trp Phe Asn Glu Cys Gly Leu Ala Asn Leu  
                   180                  185                  190  
 Asn Gly Ile His His Phe Ser Gly Lys Leu Leu Ala Thr Gly Ile Gln  
                   195                  200                  205  
 Trp Gly Thr Trp Thr Lys Asn Asn Ser Pro Val Lys Ile Lys Ser Val  
                   210                  215                  220  
 Ser Met Lys Ile Arg Arg Met Tyr Asn Pro Tyr Lys Phe  
                   225                  230                  235

<210> 22  
 <211> 45  
 <212> PRT  
 <213> Homo sapiens

<400> 22  
 Met Met Ser Pro Ser Gln Ala Ser Leu Leu Phe Leu Asn Val Cys Ile  
           1                  5                  10                  15  
 Phe Ile Cys Gly Glu Val Val Gln Gly Asn Cys Val His His Ser Thr  
                   20                  25                  30  
 Asp Ser Ser Val Val Asn Ile Val Glu Asp Gly Ser Asn  
           35                  40                  45

1

1

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/36061

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(7) : Please See Extra Sheet. US CL : 435/69.1, 69.2, 183, 252.3, 320.1, 325, ; 530/350; 536/25.1 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/69.1, 69.2, 183, 252.3, 320.1, 325, ; 530/350; 536/25.1 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) STN (Bioscience), East (all databases), sequence search for SEQ ID NO:2, search terms: angiopoietin, prothrombinase, fg12.		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P — Y,P	Database GenEmbl on GenCore, Compugen Ltd. Acc. No. AC015500, BIRREN et al. Homo Sapiens, clone RP11-21G19, Direct Submission, March 2000.	1, 2, 4, 31-34 — 6-9, 24, 26, 36-39, 49
A,P	LEVY et al. Molecular and Functional Analysis of the Human Prothrombinase Gene (HFGL2) and its Role in Viral Hepatitis. Am. J. Path. April 2000. Vol. 156, No. 4, pp. 1217-1225.	1-4, 6-16, 24-26, 31-34, 36-41, 49-50, 55-60
A	GALE et al. Growth factors acting via endothelial cell-specific receptor tyrosine kinases: VEGFs, Angiopoietins, and ephrins in vascular development. Genes & Dev. 1999, Vol. 13, pp. 1055-1066.	1-4, 6-16, 24-26, 31-34, 36-41, 49-50, 55-60
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family	
Date of the actual completion of the international search 05 JULY 2001		Date of mailing of the international search report 07 AUG 2001
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-9230		Authorized officer HOLLY SCHNITZER Telephone No. (703) 305-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/35061

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Database GenEmbl on GenCore, Compugen Ltd. Acc. No. AX079973, BALLINGER et al. Novel angiopoietin materials and methods, WO 0105825, January 25, 2001, see especially title.	1-4, 6-16, 24-26, 31-34, 36-41, 49-50, 55-60

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/35061

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 27, 28, 45, 52, AND 53  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:  
1-4, 6-16, 24-25, 31-34, 36-41, 49, 50, 55-60 as they relate to SEQ ID NO:2

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US00/35061

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (7):

Co7H 21/02, 21/04; Co7K 14/00; C12N 1/21, 9/00, 15/00, 15/68, 15/85, 15/86

## BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-4, 6-16, 24-26, 31-34, 36-41, 49-50, and 55-60, drawn to a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 2-3, 10, 12-13, and 15, a recombinant vector, a host cell, a method of producing the encoded polypeptide, and a polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence selected from the group consisting of SEQ ID NO: 4-9, 14, and 16-18. Applicant must elect one nucleic acid sequence and one protein sequence (see species election below).

Group II, claim(s) 17 and 42, drawn to an antibody that selectively binds the claimed polypeptides.

Group III, claim(s) 18 and 43, drawn to a method of detecting a polynucleotide by hybridization assay.

Group IV, claims 19-20 and 44 drawn to a method of detecting a polynucleotide by PCR.

Group V, claims 21-22 and 46-47, drawn to a protein binding assay.

Group VI, claims 23 and 48, drawn to a method of identifying a compound that binds the claimed polypeptide by reporter gene expression.

Group VII, claims 1-6, 26, 31, 34-36, and 51, drawn to a nucleic acid array comprising the claimed polynucleotides.

Group VIII, claims 29a and c, drawn to a method of treatment by administering an agonist of the claimed polypeptides.

Group IX, claims 29b and 54, drawn to a method of treatment by administering the claimed polypeptide.

Group X, claim 29c, drawn to a method of gene therapy using the claimed polynucleotides.

Group XI, claim 30a, drawn to a method of treatment using an antagonist.

Group XII, claim 30b, drawn to a method of treatment using antisense therapy.

The inventions listed as Groups I-XII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Pursuant to 37 C.F.R. 1.475(d), the ISA/US considers that where multiple products and processes are claimed, the main invention shall consist of the first invention of the category first mentioned in the claims and the first invention of each of the other categories related thereto. Accordingly, the main invention (Group I) comprises the first recited product; the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO: 2-3, 10, 12-13, and 15, a recombinant vector, a host cell, a method of producing the encoded polypeptide, and a polypeptide comprising an amino acid sequence which is at least 80% identical to the amino acid sequence selected from the group consisting of SEQ ID NO: 4-9, 14, and 16-18. Further pursuant to 37 C.F.R. 1.475(d), the ISA/US considers that any feature which the subsequently recited products and methods share with the main invention does not constitute a special technical feature within the meaning of PCT Rule 13.2 and that each of such products and methods accordingly defines a separate invention.

This application contains claims directed to more than one species of the generic invention. These species are deemed to lack Unity of Invention because they are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for more than one species to be searched, the appropriate additional search fees must be paid. The species are as follows:

- I. The polynucleotide of SEQ ID NO: 2,
- II. The polynucleotide of SEQ ID NO: 3 and the polypeptide it encodes, SEQ ID NOs: 4 and 5-9.
- III. The polynucleotide of SEQ ID NO: 10,
- IV. The polynucleotide of SEQ ID NO: 12,
- V. The polynucleotide of SEQ ID NO: 13 and the polypeptide it encodes, SEQ ID NO: 14, 5-6, and 16-18,
- VI. The polynucleotide of SEQ ID NO: 15.

All of the claims are deemed to correspond to the species listed above.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/35061

The species listed above do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, the species lack the same or corresponding special technical features for the following reasons:

The polynucleotides with the nucleotide sequences set forth in SEQ ID NOs: 2-3, 10, 12-13, and 15 and the polypeptides they encode have different structural and functional features.

Therefore, Group I will be searched with respect to species I (the polynucleotide of SEQ ID NO:2). The appropriate fees must be paid for searches of additional groups and species.

In the absence of any response from the applicant, this Authority will establish the International Preliminary Examination Report based on the main invention. The claims drawn to the main invention are as follows:

Claims 1-4, 6-16, 24-25, 31-34, 36-41, 49-50, and 55-60 as they relate to SEQ ID NO: 2.